

PCT

WORLD INTELLECTUAL PROPERTY ORGANIZATION
International Bureau

INTERNATIONAL APPLICATION PUBLISHED UNDER THE PATENT COOPERATION TREATY (PCT)

(51) International Patent Classification ⁷ : C07K 14/47, C12N 5/10, 5/16, 15/12, 15/63, 15/64	A1	(11) International Publication Number: WO 00/55199 (43) International Publication Date: 21 September 2000 (21.09.00)
(21) International Application Number: PCT/US00/06014 (22) International Filing Date: 9 March 2000 (09.03.00) (30) Priority Data: 60/124,095 12 March 1999 (12.03.99) US 60/138,598 11 June 1999 (11.06.99) US 60/168,665 3 December 1999 (03.12.99) US (71) Applicant (for all designated States except US): HUMAN GENOME SCIENCES, INC. [US/US]; 9410 Key West Avenue, Rockville, MD 20850 (US). (72) Inventors; and (75) Inventors/Applicants (for US only): ROSEN, Craig, A. [US/US]; 22400 Rolling Hill Road, Laytonsville, MD 20882 (US). RUBEN, Steven, M. [US/US]; 18528 Heritage Hills Drive, Laytonsville, MD 20882 (US). KOMAT-SOULIS, George [US/US]; 9518 Garwood Street, Silver Spring, MD 20901 (US). (74) Agents: HOOVER, Kenley, K. et al.; Human Genome Sciences, Inc., 9410 Key West Avenue, Rockville, MD 20850 (US).		(81) Designated States: AL, AM, AT, AU, AZ, BA, BB, BG, BR, BY, CA, CH, CN, CU, CZ, DE, DK, EE, ES, FI, GB, GE, GH, GM, HR, HU, ID, IL, IN, IS, JP, KE, KG, KP, KR, KZ, LC, LK, LR, LS, LT, LU, LV, MD, MG, MK, MN, MW, MX, NO, NZ, PL, PT, RO, RU, SD, SE, SG, SI, SK, SL, TJ, TM, TR, TT, UA, UG, US, UZ, VN, YU, ZW, ARIPO patent (GH, GM, KE, LS, MW, SD, SL, SZ, TZ, UG, ZW), Eurasian patent (AM, AZ, BY, KG, KZ, MD, RU, TJ, TM), European patent (AT, BE, CH, CY, DE, DK, ES, FI, FR, GB, GR, IE, IT, LU, MC, NL, PT, SE), OAPI patent (BF, BJ, CF, CG, CI, CM, GA, GN, GW, ML, MR, NE, SN, TD, TG). Published <i>With international search report.</i> <i>Before the expiration of the time limit for amending the claims and to be republished in the event of the receipt of amendments.</i>
(54) Title: 47 HUMAN SECRETED PROTEINS (57) Abstract <p>The present invention relates to novel human secreted proteins and isolated nucleic acids containing the coding regions of the genes encoding such proteins. Also provided are vectors, host cells, antibodies, and recombinant methods for producing human secreted proteins. The invention further relates to diagnostic and therapeutic methods useful for diagnosing and treating disorders related to these novel human secreted proteins.</p>		

FOR THE PURPOSES OF INFORMATION ONLY

Codes used to identify States party to the PCT on the front pages of pamphlets publishing international applications under the PCT.

AL	Albania	ES	Spain	LS	Lesotho	SI	Slovenia
AM	Armenia	FI	Finland	LT	Lithuania	SK	Slovakia
AT	Austria	FR	France	LU	Luxembourg	SN	Senegal
AU	Australia	GA	Gabon	LV	Latvia	SZ	Swaziland
AZ	Azerbaijan	GB	United Kingdom	MC	Monaco	TD	Chad
BA	Bosnia and Herzegovina	GE	Georgia	MD	Republic of Moldova	TG	Togo
BB	Barbados	GH	Ghana	MG	Madagascar	TJ	Tajikistan
BE	Belgium	GN	Guinea	MK	The former Yugoslav Republic of Macedonia	TM	Turkmenistan
BF	Burkina Faso	GR	Greece	ML	Mali	TR	Turkey
BG	Bulgaria	HU	Hungary	MN	Mongolia	TT	Trinidad and Tobago
BJ	Benin	IE	Ireland	MR	Mauritania	UA	Ukraine
BR	Brazil	IL	Israel	MW	Malawi	UG	Uganda
BY	Belarus	IS	Iceland	MX	Mexico	US	United States of America
CA	Canada	IT	Italy	NE	Niger	UZ	Uzbekistan
CF	Central African Republic	JP	Japan	NL	Netherlands	VN	Viet Nam
CG	Congo	KE	Kenya	NO	Norway	YU	Yugoslavia
CH	Switzerland	KG	Kyrgyzstan	NZ	New Zealand	ZW	Zimbabwe
CI	Côte d'Ivoire	KP	Democratic People's Republic of Korea	PL	Poland		
CM	Cameroon	KR	Republic of Korea	PT	Portugal		
CN	China	KZ	Kazakhstan	RO	Romania		
CU	Cuba	LC	Saint Lucia	RU	Russian Federation		
CZ	Czech Republic	LI	Liechtenstein	SD	Sudan		
DE	Germany	LK	Sri Lanka	SE	Sweden		
DK	Denmark	LR	Liberia	SG	Singapore		
EE	Estonia						

47 Human Secreted Proteins

Field of the Invention

This invention relates to newly identified polynucleotides and the polypeptides encoded by these polynucleotides, uses of such polynucleotides and polypeptides, and their production.

Background of the Invention

Unlike bacterium, which exist as a single compartment surrounded by a membrane, human cells and other eucaryotes are subdivided by membranes into many functionally distinct compartments. Each membrane-bounded compartment, or organelle, contains different proteins essential for the function of the organelle. The cell uses "sorting signals," which are amino acid motifs located within the protein, to target proteins to particular cellular organelles.

One type of sorting signal, called a signal sequence, a signal peptide, or a leader sequence, directs a class of proteins to an organelle called the endoplasmic reticulum (ER). The ER separates the membrane-bounded proteins from all other types of proteins. Once localized to the ER, both groups of proteins can be further directed to another organelle called the Golgi apparatus. Here, the Golgi distributes the proteins to vesicles, including secretory vesicles, the cell membrane, lysosomes, and the other organelles.

Proteins targeted to the ER by a signal sequence can be released into the extracellular space as a secreted protein. For example, vesicles containing secreted proteins can fuse with the cell membrane and release their contents into the extracellular space - a process called exocytosis. Exocytosis can occur constitutively or after receipt of a triggering signal. In the latter case, the proteins are stored in secretory vesicles (or secretory granules) until exocytosis is triggered. Similarly, proteins residing on the cell membrane can also be secreted into the extracellular space by proteolytic cleavage of a "linker" holding the protein to the membrane.

Despite the great progress made in recent years, only a small number of genes encoding human secreted proteins have been identified. These secreted proteins include the commercially valuable human insulin, interferon, Factor VIII, human

growth hormone, tissue plasminogen activator, and erythropoietin. Thus, in light of the pervasive role of secreted proteins in human physiology, a need exists for identifying and characterizing novel human secreted proteins and the genes that encode them. This knowledge will allow one to detect, to treat, and to prevent
5 medical disorders by using secreted proteins or the genes that encode them.

Summary of the Invention

The present invention relates to novel polynucleotides and the encoded polypeptides. Moreover, the present invention relates to vectors, host cells,
10 antibodies, and recombinant and synthetic methods for producing the polypeptides and polynucleotides. Also provided are diagnostic methods for detecting disorders and conditions related to the polypeptides and polynucleotides, and therapeutic methods for treating such disorders and conditions. The invention further relates to screening methods for identifying binding partners of the polypeptides.

15

Detailed Description

Definitions

The following definitions are provided to facilitate understanding of certain terms used throughout this specification.

20 In the present invention, "isolated" refers to material removed from its original environment (e.g., the natural environment if it is naturally occurring), and thus is altered "by the hand of man" from its natural state. For example, an isolated polynucleotide could be part of a vector or a composition of matter, or could be contained within a cell, and still be "isolated" because that vector, composition of
25 matter, or particular cell is not the original environment of the polynucleotide. The term "isolated" does not refer to genomic or cDNA libraries, whole cell total or mRNA preparations, genomic DNA preparations (including those separated by electrophoresis and transferred onto blots), sheared whole cell genomic DNA preparations or other compositions where the art demonstrates no distinguishing
30 features of the polynucleotide/sequences of the present invention.

In the present invention, a "secreted" protein refers to those proteins capable of being directed to the ER, secretory vesicles, or the extracellular space as a result of a signal sequence, as well as those proteins released into the extracellular space without necessarily containing a signal sequence. If the secreted protein is released
5 into the extracellular space, the secreted protein can undergo extracellular processing to produce a "mature" protein. Release into the extracellular space can occur by many mechanisms, including exocytosis and proteolytic cleavage.

In specific embodiments, the polynucleotides of the invention are at least 15, at least 30, at least 50, at least 100, at least 125, at least 500, or at least 1000
10 continuous nucleotides but are less than or equal to 300 kb, 200 kb, 100 kb, 50 kb, 15 kb, 10 kb, 7.5 kb, 5 kb, 2.5 kb, 2.0 kb, or 1 kb, in length. In a further embodiment, polynucleotides of the invention comprise a portion of the coding sequences, as disclosed herein, but do not comprise all or a portion of any intron. In another embodiment, the polynucleotides comprising coding sequences do not contain coding
15 sequences of a genomic flanking gene (i.e., 5' or 3' to the gene of interest in the genome). In other embodiments, the polynucleotides of the invention do not contain the coding sequence of more than 1000, 500, 250, 100, 50, 25, 20, 15, 10, 5, 4, 3, 2, or 1 genomic flanking gene(s).

As used herein, a "polynucleotide" refers to a molecule having a nucleic acid
20 sequence contained in SEQ ID NO:X or the cDNA contained within the clone deposited with the ATCC. For example, the polynucleotide can contain the nucleotide sequence of the full length cDNA sequence, including the 5' and 3' untranslated sequences, the coding region, with or without the signal sequence, the secreted protein coding region, as well as fragments, epitopes, domains, and variants
25 of the nucleic acid sequence. Moreover, as used herein, a "polypeptide" refers to a molecule having the translated amino acid sequence generated from the polynucleotide as broadly defined.

In the present invention, the full length sequence identified as SEQ ID NO:X was often generated by overlapping sequences contained in multiple clones (contig
30 analysis). A representative clone containing all or most of the sequence for SEQ ID NO:X was deposited with the American Type Culture Collection ("ATCC"). As

shown in Table 1, each clone is identified by a cDNA Clone ID (Identifier) and the ATCC Deposit Number. The ATCC is located at 10801 University Boulevard, Manassas, Virginia 20110-2209, USA. The ATCC deposit was made pursuant to the terms of the Budapest Treaty on the international recognition of the deposit of microorganisms for purposes of patent procedure.

A "polynucleotide" of the present invention also includes those polynucleotides capable of hybridizing, under stringent hybridization conditions, to sequences contained in SEQ ID NO:X, the complement thereof, or the cDNA within the clone deposited with the ATCC. "Stringent hybridization conditions" refers to an overnight incubation at 42 degree C in a solution comprising 50% formamide, 5x SSC (750 mM NaCl, 75 mM trisodium citrate), 50 mM sodium phosphate (pH 7.6), 5x Denhardt's solution, 10% dextran sulfate, and 20 µg/ml denatured, sheared salmon sperm DNA, followed by washing the filters in 0.1x SSC at about 65 degree C.

Also contemplated are nucleic acid molecules that hybridize to the polynucleotides of the present invention at lower stringency hybridization conditions. Changes in the stringency of hybridization and signal detection are primarily accomplished through the manipulation of formamide concentration (lower percentages of formamide result in lowered stringency); salt conditions, or temperature. For example, lower stringency conditions include an overnight incubation at 37 degree C in a solution comprising 6X SSPE (20X SSPE = 3M NaCl; 0.2M NaH_2PO_4 ; 0.02M EDTA, pH 7.4), 0.5% SDS, 30% formamide, 100 µg/ml salmon sperm blocking DNA; followed by washes at 50 degree C with 1XSSPE, 0.1% SDS. In addition, to achieve even lower stringency, washes performed following stringent hybridization can be done at higher salt concentrations (e.g. 5X SSC).

Note that variations in the above conditions may be accomplished through the inclusion and/or substitution of alternate blocking reagents used to suppress background in hybridization experiments. Typical blocking reagents include Denhardt's reagent, BLOTTO, heparin, denatured salmon sperm DNA, and commercially available proprietary formulations. The inclusion of specific blocking

reagents may require modification of the hybridization conditions described above, due to problems with compatibility.

Of course, a polynucleotide which hybridizes only to polyA⁺ sequences (such as any 3' terminal polyA⁺ tract of a cDNA shown in the sequence listing), or to a
5 complementary stretch of T (or U) residues, would not be included in the definition of "polynucleotide," since such a polynucleotide would hybridize to any nucleic acid molecule containing a poly (A) stretch or the complement thereof (e.g., practically any double-stranded cDNA clone generated using oligo dT as a primer).

The polynucleotide of the present invention can be composed of any
10 polyribonucleotide or polydeoxribonucleotide, which may be unmodified RNA or DNA or modified RNA or DNA. For example, polynucleotides can be composed of single- and double-stranded DNA, DNA that is a mixture of single- and double-stranded regions, single- and double-stranded RNA, and RNA that is mixture of single- and double-stranded regions, hybrid molecules comprising DNA and RNA
15 that may be single-stranded or, more typically, double-stranded or a mixture of single- and double-stranded regions. In addition, the polynucleotide can be composed of triple-stranded regions comprising RNA or DNA or both RNA and DNA. A polynucleotide may also contain one or more modified bases or DNA or RNA backbones modified for stability or for other reasons. "Modified" bases include, for
20 example, tritylated bases and unusual bases such as inosine. A variety of modifications can be made to DNA and RNA; thus, "polynucleotide" embraces chemically, enzymatically, or metabolically modified forms.

The polypeptide of the present invention can be composed of amino acids joined to each other by peptide bonds or modified peptide bonds, i.e., peptide
25 isosteres, and may contain amino acids other than the 20 gene-encoded amino acids. The polypeptides may be modified by either natural processes, such as posttranslational processing, or by chemical modification techniques which are well known in the art. Such modifications are well described in basic texts and in more detailed monographs, as well as in a voluminous research literature. Modifications
30 can occur anywhere in a polypeptide, including the peptide backbone, the amino acid side-chains and the amino or carboxyl termini. It will be appreciated that the same

type of modification may be present in the same or varying degrees at several sites in a given polypeptide. Also, a given polypeptide may contain many types of modifications. Polypeptides may be branched, for example, as a result of ubiquitination, and they may be cyclic, with or without branching. Cyclic, branched, and branched cyclic polypeptides may result from posttranslation natural processes or may be made by synthetic methods. Modifications include acetylation, acylation, ADP-ribosylation, amidation, covalent attachment of flavin, covalent attachment of a heme moiety, covalent attachment of a nucleotide or nucleotide derivative, covalent attachment of a lipid or lipid derivative, covalent attachment of phosphatidylinositol, cross-linking, cyclization, disulfide bond formation, demethylation, formation of covalent cross-links, formation of cysteine, formation of pyroglutamate, formylation, gamma-carboxylation, glycosylation, GPI anchor formation, hydroxylation, iodination, methylation, myristoylation, oxidation, pegylation, proteolytic processing, phosphorylation, prenylation, racemization, selenoylation, sulfation, transfer-RNA mediated addition of amino acids to proteins such as arginylation, and ubiquitination. (See, for instance, PROTEINS - STRUCTURE AND MOLECULAR PROPERTIES, 2nd Ed., T. E. Creighton, W. H. Freeman and Company, New York (1993); POSTTRANSLATIONAL COVALENT MODIFICATION OF PROTEINS, B. C. Johnson, Ed., Academic Press, New York, pgs. 1-12 (1983); Seifter et al., Meth Enzymol 182:626-646 (1990); Rattan et al., Ann NY Acad Sci 663:48-62 (1992).)

"SEQ ID NO:X" refers to a polynucleotide sequence while "SEQ ID NO:Y" refers to a polypeptide sequence, both sequences identified by an integer specified in Table 1.

"A polypeptide having biological activity" refers to polypeptides exhibiting activity similar, but not necessarily identical to, an activity of a polypeptide of the present invention, including mature forms, as measured in a particular biological assay, with or without dose dependency. In the case where dose dependency does exist, it need not be identical to that of the polypeptide, but rather substantially similar to the dose-dependence in a given activity as compared to the polypeptide of the present invention (i.e., the candidate polypeptide will exhibit greater activity or not more than about 25-fold less and, preferably, not more than about tenfold less

activity, and most preferably, not more than about three-fold less activity relative to the polypeptide of the present invention.)

Many proteins (and translated DNA sequences) contain regions where the amino acid composition is highly biased toward a small subset of the available
5 residues. For example, membrane spanning domains and signal peptides (which are also membrane spanning) typically contain long stretches where Leucine (L), Valine (V), Alanine (A), and Isoleucine (I) predominate. Poly-Adenosine tracts (polyA) at the end of cDNAs appear in forward translations as poly-Lysine (poly-K) and poly-Phenylalanine (poly-F) when the reverse complement is translated. These regions are
10 often referred to as "low complexity" regions.

Such regions can cause database similarity search programs such as BLAST to find high-scoring sequence matches that do not imply true homology. The problem is exacerbated by the fact that most weight matrices (used to score the alignments generated by BLAST) give a match between any of a group of hydrophobic amino
15 acids (L, V and I) that are commonly found in certain low complexity regions almost as high a score as for exact matches.

In order to compensate for this, BLASTX.2 (version 2.0a5MP-WashU) employs two filters ("seg" and "xnu") which "mask" the low complexity regions in a particular sequence. These filters parse the sequence for such regions, and create a
20 new sequence in which the amino acids in the low complexity region have been replaced with the character "X". This is then used as the input sequence (sometimes referred to herein as "Query" and/or "Q") to the BLASTX program. While this regime helps to ensure that high-scoring matches represent true homology, there is a negative consequence in that the BLASTX program uses the query sequence that has
25 been masked by the filters to draw alignments.

Thus, a stretch of "X"s in an alignment shown in the following application does not necessarily indicate that either the underlying DNA sequence or the translated protein sequence is unknown or uncertain. Nor is the presence of such stretches meant to indicate that the sequence is identical or not identical to the
30 sequence disclosed in the alignment of the present invention. Such stretches may simply indicate that the BLASTX program masked amino acids in that region due to

the detection of a low complexity region, as defined above. In all cases, the reference sequence(s) (sometimes referred to herein as "Subject", "Sbjct", and/or "S") indicated in the specification, sequence table (Table 1), and/or the deposited clone is (are) the definitive embodiment(s) of the present invention, and should not be construed as
5 limiting the present invention to the partial sequence shown in an alignment, unless specifically noted otherwise herein.

Polynucleotides and Polypeptides of the Invention

10 FEATURES OF PROTEIN ENCODED BY GENE NO: 1

Many polynucleotide sequences, such as EST sequences, are publicly available and accessible through sequence databases. Some of these sequences are related to SEQ ID NO:11 and may have been publicly available prior to conception of the present invention. Preferably, such related polynucleotides are specifically
15 excluded from the scope of the present invention. To list every related sequence would be cumbersome. Accordingly, preferably excluded from the present invention are one or more polynucleotides comprising a nucleotide sequence described by the general formula of a-b, where a is any integer between 1 to 1215 of SEQ ID NO:11, b is an integer of 15 to 1229, where both a and b correspond to the positions of
20 nucleotide residues shown in SEQ ID NO:11, and where b is greater than or equal to a + 14.

FEATURES OF PROTEIN ENCODED BY GENE NO: 2

The computer algorithm BLASTX has been used to determine that the
25 translation product of this gene shares sequence homology with, as a non-limiting example, the sequence accessible through the following database accession no. gil50313 (all information available through the recited accession number is incorporated herein by reference) which is described therein as "carboxypeptidase H [Mus musculus]". A partial alignment demonstrating the observed homology is shown
30 immediately below.


```

>gi|50313 carboxypeptidase H [Mus musculus] >pir|S16383|S16383
carboxypeptidase
      H (EC 3.4.17.10) - mouse
      Length = 477
5      Plus Strand HSPs:

      Score = 549 (193.3 bits), Expect = 3.1e-99, Sum P(3) = 3.1e-99
      Identities = 102/109 (93%), Positives = 105/109 (96%), Frame = +2
10      Query:   11 NNHLLKNMKKIVDQNTKLAPETKAVIHWIMDIPFVLSANLHGGDLVANYPYDETRSGSAH
      190
      NNHLLKN+KKIVDQN+KLAPETKAVIHWIMDIPFVLSANLHGGDLVANYPYDETRSG+AH
      Sbjet:   208 NNHLLKNLKKIVDQNSKLAPETKAVIHWIMDIPFVLSANLHGGDLVANYPYDETRSGTAH
15      267

      Query:   191 EYSSSPDDAIFQSLARAYSSFPAMSDPNRPPCRKNDDSSSFVDGTTNG 337
      EYSS PDDAIFQSLARAYSSFP MSDPNRPPC KNDDSSSF DGTNG
      Sbjet:   268 EYSSCPDDAIFQSLARAYSSFPVMSDPNRPPCCKNDDSSSFDDGTTNG 316
20

      Score = 382 (134.5 bits), Expect = 3.1e-99, Sum P(3) = 3.1e-99
      Identities = 76/98 (77%), Positives = 83/98 (84%), Frame = +1

      Query:   340 AWYSVPGGMQDFNYLSS-NCFEITVELSCEKFPPEETLKTYWEDNKNSLISYLEQIHRGV
25      516
      AWYSVPGGMQDFNYLSS NCFEITVEL+CEKFPPEETLK+YWEDNKNSLI+YLEQIHRGV
      Sbjet:   318 AWYSVPGGMQDFNYLSSNCFEITVELTCEKFPPEETLKSYWEDNKNSLINYLEQIHRGV
      377

      Query:   517 KGFVRDLQGNFIAMAT-NLRGRNRPRCIYIRKG--WRL 621
      KGFVRDLQGNP A AT ++ G + + G WRL
      Sbjet:   378 KGFVRDLQGNPNANATISVDGIDHDVTSKDGWDYWRL 415
30

```

The segment of gi|50313 that is shown as "Sbjet" above is set out as
 35 sequence(s) SEQ ID NO. 105 and SEQ ID NO. 107. Based on the structural similarity
 these homologous polypeptides are expected to share at least some biological
 activities. Such activities are known in the art and described elsewhere herein. Assays
 for determining such activities are also known in the art, some of which have been
 described elsewhere herein. Preferred polypeptides of the invention comprise
 40 polypeptides having the amino acid sequence(s) set out as SEQ ID NO. 106 and SEQ
 ID NO. 108 which correspond to the Query sequence in the alignment shown above
 (gaps introduced in the sequence by the computer are, of course, removed).

It has been discovered that this gene is expressed primarily in the following
 tissues: Human Amygdala and to a lesser extent in Nine Week Old Early Stage
 45 Human, Soares infant brain 1NIB, Human Prostate, NCI_CGAP_Lu5, Human Testes,
 Human Chondrosarcoma, Morton Fetal Cochlea, normalized infant brain cDNA, 12
 Week Old Early Stage Human, II, Human Cerebellum, Human Substantia Nigra,
 Human 8 Week Whole Embryo, Soares_parathyroid_tumor_NbHPA,

- Soares_fetal_liver_spleen_1NFLS_S1, Hippocampus, Alzheimer Subtracted, 12 Week Early Stage Human II; Reexcision, Human Adrenal Gland Tumor, H. Frontal cortex,epileptic;re-excision, Stratagene lung (#937210), H. Epididymus, cauda, Human Amygdala;re-excision, Healing groin wound - zero hr post-incision (control),
- 5 Human Osteosarcoma, Temporal cortex-Alzheimzer; subtracted, Human Hypothalamus,Schizophrenia, Human umbilical vein endothelial cells, IL-4 induced, Spinal cord, 12 Week Old Early Stage Human, Soares_multiple_sclerosis_2NbHMSP, Human Fetal Lung III, Human Fetal Heart, NCI_CGAP_Pr3, Colon Normal III, Stratagene endothelial cell 937223, Soares fetal
- 10 liver spleen 1NFLS, Soares_pineal_gland_N3HPG, Human Brain, Human Fetal Lung, Whole 6 Week Old Embryo, Dermatofibrosarcoma Protuberance, Soares_fetal_lung_NbHL19W, Stratagene fetal retina 937202, Human Manic Depression Tissue, Soares_total_fetus_Nb2HF8_9w, Human Brain, Striatum, human ovarian cancer, NCI_CGAP_Pr18, Soares_pregnant_uterus_NbHPU, Human
- 15 Hippocampus, Soares_fetal_lung_NbHL19W, Soares_fetal_heart_NbHH19W, NTERA2, control, NCI_CGAP_Pr3, Human heart cDNA (YNakamura), Human retina cDNA randomly primed sublibrary, NCI_CGAP_Thy1, NCI_CGAP_GC4, Osteoblasts, Human Hippocampus, prescreened, Human Fetal Brain, H. Adult Spleen, ziplox, Stratagene hNT neuron (#937233), HUman Fetal Brain, normalized 100024F,
- 20 Human Hippocampus, Human Fetal Brain, normalized c5-11-26, Human Fetal Brain, normalized A5002F, Brain; normal, Larynx Normal, Pharynx Carcinoma, Human Normal Cartilage Fraction IV, Normal Prostate, Brain Amygdala Depression, Human Normal Cartilage Fraction II, Human Prostate, subtracted, Human 8 Week Whole Embryo, subtracted, STRIATUM DEPRESSION, Soares_testis_NHT, Frontal Lobe,
- 25 Dementia, H. Striatum Depression, subt, Human Cerebellum, subtracted, Human Cardiomyopathy, subtracted, H. Atrophic Endometrium, Human Thyroid, 1-NIB, Normalized infant brain, Bento Soares, Soares_testis_NHT, Soares_NFL_T_GBC_S1, Soares_fetal_heart_NbHH19W, Soares_total_fetus_Nb2HF8_9w, Stratagene ovary (#937217), H. Whole Brain #2, re-
- 30 excision, Human Whole Brain, re-excision, Human Epididymus, Stomach cancer (human);re-excision, Morton Fetal, Human Hypothalamus,schizophrenia, re-excision,

Human Synovium, Human Colon Cancer;re-excision, human corpus colosum,
 Healing groin wound; 7.5 hours post incision, H. Kidney Cortex, subtracted, Human
 Whole Brain #2 - Oligo dT > 1.5Kb, Human endometrial stromal cells-treated with
 progesterone, Human Frontal Cortex, Schizophrenia, Soares_fetal_heart_NbHH19W,
 5 Human Osteoclastoma, re-excision, Human endometrial stromal cells, Human Ovary,
 Soares_fetal_heart_NbHH19W, H. Meningima, M1, Testis 1, Soares_NbHFB,
 Stratagene lung carcinoma 937218, H. Kidney Medulla, re-excision, Human Fetal
 Kidney, NCI_CGAP_Lu5, NCI_CGAP_Alv1, Human Umbilical Vein Endothelial
 Cells, uninduced, Merkel Cells, Human Thymus Stromal Cells, Human Whole Six
 10 Week Old Embryo, Human Liver, normal, NCI_CGAP_Ew1, Fetal Heart, Early
 Stage Human Brain, Colon Tumor II, Human Testes, Reexcision, human tonsils,
 Human adult (K.Okubo), Human fetal heart, Lambda ZAP Express, NCI_CGAP_Br2,
 NCI_CGAP_Ew1, NCI_CGAP_GC5, NCI_CGAP_Pr6, NCI_CGAP_Pr10, Smooth
 muscle,control, Human Endometrial Tumor, Hodgkin's Lymphoma II, Human
 15 pancreatic islet, NCI_CGAP_Br2, NCI_CGAP_Kid3, NCI_CGAP_Kid5,
 NCI_CGAP_Brn23.

Preferred epitopes include those comprising a sequence shown in SEQ ID NO.
 59 as residues: Gln-7 to Pro-13, Pro-43 to Gly-50, Glu-54 to Asp-60, Ala-77 to Ser-
 94, Ser-134 to Leu-139.

20 The translation product of this gene was shown to have homology to
 Carboxypeptidase H which is a putative post-translational processing enzyme that
 removes basic amino acid residues from intermediates during protein hormone
 biosynthesis. Moreover, the tissue distribution in neural tissues suggests the protein
 product of this clone is useful for the detection, treatment, and/or prevention of
 25 neurodegenerative disease states, behavioral disorders, or inflammatory conditions
 which include, but are not limited to Alzheimerís Disease, Parkinsonís Disease,
 Huntingtonís Disease, Tourette Syndrome, meningitis, encephalitis, demyelinating
 diseases, peripheral neuropathies, neoplasia, trauma, congenital malformations, spinal
 cord injuries, ischemia and infarction, aneurysms, hemorrhages, schizophrenia,
 30 mania, dementia, paranoia, obsessive compulsive disorder, depression, panic disorder,
 learning disabilities, ALS, psychoses, autism, and altered behaviors, including

disorders in feeding, sleep patterns, balance, and perception. In addition, elevated expression of this gene product in regions of the brain suggests it plays a role in normal neural function. Potentially, this gene product is involved in synapse formation, neurotransmission, learning, cognition, homeostasis, or neuronal differentiation or survival.

Moreover, the expression within embryonic tissue and other cellular sources marked by proliferating cells suggests this protein may play a role in the regulation of cellular division, and may show utility in the diagnosis and treatment of cancer and other proliferative disorders. Similarly, developmental tissues rely on decisions involving cell differentiation and/or apoptosis in pattern formation. Dysregulation of apoptosis can result in inappropriate suppression of cell death, as occurs in the development of some cancers, or in failure to control the extent of cell death, as is believed to occur in acquired immunodeficiency and certain neurodegenerative disorders, such as spinal muscular atrophy (SMA). Therefore, the polynucleotides and polypeptides of the present invention are useful in treating, detecting, and/or preventing said disorders and conditions, in addition to other types of degenerative conditions. Thus this protein may modulate apoptosis or tissue differentiation and would be useful in the detection, treatment, and/or prevention of degenerative or proliferative conditions and diseases. Protein, as well as, antibodies directed against the protein may show utility as a tumor marker and/or immunotherapy targets for the above listed tissues.

Many polynucleotide sequences, such as EST sequences, are publicly available and accessible through sequence databases. Some of these sequences are related to SEQ ID NO:12 and may have been publicly available prior to conception of the present invention. Preferably, such related polynucleotides are specifically excluded from the scope of the present invention. To list every related sequence would be cumbersome. Accordingly, preferably excluded from the present invention are one or more polynucleotides comprising a nucleotide sequence described by the general formula of a-b, where a is any integer between 1 to 1524 of SEQ ID NO:12, b is an integer of 15 to 1538, where both a and b correspond to the positions of

nucleotide residues shown in SEQ ID NO:12, and where b is greater than or equal to a + 14.

FEATURES OF PROTEIN ENCODED BY GENE NO: 3

5 It has been discovered that this gene is expressed primarily in the following tissues: Human Activated Monocytes, Human Chondrosarcoma, and Human Placenta.

Many polynucleotide sequences, such as EST sequences, are publicly available and accessible through sequence databases. Some of these sequences are related to SEQ ID NO:13 and may have been publicly available prior to conception of
10 the present invention. Preferably, such related polynucleotides are specifically excluded from the scope of the present invention. To list every related sequence would be cumbersome. Accordingly, preferably excluded from the present invention are one or more polynucleotides comprising a nucleotide sequence described by the general formula of a-b, where a is any integer between 1 to 1018 of SEQ ID NO:13, b
15 is an integer of 15 to 1032, where both a and b correspond to the positions of nucleotide residues shown in SEQ ID NO:13, and where b is greater than or equal to a + 14.

FEATURES OF PROTEIN ENCODED BY GENE NO: 4

20 The computer algorithm BLASTX has been used to determine that the translation product of this gene shares sequence homology with, as a non-limiting example, the sequence accessible through the following database accession no. gil1399105 (all information available through the recited accession number is incorporated herein by reference) which is described therein as "phosphatidylinositol
25 (4,5)bispophosphate 5-phosphatase homolog". A partial alignment demonstrating the observed homology is shown immediately below.

```
30 >gi|1399105 phosphatidylinositol (4,5)bispophosphate 5-phosphatase homolog
[Homo sapiens]
Length = 397

Plus Strand HSPs:

35 Score = 1314 (462.6 bits), Expect = 1.7e-133, P = 1.7e-133
Identities = 260/382 (68%), Positives = 262/382 (68%), Frame = +2
```

Query: 404 LHFVKFAIDSDQLHQLWEKDQLNMAKNTWPILKGFQEGPLNFAPTFFKFDVGTNKYDTSK
 583
 5 Sbjct: 4 LHFVKFAIDSDQLHQLWEKDQLNMAKNTWPILKGFQEGPLNFAPTFFKFDVGTNKYDTSK
 63
 Query: 584 KRKPAWTDRLWKVKAXXXXXXXXXXKSHRLQVTQHSYRSHMEYTVSDHKPVXAQFLLQF
 763
 10 Sbjct: 64 KRKPAWTDRLWKVKAPGGGSPSPSGRKSRLQVTQHSYRSHMEYTVSDHKPVAAQFLLQF
 123
 Query: 764 AFQGRHXXXXXXXXXXXXXPSRRW*GYRXETXFXRSSWDWIGLYRVGFRHCKDYVXYVW
 943
 15 Sbjct: 124 AF+ P + YR ET F RSSWDWIGLYRVGFRHCKDYV YVW
 182
 Query: 944 AKHEDVDGNTYQVTFSEESLPKGHGDFILGYXSHNHSILIGITEPFQIXXXXXXXXXXXXX
 1123
 20 Sbjct: 183 AKHEDVDGNTYQVTFSEESLPKGHGDFILGY SHNHSILIGITEPFQI
 242
 25 Query: 1124 XXXXXXXXXXXXXXXLELLAXXXXXXXXXXXXXXXXXXXXXLXARFPGLALRPSSREXXXXXX
 1303
 Sbjct: 243 DSSGTSSEGEDDSTLELLAPKSRSPSPGSKRHRSRSPGLARFPGLALRPSSRERRGASR
 30 302
 Query: 1304 XXXXXXXXXXXXVAPDXXXXXXXXXXXXXXXXXXXXXWAFPPAVPRSLGLLPALRLETVDP
 1483
 35 Sbjct: 303 VAPD WAFPPAVPRSLGLLPALRLETVDP
 362
 Query: 1484 GGGGSWGPDREALAPNSLSPSP 1549
 40 Sbjct: 363 GGGGSWGPDREALAPNSLSPSP 384

The segment of gil1399105 that is shown as "Sbjct" above is set out as
 sequence(s) SEQ ID NO. 109. Based on the structural similarity these homologous
 polypeptides are expected to share at least some biological activities. Such activities
 45 are known in the art and described elsewhere herein. Assays for determining such
 activities are also known in the art, some of which have been described elsewhere
 herein. Preferred polypeptides of the invention comprise polypeptides having the
 amino acid sequence(s) set out as SEQ ID NO. 110 which correspond to the Query
 sequence in the alignment shown above (gaps introduced in the sequence by the
 50 computer are, of course, removed).

It has been discovered that this gene is expressed primarily in the following
 tissues: Soares' infant brain 1NIB and to a lesser extent in Human Cerebellum,

subtracted, Human Whole Six Week Old Embryo, Stratagene neuroepithelium (#937231), Human Testes Tumor, Human Cerebellum.

Preferred epitopes include those comprising a sequence shown in SEQ ID NO. 61 as residues: Asp-20 to Thr-26, Leu-30 to Gly-38, Asp-63 to Phe-72, Gly-160 to Trp-175, Gly-189 to Ser-197, Thr-214 to Val-221.

The tissue distribution in neural tissues suggests the protein product of this clone is useful for the detection, treatment, and/or prevention of neurodegenerative disease states, behavioral disorders, or inflammatory conditions which include, but are not limited to Alzheimer's Disease, Parkinson's Disease, Huntington's Disease, Tourette Syndrome, meningitis, encephalitis, demyelinating diseases, peripheral neuropathies, neoplasia, trauma, congenital malformations, spinal cord injuries, ischemia and infarction, aneurysms, hemorrhages, schizophrenia, mania, dementia, paranoia, obsessive compulsive disorder, depression, panic disorder, learning disabilities, ALS, psychoses, autism, and altered behaviors, including disorders in feeding, sleep patterns, balance, and perception. In addition, elevated expression of this gene product in regions of the brain suggests it plays a role in normal neural function. Potentially, this gene product is involved in synapse formation, neurotransmission, learning, cognition, homeostasis, or neuronal differentiation or survival. Moreover, the expression within infant tissue and other cellular sources marked by proliferating cells suggests this protein may play a role in the regulation of cellular division, and may show utility in the diagnosis and treatment of cancer and other proliferative disorders. Similarly, developmental tissues rely on decisions involving cell differentiation and/or apoptosis in pattern formation. Dysregulation of apoptosis can result in inappropriate suppression of cell death, as occurs in the development of some cancers, or in failure to control the extent of cell death, as is believed to occur in acquired immunodeficiency and certain neurodegenerative disorders, such as spinal muscular atrophy (SMA). Therefore, the polynucleotides and polypeptides of the present invention are useful in treating, detecting, and/or preventing said disorders and conditions, in addition to other types of degenerative conditions. Thus this protein may modulate apoptosis or tissue differentiation and would be useful in the detection, treatment, and/or prevention of degenerative or

proliferative conditions and diseases. Protein, as well as, antibodies directed against the protein may show utility as a tumor marker and/or immunotherapy targets for the above listed tissues.

Many polynucleotide sequences, such as EST sequences, are publicly
5 available and accessible through sequence databases. Some of these sequences are related to SEQ ID NO:14 and may have been publicly available prior to conception of the present invention. Preferably, such related polynucleotides are specifically excluded from the scope of the present invention. To list every related sequence would be cumbersome. Accordingly, preferably excluded from the present invention
10 are one or more polynucleotides comprising a nucleotide sequence described by the general formula of a-b, where a is any integer between 1 to 1882 of SEQ ID NO:14, b is an integer of 15 to 1896, where both a and b correspond to the positions of nucleotide residues shown in SEQ ID NO:14, and where b is greater than or equal to a + 14.

15

FEATURES OF PROTEIN ENCODED BY GENE NO: 5

It has been discovered that this gene is expressed primarily in the following tissues: Human Whole Brain, re-excision, Human Cerebellum.

Many polynucleotide sequences, such as EST sequences, are publicly
20 available and accessible through sequence databases. Some of these sequences are related to SEQ ID NO:15 and may have been publicly available prior to conception of the present invention. Preferably, such related polynucleotides are specifically excluded from the scope of the present invention. To list every related sequence would be cumbersome. Accordingly, preferably excluded from the present invention
25 are one or more polynucleotides comprising a nucleotide sequence described by the general formula of a-b, where a is any integer between 1 to 1216 of SEQ ID NO:15, b is an integer of 15 to 1230, where both a and b correspond to the positions of nucleotide residues shown in SEQ ID NO:15, and where b is greater than or equal to a + 14.

30

FEATURES OF PROTEIN ENCODED BY GENE NO: 6

The computer algorithm BLASTX has been used to determine that the translation product of this gene shares sequence homology with, as a non-limiting example, the sequence accessible through the following database accession no. gnl|PID|d1029109 (all information available through the recited accession number is incorporated herein by reference) which is described therein as "TIP49 [Homo sapiens]". A partial alignment demonstrating the observed homology is shown immediately below.

```

10      >gnl|PID|d1029109 (AB012122) TIP49 [Homo sapiens] >gi|3243035 (AF070735)
      RuvB-like protein RUVBL1 [Homo sapiens] >gnl|PID|e1352152
      (AJ010058) Nuclear matrix protein NMP238 [Homo sapiens]
      >gi|4151525
      (AF099084) Pontin52 [Homo sapiens] >sp|D1029109|D1029109
15      TIP49.
      >sp|G3243035|G3243035 RUVB-LIKE PROTEIN RUVBL1.
      >sp|E1352152|E1352152 NUCLEAR MATRIX PROTEIN NMP238.
      Length = 456

20      Plus Strand HSPs:

      Score = 613 (215.8 bits), Expect = 1.3e-222, Sum P(5) = 1.3e-222
      Identities = 118/119 (99%), Positives = 119/119 (100%), Frame = +2

25      Query:   473 EVTELTPCETENPMGGYGKTISHVIIGLKTAKGTKQLKLDPSIFESLQKERVEAGDVIYI
      652
      Sbjct:   134 EVTELTPCETENPMGGYGKTISHVIIGLKTAKGTKQLKLDPSIFESLQKERVEAGDVIYI
      193

30      Query:   653 EANSgAVKRQGRCDTYATEFDLEAEYVPLPKGDVHKKKEIIQDVTLHDLVDANARPRG
      829
      Sbjct:   194 EANSgAVKRQGRCDTYATEFDLEAEYVPLPKGDVHKKKEIIQDVTLHDLVDANARPG
      252

35      Score = 589 (207.3 bits), Expect = 1.3e-222, Sum P(5) = 1.3e-222
      Identities = 119/121 (98%), Positives = 120/121 (99%), Frame = +1

40      Query:   76 MKIEEVKSTTKTQRIASHSHVKGLGLDESLAKQAASGLVGQENAREACGVIVELIKSKK
      255
      Sbjct:    1 MKIEEVKSTTKTQRIASHSHVKGLGLDESLAKQAASGLVGQENAREACGVIVELIKSKK
      60

45      Query:   256 MAGRAVLLAGPPGTGKTALALAIAQELGSKVPFCPMVGSEVYSTEIKKTEVLMENFRRPL
      435
      Sbjct:    61 MAGRAVLLAGPPGTGKTALALAIAQELGSKVPFCPMVGSEVYSTEIKKTEVLMENFRR +
      120

50      Query:   436 G 438
      G
      Sbjct:   121 G 121

55      Score = 586 (206.3 bits), Expect = 1.3e-222, Sum P(5) = 1.3e-222
      Identities = 117/128 (91%), Positives = 122/128 (95%), Frame = +1

```

Query: 838 ILSMMGQLMKPRRQK-SRQTSGEINKVVNKYIDQGIAELVPGVLFVDEVHMLDIECFITYL
 1014
 5 Sbjct: 256 ILSMMGQLMKP++ + + + GEINKVVNKYIDQGIAELVPGVLFVDEVHMLDIECFITYL
 315
 Query: 1015 HRALESSIAPIVIFASNRGNCVIRGTEDITSPHGIPLXLLDRVMIIRTMLYTPQEMKQII
 1194
 10 Sbjct: 316 HRALESSIAPIVIFASNRGNCVIRGTEDITSPHGIPL LLDVRMIIRTMLYTPQEMKQII
 375
 Query: 1195 KIRAQTEG 1218
 15 KIRAQTEG
 Sbjct: 376 KIRAQTEG 383

The segment of gnlIPIDd1029109 that is shown as "Sbjct" above is set out as
 20 sequence(s) SEQ ID NO. 111, SEQ ID NO. 113, and SEQ ID NO. 115. Based on the
 structural similarity these homologous polypeptides are expected to share at least
 some biological activities. Such activities are known in the art and described
 elsewhere herein. Assays for determining such activities are also known in the art,
 some of which have been described elsewhere herein. Preferred polypeptides of the
 25 invention comprise polypeptides having the amino acid sequence(s) set out as SEQ ID
 NO. 112, SEQ ID NO. 114, and SEQ ID NO. 116 which correspond to the Query
 sequence in the alignment shown above (gaps introduced in the sequence by the
 computer are, of course, removed).

It has been discovered that this gene is expressed primarily in the following
 30 tissues: Soares fetal liver spleen 1NFLS and to a lesser extent in Soares adult brain
 N2b5HB55Y, Human Chondrosarcoma, Stratagene colon HT29 (#937221), Activated
 T-cell(12h)/Thiouridine-re-excision, Pancreas Islet Cell Tumor, Human Testes
 Tumor, Activated T-Cell (12hs)/Thiouridine labelledEco, HUMAN B CELL
 LYMPHOMA, Nine Week Old Early Stage Human, Stratagene pancreas (#937208),
 35 Human Fetal Brain, Human colon carcinoma (HCC) cell line, remake, Human
 Primary Breast Cancer Reexcision, Soares_fetal_lung_NbHL19W, Human B Cell
 8866, Weizmann Olfactory Epithelium, NCI_CGAP_GC4, NCI_CGAP_Co12,
 NCI_CGAP_GCB1, NCI_CGAP_Brn25, Resting T-Cell; re-excision, Jurkat T-Cell,
 S phase, HUMAN JURKAT MEMBRANE BOUND POLYSOMES, Human Adult
 40 Testes, Large Inserts, Reexcision, Smooth muscle, serum treated, Human Placenta,

Human Placenta, Endothelial-induced, Endothelial cells-control,
 Soares_pregnant_uterus_NbHPU, Smooth muscle,control, Human Testes, Bone
 Marrow Cell Line (RS4;11), Human 8 Week Whole Embryo, Stomach Tumour,
 Human retina cDNA Tsp509I-cleaved sublibrary, NCI_CGAP_HN4,
 5 NCI_CGAP_Lu5, NCI_CGAP_Ov2, NCI_CGAP_Kid3, NCI_CGAP_Kid6,
 NCI_CGAP_Pr24, NCI_CGAP_Pr25, NCI_CGAP_Brn23, Saos2 Cells; Untreated,
 Morton Fetal Cochlea, Soares_fetal_lung_NbHL19W, H. Striatum Depression, sublt,
 Human Adult Retina, Human Thyroid, Early Stage Human Lung, subtracted, Human
 T-cell lymphoma;re-excision, Cem cells cyclohexamide treated, Human Normal
 10 Breast, Ovarian Tumor 10-3-95, Stomach cancer (human);re-excision, Myoloid
 Progenitor Cell Line, Human Thymus, Human Dermal Endothelial Cells,untreated,
 Human Brain, Striatum, L428, Human Fetal Kidney, Stratagene fetal spleen
 (#937205), NCI_CGAP_Br2, NCI_CGAP_Co9, NCI_CGAP_Lu1, Human Umbilical
 Vein Endothelial Cells, uninduced, Human Uterine Cancer, Human umbilical vein
 15 endothelial cells, IL-4 induced, Epithelial-TNFa and INF induced, Human Thymus,
 H. Frontal cortex,epileptic;re-excision, Human T-Cell Lymphoma, Brain frontal
 cortex, Soares breast 3NbHBst, Adipocytes, Human Synovial Sarcoma, Human Fetal
 Lung III, human tonsils, NCI_CGAP_GCB1, Colon Normal III,
 Soares_fetal_liver_spleen_1NFLS_S1, Human Microvascular Endothelial Cells, fract.
 20 A, Stratagene colon (#937204), Hodgkin's Lymphoma II, Keratinocyte, Human
 Cerebellum.

Preferred epitopes include those comprising a sequence shown in SEQ ID NO.
 63 as residues: Gln-40 to Cys-48.

The tissue distribution suggests the protein product of this clone is useful for
 25 the diagnosis and treatment of a variety of immune system disorders. Moreover, the
 expression of this gene product suggests a role in regulating the proliferation;
 survival; differentiation; and/or activation of hematopoietic cell lineages, including
 blood stem cells. This gene product may be involved in the regulation of cytokine
 production, antigen presentation, or other processes suggesting a usefulness in the
 30 treatment of cancer (e.g. by boosting immune responses).

Since the gene is expressed in cells of lymphoid origin, the natural gene product may be involved in immune functions. Therefore it may be also used as an agent for immunological disorders including arthritis, asthma, immunodeficiency diseases such as AIDS, leukemia, rheumatoid arthritis, granulomatous disease, inflammatory bowel disease, sepsis, acne, neutropenia, neutrophilia, psoriasis, hypersensitivities, such as T-cell mediated cytotoxicity; immune reactions to transplanted organs and tissues, such as host-versus-graft and graft-versus-host diseases, or autoimmunity disorders, such as autoimmune infertility, lense tissue injury, demyelination, systemic lupus erythematosus, drug induced hemolytic anemia, rheumatoid arthritis, Sjogren's disease, scleroderma and tissues. Moreover, the protein may represent a secreted factor that influences the differentiation or behavior of other blood cells, or that recruits hematopoietic cells to sites of injury. In addition, this gene product may have commercial utility in the expansion of stem cells and committed progenitors of various blood lineages, and in the differentiation and/or proliferation of various cell types. Protein, as well as, antibodies directed against the protein may show utility as a tumor marker and/or immunotherapy targets for the above listed tissues.

Many polynucleotide sequences, such as EST sequences, are publicly available and accessible through sequence databases. Some of these sequences are related to SEQ ID NO:16 and may have been publicly available prior to conception of the present invention. Preferably, such related polynucleotides are specifically excluded from the scope of the present invention. To list every related sequence would be cumbersome. Accordingly, preferably excluded from the present invention are one or more polynucleotides comprising a nucleotide sequence described by the general formula of a-b, where a is any integer between 1 to 1730 of SEQ ID NO:16, b is an integer of 15 to 1744, where both a and b correspond to the positions of nucleotide residues shown in SEQ ID NO:16, and where b is greater than or equal to a + 14.

30 FEATURES OF PROTEIN ENCODED BY GENE NO: 7

The computer algorithm BLASTX has been used to determine that the translation product of this gene shares sequence homology with, as a non-limiting example, the sequence accessible through the following database accession no. gil2580578 (all information available through the recited accession number is incorporated herein by reference) which is described therein as "ubiquitous TPR motif". A partial alignment demonstrating the observed homology is shown immediately below.

```

10  >gi|2580578 (AF000996) ubiquitous TPR motif, Y isoform [Homo sapiens]
      Length = 1079

      Minus Strand HSPs:

15  Score = 257 (90.5 bits), Expect = 6.1e-18, P = 6.1e-18
      Identities = 53/81 (65%), Positives = 56/81 (69%), Frame = -2

      Query:   999 AGVQWLNLSSLQPPHPGFK*FSCLSLPCSWDYRSAPPGPGNFCIFRRDGVRHVGQNSLEL
820
      AG+QW +LSSLQPP PGFK FS LSLP SW+YR P P NFCIF G HVGQ LEL
20  Sbjct:   996 AGMQWCDLSSLQPPPPGFKRFSHLPLNSWNYRHLPSCPTNFCIFVETGFHHVGQACLEL
1055

      Query:   819 LNSGDPALAPQSGGTTGVSH 757
      L SG A A QS G TGVSH
25  Sbjct:  1056 LTSGGLLASASQSAGITGVSH 1076

      Score = 140 (49.3 bits), Expect = 3.5e-05, P = 3.5e-05
      Identities = 30/43 (69%), Positives = 33/43 (76%), Frame = -3

30  Query:  1553 FVFVVQMGLHHVGQAGLKLPTSGDPPASASQSAGITGVSHHAQ 1425
      F V+ G HHVGQA L+L TSG ASASQSAGITGVSHHA+
      Sbjct:  1037 FCIFVETGFHHVGQACLELLTSGGLLASASQSAGITGVSHHAR 1079

35  Score = 122 (42.9 bits), Expect = 0.00073, Sum P(2) = 0.00073
      Identities = 22/39 (56%), Positives = 27/39 (69%), Frame = -2

      Query:  1647 MQPPPPGLKQFSRLSPSSWDYRCVPPRLANFCICSTNG 1531
      +QPPPPG K+FS LS P+SW+YR +P NFCI G
40  Sbjct:  1006 LQPPPPGFKRFSHLPLNSWNYRHLPSCPTNFCIFVETG 1044

      Score = 54 (19.0 bits), Expect = 0.00073, Sum P(2) = 0.00073
      Identities = 12/17 (70%), Positives = 12/17 (70%), Frame = -3

45  Query:   932 ASASHVAGTTGVHHQAR 882
      ASAS AG TGV H AR
      Sbjct:  1063 ASASQSAGITGVSHHAR 1079

```

The segment of gil2580578 that is shown as "Sbjct" above is set out as sequence(s) SEQ ID NO. 117, SEQ ID NO. 119, SEQ ID NO. 121, and SEQ ID NO. 123. Based on the structural similarity these homologous polypeptides are expected to

share at least some biological activities. Such activities are known in the art and described elsewhere herein. Assays for determining such activities are also known in the art, some of which have been described elsewhere herein. Preferred polypeptides of the invention comprise polypeptides having the amino acid sequence(s) set out as
5 SEQ ID NO. 118, SEQ ID NO. 120, SEQ ID NO. 122, and SEQ ID NO. 124 which correspond to the Query sequence in the alignment shown above (gaps introduced in the sequence by the computer are, of course, removed).

It has been discovered that this gene is expressed primarily in the following tissues: Human Microvascular Endothelial Cells, fract. A, Human Cerebellum.

10 The tissue distribution in microvascular tissue suggests the protein is useful in the detection, treatment, and/or prevention of vascular conditions, which include, but are not limited to, microvascular disease, vascular leak syndrome, aneurysm, stroke, atherosclerosis, arteriosclerosis, or embolism. Protein, as well as, antibodies directed against the protein may show utility as a tumor marker and/or immunotherapy targets
15 for the above listed tissues.

Many polynucleotide sequences, such as EST sequences, are publicly available and accessible through sequence databases. Some of these sequences are related to SEQ ID NO:17 and may have been publicly available prior to conception of the present invention. Preferably, such related polynucleotides are specifically
20 excluded from the scope of the present invention. To list every related sequence would be cumbersome. Accordingly, preferably excluded from the present invention are one or more polynucleotides comprising a nucleotide sequence described by the general formula of a-b, where a is any integer between 1 to 1700 of SEQ ID NO:17, b is an integer of 15 to 1714, where both a and b correspond to the positions of
25 nucleotide residues shown in SEQ ID NO:17, and where b is greater than or equal to a + 14.

FEATURES OF PROTEIN ENCODED BY GENE NO: 8

It has been discovered that this gene is expressed primarily in the following
30 tissues: Human Cerebellum, Human Greater Omentum, fll remake, human tonsils.

Many polynucleotide sequences, such as EST sequences, are publicly available and accessible through sequence databases. Some of these sequences are related to SEQ ID NO:18 and may have been publicly available prior to conception of the present invention. Preferably, such related polynucleotides are specifically excluded from the scope of the present invention. To list every related sequence would be cumbersome. Accordingly, preferably excluded from the present invention are one or more polynucleotides comprising a nucleotide sequence described by the general formula of a-b, where a is any integer between 1 to 1405 of SEQ ID NO:18, b is an integer of 15 to 1419, where both a and b correspond to the positions of nucleotide residues shown in SEQ ID NO:18, and where b is greater than or equal to a + 14.

FEATURES OF PROTEIN ENCODED BY GENE NO: 9

The computer algorithm BLASTX has been used to determine that the translation product of this gene shares sequence homology with, as a non-limiting example, the sequence accessible through the following database accession no. gnl|PID|e1345389 (all information available through the recited accession number is incorporated herein by reference) which is described therein as "a yeast hypothetical protein". A partial alignment demonstrating the observed homology is shown immediately below.

```

>gnl|PID|e1345389 similarity with the yeast hypothetical 251.0 KD protein
YCR32W (Swiss-Prot accession number P25356); cDNA EST
EMBL:D34073
comes from this gene; cDNA EST EMBL:D32376 comes from this
gene;
cDNA EST EMBL:D37092 comes from this gene; cDNA EST EMBL>
>gnl|PID|e1346292 similarity with the yeast hypothetical 251.0
KD
protein YCR32W (Swiss-Prot accession number P25356); cDNA EST
EMBL:D34073 comes from this gene; cDNA EST EMBL:D32376 comes
from
this gene; cDNA EST EMBL:D37092 comes from this gene; cDNA EST
EMBL> >gnl|PID|e1346292 similarity with the yeast hypothetical
251.0 KD protein YCR32W (Swiss-Prot accession number P25356);
cDNA
EST EMBL:D34073 comes from this gene; cDNA EST EMBL:D32376
comes
from this gene; cDNA EST EMBL:D37092 comes from this gene;
cDNA EST
EMBL> >sp|Q19317|Q19317 F10F2.1 PROTEIN. >sp|Q20061|Q20061
F10F2.1
(FRAGMENT). (SUB 1463-2528)
Length = 2528

```

Plus Strand HSPs:

Score = 516 (181.6 bits), Expect = 4.2e-47, P = 4.2e-47
 Identities = 98/257 (38%), Positives = 161/257 (62%), Frame = +3

5 Query: 3 DKSFRVYSTETGKLTQIVFGHWDVVTCLARSESYIGGDCYIVSGSRDATLLWYWSGRHH
 182

10 D SFR+ T++G++ Q V+GH DVVTC+ARSE+ + DCY+V+GS D T++LW+W+G
 Sbjct: 2267 DYSFRIVDTDSGRVRQAVYGHGDVVTCIARSETSLFSDCYVVTGSMDCTVVLWHWNGTGT
 2326

15 Query: 183 II-GDNPNSSDYPAPRAVLTHGDHEVVCVSVCAELGLVISGAKEGPCLVHTITGDLLRAL
 359

 I G+ + P+PR++LTGH+ + + V AE GLV+SG ++G L+HT + DLLR +
 Sbjct: 2327 FIAGEYNQPGEVPSPRSILTGHEASISALCVSAEHGLVVSGCEDGVILIHTTSSDLLRRI
 2386

20 Query: 360 EGPENCLFPRLISVSSEGHCIYYERGRFSNFSINGKLLAQMEINDSTRAILLSSDGQN
 539

 G + + +L S+S E + ++ R +S K L ++ ++D + ++ DG+
 Sbjct: 2387 RG--HGIVTQL-SMSRECILLSLFDKRMVTYSATAKKLDEVLDVKIECVTVTRDGEFA
 2443

25 Query: 540 VTGGDNGVVEVWQACDFKQLYIYPGCDAGIRAMDLSHDQRTLITGMASGSIVAFNIDFNR
 719

 VTG NG + +W+ +LY Y ++ +R++ + R ++ G+ SG+IV FN DFNR
 Sbjct: 2444 VTGAVNGRINIWRMFPLTKLYTYQPLNSAVRSVAVVASHRFILGGLDSGAIVVFNADFN
 2503

30 Query: 720 WHYEHQNRVYR*RKNQKP 773
 WHYE+++RY + KP
 Sbjct: 2504 WHYEYKHRYIQNTSAAKP 2521

35 The segment of gnlIPIDle1345389 that is shown as "Sbjct" above is set out as
 sequence(s) SEQ ID NO. 125. Based on the structural similarity these homologous
 polypeptides are expected to share at least some biological activities. Such activities
 are known in the art and described elsewhere herein. Assays for determining such
 activities are also known in the art, some of which have been described elsewhere
 40 herein. Preferred polypeptides of the invention comprise polypeptides having the
 amino acid sequence(s) set out as SEQ ID NO. 126 which correspond to the Query
 sequence in the alignment shown above (gaps introduced in the sequence by the
 computer are, of course, removed).

45 It has been discovered that this gene is expressed primarily in the following
 tissues: Soares infant brain 1NIB and to a lesser extent in
 Soares_total_fetus_Nb2HF8_9w, Temporal cortex-Alzheimer; subtracted, Human
 Brain, Striatum, human ovarian cancer, Human Adrenal Gland Tumor, Human
 Amygdala, Nine Week Old Early Stage Human, Human Cerebellum.

The tissue distribution in neural tissues suggests the protein product of this clone is useful for the detection, treatment, and/or prevention of neurodegenerative disease states, behavioral disorders, or inflammatory conditions which include, but are not limited to Alzheimer's Disease, Parkinson's Disease, Huntington's Disease, 5 Tourette Syndrome, meningitis, encephalitis, demyelinating diseases, peripheral neuropathies, neoplasia, trauma, congenital malformations, spinal cord injuries, ischemia and infarction, aneurysms, hemorrhages, schizophrenia, mania, dementia, paranoia, obsessive compulsive disorder, depression, panic disorder, learning disabilities, ALS, psychoses, autism, and altered behaviors, including disorders in 10 feeding, sleep patterns, balance, and perception. In addition, elevated expression of this gene product in regions of the brain suggests it plays a role in normal neural function. Potentially, this gene product is involved in synapse formation, neurotransmission, learning, cognition, homeostasis, or neuronal differentiation or survival. Protein, as well as, antibodies directed against the protein may show utility 15 as a tumor marker and/or immunotherapy targets for the above listed tissues.

Many polynucleotide sequences, such as EST sequences, are publicly available and accessible through sequence databases. Some of these sequences are related to SEQ ID NO:19 and may have been publicly available prior to conception of the present invention. Preferably, such related polynucleotides are specifically 20 excluded from the scope of the present invention. To list every related sequence would be cumbersome. Accordingly, preferably excluded from the present invention are one or more polynucleotides comprising a nucleotide sequence described by the general formula of a-b, where a is any integer between 1 to 1151 of SEQ ID NO:19, b is an integer of 15 to 1165, where both a and b correspond to the positions of 25 nucleotide residues shown in SEQ ID NO:19, and where b is greater than or equal to a + 14.

FEATURES OF PROTEIN ENCODED BY GENE NO: 10

It has been discovered that this gene is expressed primarily in Human 30 Cerebellum..

Many polynucleotide sequences, such as EST sequences, are publicly available and accessible through sequence databases. Some of these sequences are related to SEQ ID NO:20 and may have been publicly available prior to conception of the present invention. Preferably, such related polynucleotides are specifically excluded from the scope of the present invention. To list every related sequence would be cumbersome. Accordingly, preferably excluded from the present invention are one or more polynucleotides comprising a nucleotide sequence described by the general formula of a-b, where a is any integer between 1 to 2266 of SEQ ID NO:20, b is an integer of 15 to 2280, where both a and b correspond to the positions of nucleotide residues shown in SEQ ID NO:20, and where b is greater than or equal to a + 14.

FEATURES OF PROTEIN ENCODED BY GENE NO: 11

It has been discovered that this gene is expressed primarily in the following tissues: Human Chondrosarcoma and to a lesser extent in Human Osteosarcoma, Bone marrow, Hodgkin's Lymphoma II.

Preferred epitopes include those comprising a sequence shown in SEQ ID NO. 68 as residues: Lys-22 to His-27, Ser-31 to Gly-38.

Many polynucleotide sequences, such as EST sequences, are publicly available and accessible through sequence databases. Some of these sequences are related to SEQ ID NO:21 and may have been publicly available prior to conception of the present invention. Preferably, such related polynucleotides are specifically excluded from the scope of the present invention. To list every related sequence would be cumbersome. Accordingly, preferably excluded from the present invention are one or more polynucleotides comprising a nucleotide sequence described by the general formula of a-b, where a is any integer between 1 to 663 of SEQ ID NO:21, b is an integer of 15 to 677, where both a and b correspond to the positions of nucleotide residues shown in SEQ ID NO:21, and where b is greater than or equal to a + 14.

FEATURES OF PROTEIN ENCODED BY GENE NO: 12

It has been discovered that this gene is expressed primarily in the following tissues: Human Chondrosarcoma, NTERA2 teratocarcinoma cell line+retinoic acid (14 days), Human Osteoclastoma.

Many polynucleotide sequences, such as EST sequences, are publicly
5 available and accessible through sequence databases. Some of these sequences are related to SEQ ID NO:22 and may have been publicly available prior to conception of the present invention. Preferably, such related polynucleotides are specifically excluded from the scope of the present invention. To list every related sequence would be cumbersome. Accordingly, preferably excluded from the present invention
10 are one or more polynucleotides comprising a nucleotide sequence described by the general formula of a-b, where a is any integer between 1 to 844 of SEQ ID NO:22, b is an integer of 15 to 858, where both a and b correspond to the positions of nucleotide residues shown in SEQ ID NO:22, and where b is greater than or equal to a + 14.

15

FEATURES OF PROTEIN ENCODED BY GENE NO: 13

It has been discovered that this gene is expressed primarily in the following tissues: Human Chondrosarcoma and to a lesser extent in Jurkat T-cell G1 phase, Soares breast 2NbHBst, Primary Dendritic cells,frac 2, Soares infant brain 1NIB.

20 Many polynucleotide sequences, such as EST sequences, are publicly available and accessible through sequence databases. Some of these sequences are related to SEQ ID NO:23 and may have been publicly available prior to conception of the present invention. Preferably, such related polynucleotides are specifically excluded from the scope of the present invention. To list every related sequence
25 would be cumbersome. Accordingly, preferably excluded from the present invention are one or more polynucleotides comprising a nucleotide sequence described by the general formula of a-b, where a is any integer between 1 to 1071 of SEQ ID NO:23, b is an integer of 15 to 1085, where both a and b correspond to the positions of nucleotide residues shown in SEQ ID NO:23, and where b is greater than or equal to a
30 + 14.

FEATURES OF PROTEIN ENCODED BY GENE NO: 14

It has been discovered that this gene is expressed primarily in the following tissues: Human fetal brain (TFujiwara) and to a lesser extent in Soares infant brain 1NIB, NCI_CGAP_GCB1, Soares_NbHFB, Stratagene endothelial cell 937223,

5 Pancreas Islet Cell Tumor, Soares_placenta_8to9weeks_2NbHP8to9W, HEL cell line, Stratagene HeLa cell s3 937216, Human endometrial stromal cells-treated with estradiol, Human Osteosarcoma, TF-1 Cell Line GM-CSF Treated, HM3, Human Pancreas Tumor; Reexcision, Soares_pregnant_uterus_NbHPU, Stratagene colon (#937204), Human Chondrosarcoma, Stratagene lung (#937210), NCI_CGAP_Lu5,

10 NCI_CGAP_HSC1, NCI_CGAP_Kid5, Soares breast 3NbHBst, Soares_multiple_sclerosis_2NbHMSP, Human Fetal Lung III, Osteoblasts, Soares_placenta Nb2HP, Soares_fetal_heart_NbHH19W, Soares_placenta_8to9weeks_2NbHP8to9W, Primary Dendritic Cells, lib 1, Saos2, Dexamethosome Treated, Human Fetal Lung, Human Fetal Brain, Smooth Muscle

15 Serum Treated, Norm, H. Epididymus, caput & corpus, Human Primary Breast Cancer, Human Lung, Early Stage Human Lung, subtracted, Soares_fetal_heart_NbHH19W, Apoptotic T-cell, re-excision, B Cell lymphoma, Stratagene pancreas (#937208), Stratagene ovarian cancer (#937219), Human endometrial stromal cells-treated with progesterone, HL-60, PMA 4H, re-excision,

20 wilm's tumor, Human Manic Depression Tissue, Fetal Liver, subtraction II, Human Prostate, Soares_NFL_T_GBC_S1, Breast Cancer Cell line, angiogenic, Human Primary Breast Cancer Reexcision, Human Ovarian Cancer Reexcision, Human Hypothalamus, Schizophrenia, Soares_testis_NHT, Soares_NFL_T_GBC_S1, Soares_fetal_heart_NbHH19W, Soares_pregnant_uterus_NbHPU, Stratagene hNT

25 neuron (#937233), Human Adipose, Bone Marrow Stromal Cell, untreated, Human Fetal Brain, NCI_CGAP_Br2, NCI_CGAP_Ew1, NCI_CGAP_GC4, NCI_CGAP_Pr4, NCI_CGAP_CLL1, NCI_CGAP_Kid3, Smooth muscle, serum induced, re-exc, Macrophage-oxLDL; re-excision, Colon Tumor, Human Substantia Nigra, Human Synovial Sarcoma, Human Testes, Reexcision, Human Placenta,

30 Human Adult Pulmonary; re-excision, NCI_CGAP_Pr9, NCI_CGAP_Pr21, Colon

Normal III, Activated T-cell(12h)/Thiouridine-re-excision, T cell helper II, Human 8 Week Whole Embryo, Human Cerebellum, Soares fetal liver spleen 1NFLS.

Many polynucleotide sequences, such as EST sequences, are publicly available and accessible through sequence databases. Some of these sequences are related to SEQ ID NO:24 and may have been publicly available prior to conception of the present invention. Preferably, such related polynucleotides are specifically excluded from the scope of the present invention. To list every related sequence would be cumbersome. Accordingly, preferably excluded from the present invention are one or more polynucleotides comprising a nucleotide sequence described by the general formula of a-b, where a is any integer between 1 to 1408 of SEQ ID NO:24, b is an integer of 15 to 1422, where both a and b correspond to the positions of nucleotide residues shown in SEQ ID NO:24, and where b is greater than or equal to a + 14.

15 **FEATURES OF PROTEIN ENCODED BY GENE NO: 15**

It has been discovered that this gene is expressed primarily in the following tissues: Human Chondrosarcoma.

Preferred epitopes include those comprising a sequence shown in SEQ ID NO. 72 as residues: Lys-23 to Lys-48.

Many polynucleotide sequences, such as EST sequences, are publicly available and accessible through sequence databases. Some of these sequences are related to SEQ ID NO:25 and may have been publicly available prior to conception of the present invention. Preferably, such related polynucleotides are specifically excluded from the scope of the present invention. To list every related sequence would be cumbersome. Accordingly, preferably excluded from the present invention are one or more polynucleotides comprising a nucleotide sequence described by the general formula of a-b, where a is any integer between 1 to 405 of SEQ ID NO:25, b is an integer of 15 to 419, where both a and b correspond to the positions of nucleotide residues shown in SEQ ID NO:25, and where b is greater than or equal to a + 14.

FEATURES OF PROTEIN ENCODED BY GENE NO: 16

It has been discovered that this gene is expressed primarily in the following tissues: Prostate, BPH, Lib 2, Duodenum, HUMAN B CELL LYMPHOMA.

Preferred epitopes include those comprising a sequence shown in SEQ ID NO.

5 73 as residues: Gln-18 to Glu-24, Ser-39 to Lys-44.

Many polynucleotide sequences, such as EST sequences, are publicly available and accessible through sequence databases. Some of these sequences are related to SEQ ID NO:26 and may have been publicly available prior to conception of the present invention. Preferably, such related polynucleotides are specifically
 10 excluded from the scope of the present invention. To list every related sequence would be cumbersome. Accordingly, preferably excluded from the present invention are one or more polynucleotides comprising a nucleotide sequence described by the general formula of a-b, where a is any integer between 1 to 1517 of SEQ ID NO:26, b is an integer of 15 to 1531, where both a and b correspond to the positions of
 15 nucleotide residues shown in SEQ ID NO:26, and where b is greater than or equal to a + 14.

FEATURES OF PROTEIN ENCODED BY GENE NO: 17

It has been discovered that this gene is expressed primarily in the following
 20 tissues: Human Chondrosarcoma and to a lesser extent in Human Fetal Lung III, Human Fetal Heart, Human Fetal Dura Mater, Nine Week Old Early Stage Human, Hodgkin's Lymphoma II, Human 8 Week Whole Embryo, Osteoblasts, Soares infant brain 1NIB, Human Osteosarcoma, Smooth muscle, serum treated, Smooth muscle, control, Soares fetal liver spleen 1NFLS, Human Rhabdomyosarcoma, Soares
 25 placenta Nb2HP, Bone Marrow Stromal Cell, untreated, Human Placenta, 12 Week Old Early Stage Human, Adipocytes, STROMAL -OSTEOCLASTOMA, Human Pancreas Tumor, Human Thymus Stromal Cells, NTERA2, control, Endothelial cells-control, Mesangial cell, frac 2, Human Microvascular Endothelial Cells, fract. A, Human Osteoclastoma Stromal Cells - unamplified, Human Synovial Sarcoma,
 30 Endothelial-induced, Morton Fetal Cochlea, Colon Tumor II, Human Fetal Bone, Smooth muscle, IL1b induced, Human Fetal Kidney, Soares melanocyte 2NbHM,

- Aorta endothelial cells + TNF- α , Amniotic Cells - TNF induced, Smooth muscle, serum induced, re-exc, Human T-Cell Lymphoma, Human Cerebellum, Amniotic Cells - Primary Culture, wilm's tumor, 12 Week Old Early Stage Human, II, Human Testes Tumor, Human Fetal Epithelium (Skin), Human endometrial stromal cells,
- 5 Human Placenta, Soares_fetal_heart_NbHH19W, Human Testes, Human Endometrial Tumor, HSA 172 Cells, Soares retina N2b4HR, Human Lung Cancer; re-excision, Human Chronic Synovitis, Synovial Fibroblasts (control), Monocyte activated, Human endometrial stromal cells-treated with progesterone, Human Hypothalamus, Schizophrenia, Human Adult Pulmonary; re-excision, Human Fetal
- 10 Heart, Differential (Fetal-Specific), Smooth muscle, control; re-excision, Human Stomach; re-excision, Human Osteoblasts II, Spinal cord, Stratagene lung (#937210), Soares breast 3NbHBst, Activated T-Cell (12hs)/Thiouridine labelled Eco, Human OB HOS treated (1 nM E2) fraction I, Human endometrial stromal cells-treated with estradiol, Human umbilical vein endothelial cells, IL-4 induced,
- 15 Soares_fetal_lung_NbHL19W, Colon Carcinoma, HSC172 cells, Early Stage Human Lung, subtracted; Human Adipose Tissue, re-excision, Stratagene endothelial cell 937223, Synovial hypoxia, human ovarian cancer, Human Ovarian Cancer Reexcision, Liver, Hepatoma, Soares_fetal_heart_NbHH19W, Human Whole Six Week Old Embryo, Human Substantia Nigra, Smooth Muscle Serum Treated, Norm,
- 20 Smooth muscle-ILb induced, Human T-cell lymphoma; re-excision, Brain Frontal Cortex, re-excision, Stratagene fetal spleen (#937205), Human Adipose, Ulcerative Colitis, Macrophage-oxLDL; re-excision, Human Testes, Reexcision, Bone marrow, Mesangial cell, frac 1, Human Umbilical Vein Endothelial Cells, fract. A, Stomach cancer (human); re-excision, Human Hypothalamus, schizophrenia, re-excision,
- 25 Healing groin wound; 7.5 hours post incision, NTERA2 + retinoic acid, 14 days, Stratagene endothelial cell 937223, Spinal Cord, re-excision, Stromal cell TF274, Macrophage-oxLDL, normalized infant brain cDNA, Stratagene endothelial cell 937223, Hepatocellular Tumor; re-excision, Soares_fetal_heart_NbHH19W, Soares_fetal_heart_NbHH19W, Brain frontal cortex, H Macrophage (GM-CSF
- 30 treated), re-excision, Soares_placenta_8to9weeks_2NbHP8to9W, Human Osteoclastoma, Soares_fetal_heart_NbHH19W, Weizmann Olfactory Epithelium,

- Human OB MG63 treated (10 nM E2) fraction I, Human OB HOS treated (10 nM E2) fraction I, Human Aortic Endothelium, Human Pancreatic Carcinoma, Smooth Muscle- HASTE normalized, Stratagene fetal retina 937202, Human Ovary, Prostate BPH, Human Infant Brain, Fetal Liver, subtraction II, Human Dermal Endothelial
- 5 Cells, untreated, Human Hippocampus, Stratagene endothelial cell 937223, Hemangiopericytoma, Colon Normal II, Human Fetal Kidney; Reexcision, Soares_pregnant_uterus_NbHPU, Pharynx carcinoma, Human Umbilical Vein Endothelial cells, frac B, re-excision, Human Uterus, normal, CHME Cell Line; treated 5 hrs, Soares_fetal_lung_NbHL19W, stomach cancer (human), H.
- 10 Atrophic Endometrium, Hodgkin's Lymphoma I, Adipocytes; re-excision, Human Fetal Spleen, HUMAN STOMACH, Soares_fetal_heart_NbHH19W, Human Adult Heart; re-excision, NTERA2 teratocarcinoma cell line+retinoic acid (14 days), Human Synovium, Stratagene placenta (#937225), human corpus colosum, H Female Bladder, Adult, H. Kidney Cortex, subtracted, Glioblastoma, Human Frontal Cortex,
- 15 Schizophrenia, H. Meningima, M1, Human Brain, Striatum, Merkel Cells, Soares_senescent_fibroblasts_NbHSF, Soares adult brain N2b5HB55Y, 12 Week Early Stage Human II; Reexcision, Soares breast 2NbHBst, Human Adrenal Gland Tumor, Rejected Kidney, lib 4, H. Frontal cortex, epileptic; re-excision, Soares_senescent_fibroblasts_NbHSF, Soares_multiple_sclerosis_2NbHMSP, Colon
- 20 Normal III, Stratagene endothelial cell 937223, Human Hippocampus, Human Fetal Heart, subtracted, Brain pons, Human Adult Spleen, fraction II, Human Macrophage, subtracted, Human Fetal Liver- Enzyme subtraction, Human Placenta, subtracted, Human Infant Adrenal Gland, Human Microvascular Endothelial Cells, fract. B, H Umbilical Vein Endothelial Cells, frac A, re-excision, Whole 6 Week Old Embryo,
- 25 prostate-edited, Human Fetal Brain, Human Cerebellum, subtracted, Human Placenta, H. cerebellum, Enzyme subtracted, Stratagene fibroblast (#937212), Human Skin Tumor, Human Pineal Gland, Stratagene fetal retina 937202, Stratagene neuroepithelium (#937231), Human Umbilical Vein, Endo. remake, Healing groin wound, 6.5 hours post incision, Human Osteoclastoma, re-excision, Stratagene
- 30 muscle 937209, Stratagene neuroepithelium (#937231), Human Adult Small Intestine, Human Primary Breast Cancer Reexcision, Human Adult Testes, Large Inserts,

- Reexcision, Human Heart, Soares_pregnant_uterus_NbHPU, Stratagene fibroblast (#937212), Soares_fetal_heart_NbHH19W, Stratagene liver (#937224), Soares_pregnant_uterus_NbHPU, Human Amygdala, Soares_fetal_heart_NbHH19W, HUMAN B CELL LYMPHOMA, Human Bone Marrow, treated, Human
- 5 Hippocampus, prescreened, Human Fetal Kidney, Smooth muscle-edited A, H Amygdala Depression, subtracted, Human Primary Breast Cancer, Human Fetal Kidney, Human Spleen, Human Normal Cartilage Fraction III, Human Umbilical Vein Endothelial Cells, fract. B, CHME Cell Line, untreated, Healing Abdomen Wound; 15 days post incision, Thyroid Thyroiditis, Brain; normal, Placenta, Human
- 10 Lung Cancer, subtracted, Prostate BPH, Lib 2, subtracted, Rectum tumour, Human Adult Heart, subtracted, Human Amygdala Depression, re-excision, Tongue carcinoma, Human Brain, Ea.hy.926 cell line, Thyroid Normal (SDCA2 No), Human Adult Heart, brain stem, Brain Amygdala Depression, Human Prostate Cancer, Stage B2, H. Meningioma, M6, Human Colon Cancer, subtracted, Human Fetal Lung,
- 15 Human epithelioid sarcoma, Saos2 Cells; Untreated, Human OB MG63 control fraction I, Dermatofibrosarcoma Protuberance, Human Fetal Brain, random primed, Human Colon, subtraction, H. Striatum Depression, subt, Human Gall Bladder, fraction II, Human Adult Spleen, Fetal Heart, Human OB HOS control fraction I, Human White Adipose, Human Primary Breast Cancer; re-excision, Human
- 20 Cardiomyopathy, subtracted, Soares_fetal_lung_NbHL19W, Soares_parathyroid_tumor_NbHPA, Soares_senescent_fibroblasts_NbHSF, Healing Abdomen wound; 70&90 min post incision, H. Epididymus, caput & corpus, Human Primary Breast Cancer, Human Thyroid, Human Lung, Soares_pregnant_uterus_NbHPU, Stratagene corneal stroma (#937222), Human adult
- 25 small intestine, re-excision, Human Amygdala; re-excision, H. Whole Brain #2, re-excision, Ovarian Tumor 10-3-95, Human Epididymus, pBMC stimulated w/ poly I/C, Synovial hypoxia-RSF subtracted, Human Colon; re-excision, Synovial Fibroblasts (Il1/TNF), subt, Spleen metastatic melanoma, H. Lymph node breast Cancer, KMH2, Human Prostate, Human Bone Marrow, re-excision, HUMAN
- 30 JURKAT MEMBRANE BOUND POLYSOMES, Human Uterine Cancer, Soares_parathyroid_tumor_NbHPA, Soares_fetal_heart_NbHH19W, Stratagene

muscle 937209, Stratagene hNT neuron (#937233), Human Thymus, Human Fetal Brain, Macrophage (GM-CSF treated), Stratagene NT2 neuronal precursor 937230, Human Liver, normal, Pancreas Islet Cell Tumor, Human Gall Bladder, Colon Tumor, breast lymph node CDNA library, Early Stage Human Brain, Soares_NbHFB, 5 Soares_NhHMPu_S1, human tonsils, Human fetal heart, Lambda ZAP Express, NCI_CGAP_PNS1, Stratagene fibroblast (#937212), Soares_fetal_lung_NbHL19W, Soares_fetal_liver_spleen_1NFLS_S1, Soares_senescent_fibroblasts_NbHSF, Spleen, Chronic lymphocytic leukemia, Stratagene muscle 937209, Stratagene fibroblast (#937212), Keratinocyte, Soares_fetal_lung_NbHL19W, 10 Soares_multiple_sclerosis_2NbHMSP, Soares_senescent_fibroblasts_NbHSF.

Many polynucleotide sequences, such as EST sequences, are publicly available and accessible through sequence databases. Some of these sequences are related to SEQ ID NO:27 and may have been publicly available prior to conception of the present invention. Preferably, such related polynucleotides are specifically 15 excluded from the scope of the present invention. To list every related sequence would be cumbersome. Accordingly, preferably excluded from the present invention are one or more polynucleotides comprising a nucleotide sequence described by the general formula of a-b, where a is any integer between 1 to 943 of SEQ ID NO:27, b is an integer of 15 to 957, where both a and b correspond to the positions of 20 nucleotide residues shown in SEQ ID NO:27, and where b is greater than or equal to a + 14.

FEATURES OF PROTEIN ENCODED BY GENE NO: 18

It has been discovered that this gene is expressed primarily in the following 25 tissues: Soares fetal liver spleen 1NFLS and to a lesser extent in Soares breast 2NbHBst, Soares_NhHMPu_S1, Soares breast 3NbHBst, Soares_fetal_lung_NbHL19W, Soares_multiple_sclerosis_2NbHMSP, human colon cancer, Human Lung Cancer;re-excision, Jurkat T-Cell, S phase, wilm's tumor, T Cell helper I, Soares_fetal_heart_NbHH19W, Bone Marrow Cell Line (RS4;11), 30 Hodgkin's Lymphoma II, Nine Week Old Early Stage Human, Human Cerebellum, Soares placenta Nb2HP.

Many polynucleotide sequences, such as EST sequences, are publicly available and accessible through sequence databases. Some of these sequences are related to SEQ ID NO:28 and may have been publicly available prior to conception of the present invention. Preferably, such related polynucleotides are specifically excluded from the scope of the present invention. To list every related sequence would be cumbersome. Accordingly, preferably excluded from the present invention are one or more polynucleotides comprising a nucleotide sequence described by the general formula of a-b, where a is any integer between 1 to 1376 of SEQ ID NO:28, b is an integer of 15 to 1390, where both a and b correspond to the positions of nucleotide residues shown in SEQ ID NO:28, and where b is greater than or equal to a + 14.

FEATURES OF PROTEIN ENCODED BY GENE NO: 19

It has been discovered that this gene is expressed primarily in the following tissues: Soares infant brain 1NIB and to a lesser extent in Human Cerebellum, Soares_pregnant_uterus_NbHPU, Soares fetal liver spleen 1NFLS, Human 8 Week Whole Embryo, subtracted, Human Placenta, Human Tonsils, Lib 2, Healing groin wound; 7.5 hours post incision, Soares_NhHMPu_S1, Hepatocellular Tumor; re-excision, Activated T-Cell (12hs)/Thiouridine labelledEco, Activated T-cell(12h)/Thiouridine-re-excision, Spleen, Chronic lymphocytic leukemia, Hodgkin's Lymphoma II, Primary Dendritic Cells, lib 1.

Many polynucleotide sequences, such as EST sequences, are publicly available and accessible through sequence databases. Some of these sequences are related to SEQ ID NO:29 and may have been publicly available prior to conception of the present invention. Preferably, such related polynucleotides are specifically excluded from the scope of the present invention. To list every related sequence would be cumbersome. Accordingly, preferably excluded from the present invention are one or more polynucleotides comprising a nucleotide sequence described by the general formula of a-b, where a is any integer between 1 to 1576 of SEQ ID NO:29, b is an integer of 15 to 1590, where both a and b correspond to the positions of

nucleotide residues shown in SEQ ID NO:29, and where b is greater than or equal to a + 14.

FEATURES OF PROTEIN ENCODED BY GENE NO: 20

5 The computer algorithm BLASTX has been used to determine that the translation product of this gene shares sequence homology with, as a non-limiting example, the sequence accessible through the following database accession no. gil1791257 (all information available through the recited accession number is incorporated herein by reference) which is described therein as "copine I [Homo sapiens]". A partial alignment demonstrating the observed homology is shown immediately below.

```

>gi|1791257 copine I [Homo sapiens]
      Length = 537

      Plus Strand HSPs:

      Score = 432 (152.1 bits), Expect = 5.2e-120, Sum P(4) = 5.2e-120
      Identities = 77/126 (61%), Positives = 98/126 (77%), Frame = +3

      Query:   519 PKYRDKKKKNYKSSGTVVLAQCTVEKVHTFLDYIMGGCQISFTVAIDFTASNGDPRSSQSL
      698
      P+ + KKK+YK+SGT+ + C VE ++FLDY+MGGCQI+FTV +DFT SNGDP S SL
      Sbjct:   246 PEKQQKKKSYKNSGTIRVKICRVETESFLDYVMGGCQINFVTGVDFGTSGNDPSSPDSL
      305

      Query:   699 HCLSPRQPNHYLQALRAVGGICQDYDSDKRFPAPFGGARIPPNFEVSHDFAINFDPENPE
      878
      H LSP N YL AL +VG + QDYDSDK FPAFGFGA++PP+++VSH+FA+NF+P NP
      Sbjct:   306 HYLSP TGVNEYLMALWSVGSVVQDYDSDKLFPAPFGFGAQVPPDWQVSHEFALNFNPSNPY
      365

      Query:   879 CEESQG 896
      C QG
      Sbjct:   366 CAGIQG 371

      Score = 394 (138.7 bits), Expect = 5.2e-120, Sum P(4) = 5.2e-120
      Identities = 82/140 (58%), Positives = 100/140 (71%), Frame = +1

      Query:   931 QLYGPTNVAPIINRVAEPAQREQSTGQATKYSVLLVLTGQVSDMAETRTAIVRASRLPM
      1110
      +LYGPTN APIIN VA A + G A++Y +LL+LTDG V+D+ TR A+VRAS LPM
      Sbjct:   384 RLYGPTNFAPIINHVARFAAQAAHQGTASQYFMLLLLTDGAVTDVEATREAVVRASNLPM
      443

      Query:   1111 SIIIVGVGNADFSDMRLLDGDDGPLRCPRGVPAARDIVQFVPRDFKDAAPSALAKCVLA
      1290
      S+IIVGVG ADF M LD D GPL G AARDIVQFVP+R F++A ALA+ VLA
      Sbjct:   444 SVIIVGVGGADFEAMEQLDADGGPLHTRSGQAAARDIVQFVPYRRFQNAPREALAQTVLA
      503

      Query:   1291 EVPRQVVEYYASQGISPGAP 1350
      EVP Q+V Y+ +QG +P P
  
```

5 Sbjct: 504 EVPTQLVSYFRAQGWAPLKP 523
 Score = 352 (123.9 bits), Expect = 5.2e-120, Sum P(4) = 5.2e-120
 Identities = 74/184 (40%), Positives = 109/184 (59%), Frame = +1
 10 Query: 10 RTTSPRNDTFLGSTECTLGQIVSQTKVTKPLLLKNGKTAGKSTITIVAEVSGTNDYVQL
 189 +T R+D FLG EC+LGQIVS +T PL+LK GK AG+ TIT+ A+E+ N V +
 Sbjct: 84 KTPELRDDDFLGGAECSLGQIVSSQVLTLPMLKPGKPAGRGTTITVSAQELKD-NRVVTM
 142
 15 Query: 190 TFRAYKLDNKDLFSKSDPFMEIYKTNEDQSDQLVWRTEVVKNLNPSWEPPRLSLHSLCS
 369 A LD KD KSDPF+E ++ D LV+R+EV+KNNLNP+W+ F + + C
 Sbjct: 143 EVEARNLDKKDFLGKSDPFLEFFRQG-DGKWHLVYRSEVIKNNLNPTWKRFSVPVQHFCG
 201
 20 Query: 370 CDVHRPLKFLVYDYDSSGKHDFIGEFSTTFQEMQEGTANPGQEMQWDCINPSIGTRRRIT
 549 + P++ DYDS G HD IG F ++ ++Q A P + ++CI+P +++
 Sbjct: 202 GNPSTPIQVQCSYDSDGSHDLIGTFTSLAQLQ---AVPAE---FECIHPEKQKKKSY
 255
 25 Query: 550 RAQG 561
 + G
 Sbjct: 256 KNSG 259

The segment of gil1791257 that is shown as "Sbjct" above is set out as
 sequence(s) SEQ ID NO. 127, SEQ ID NO. 129, and SEQ ID NO. 131. Based on the
 30 structural similarity these homologous polypeptides are expected to share at least
 some biological activities. Such activities are known in the art and described
 elsewhere herein. Assays for determining such activities are also known in the art,
 some of which have been described elsewhere herein. Preferred polypeptides of the
 invention comprise polypeptides having the amino acid sequence(s) set out as SEQ ID
 35 NO. 128, SEQ ID NO. 130, and SEQ ID NO. 132 which correspond to the Query
 sequence in the alignment shown above (gaps introduced in the sequence by the
 computer are, of course, removed).

It has been discovered that this gene is expressed primarily in the following
 tissues: Soares infant brain 1NIB and to a lesser extent in Human Cerebellum, Human
 40 Fetal Brain, Infant brain, Bento Soares, Frontal lobe, dementia; re-excision, Human
 Whole Brain #2 - Oligo dT > 1.5Kb, Human Infant Brain, Human
 Hypothalamus, Schizophrenia, Pancreas Islet Cell Tumor, Human Amygdala.

Preferred epitopes include those comprising a sequence shown in SEQ ID NO.
 77 as residues: Arg-30 to Gly-45, Ser-52 to Ser-65, Gly-101 to Pro-106.

The tissue distribution in neural tissues suggests the protein product of this clone is useful for the detection, treatment, and/or prevention of neurodegenerative disease states, behavioral disorders, or inflammatory conditions which include, but are not limited to Alzheimer's Disease, Parkinson's Disease, Huntington's Disease, 5 Tourette Syndrome, meningitis, encephalitis, demyelinating diseases, peripheral neuropathies, neoplasia, trauma, congenital malformations, spinal cord injuries, ischemia and infarction, aneurysms, hemorrhages, schizophrenia, mania, dementia, paranoia, obsessive compulsive disorder, depression, panic disorder, learning disabilities, ALS, psychoses, autism, and altered behaviors, including disorders in 10 feeding, sleep patterns, balance, and perception. In addition, elevated expression of this gene product in regions of the brain suggests it plays a role in normal neural function. Potentially, this gene product is involved in synapse formation, neurotransmission, learning, cognition, homeostasis, or neuronal differentiation or survival.

15 Moreover, the expression within fetal tissue and other cellular sources marked by proliferating cells suggests this protein may play a role in the regulation of cellular division, and may show utility in the diagnosis and treatment of cancer and other proliferative disorders. Similarly, developmental tissues rely on decisions involving cell differentiation and/or apoptosis in pattern formation. Dysregulation of apoptosis 20 can result in inappropriate suppression of cell death, as occurs in the development of some cancers, or in failure to control the extent of cell death, as is believed to occur in acquired immunodeficiency and certain neurodegenerative disorders, such as spinal muscular atrophy (SMA). Therefore, the polynucleotides and polypeptides of the present invention are useful in treating, detecting, and/or preventing said disorders 25 and conditions, in addition to other types of degenerative conditions. Thus this protein may modulate apoptosis or tissue differentiation and would be useful in the detection, treatment, and/or prevention of degenerative or proliferative conditions and diseases. Protein, as well as, antibodies directed against the protein may show utility as a tumor marker and/or immunotherapy targets for the above listed tissues.

30 Many polynucleotide sequences, such as EST sequences, are publicly available and accessible through sequence databases. Some of these sequences are

related to SEQ ID NO:30 and may have been publicly available prior to conception of the present invention. Preferably, such related polynucleotides are specifically excluded from the scope of the present invention. To list every related sequence would be cumbersome. Accordingly, preferably excluded from the present invention are one or more polynucleotides comprising a nucleotide sequence described by the general formula of a-b, where a is any integer between 1 to 1637 of SEQ ID NO:30, b is an integer of 15 to 1651, where both a and b correspond to the positions of nucleotide residues shown in SEQ ID NO:30, and where b is greater than or equal to a + 14.

10

FEATURES OF PROTEIN ENCODED BY GENE NO: 21

It has been discovered that this gene is expressed primarily in the following tissues: Human Neutrophils, Activated, re-excision, Human Cerebellum.

The tissue distribution in neural tissues suggests the protein product of this clone is useful for the detection, treatment, and/or prevention of neurodegenerative disease states, behavioral disorders, or inflammatory conditions which include, but are not limited to Alzheimer's Disease, Parkinson's Disease, Huntington's Disease, Tourette Syndrome, meningitis, encephalitis, demyelinating diseases, peripheral neuropathies, neoplasia, trauma, congenital malformations, spinal cord injuries, ischemia and infarction, aneurysms, hemorrhages, schizophrenia, mania, dementia, paranoia, obsessive compulsive disorder, depression, panic disorder, learning disabilities, ALS, psychoses, autism, and altered behaviors, including disorders in feeding, sleep patterns, balance, and perception. In addition, elevated expression of this gene product in regions of the brain suggests it plays a role in normal neural function. Potentially, this gene product is involved in synapse formation, neurotransmission, learning, cognition, homeostasis, or neuronal differentiation or survival.

Moreover, the expression of this gene product suggests a role in regulating the proliferation; survival; differentiation; and/or activation of hematopoietic cell lineages, including blood stem cells. This gene product may be involved in the regulation of cytokine production, antigen presentation, or other processes suggesting

a usefulness in the treatment of cancer (e.g. by boosting immune responses). Since the gene is expressed in cells of lymphoid origin, the natural gene product may be involved in immune functions. Therefore it may be also used as an agent for immunological disorders including arthritis, asthma, immunodeficiency diseases such as AIDS, leukemia, rheumatoid arthritis, granulomatous disease, inflammatory bowel disease, sepsis, acne, neutropenia, neutrophilia, psoriasis, hypersensitivities, such as T-cell mediated cytotoxicity; immune reactions to transplanted organs and tissues, such as host-versus-graft and graft-versus-host diseases, or autoimmunity disorders, such as autoimmune infertility, lense tissue injury, demyelination, systemic lupus erythematosus, drug induced hemolytic anemia, rheumatoid arthritis, Sjogren's disease, scleroderma and tissues. Moreover, the protein may represent a secreted factor that influences the differentiation or behavior of other blood cells, or that recruits hematopoietic cells to sites of injury. In addition, this gene product may have commercial utility in the expansion of stem cells and committed progenitors of various blood lineages, and in the differentiation and/or proliferation of various cell types. Protein, as well as, antibodies directed against the protein may show utility as a tumor marker and/or immunotherapy targets for the above listed tissues.

Many polynucleotide sequences, such as EST sequences, are publicly available and accessible through sequence databases. Some of these sequences are related to SEQ ID NO:31 and may have been publicly available prior to conception of the present invention. Preferably, such related polynucleotides are specifically excluded from the scope of the present invention. To list every related sequence would be cumbersome. Accordingly, preferably excluded from the present invention are one or more polynucleotides comprising a nucleotide sequence described by the general formula of a-b, where a is any integer between 1 to 1706 of SEQ ID NO:31, b is an integer of 15 to 1720, where both a and b correspond to the positions of nucleotide residues shown in SEQ ID NO:31, and where b is greater than or equal to a + 14.

30 FEATURES OF PROTEIN ENCODED BY GENE NO: 22

It has been discovered that this gene is expressed primarily in the following tissues: Human Rhabdomyosarcoma, Human Cerebellum.

Many polynucleotide sequences, such as EST sequences, are publicly available and accessible through sequence databases. Some of these sequences are related to SEQ ID NO:32 and may have been publicly available prior to conception of the present invention. Preferably, such related polynucleotides are specifically excluded from the scope of the present invention. To list every related sequence would be cumbersome. Accordingly, preferably excluded from the present invention are one or more polynucleotides comprising a nucleotide sequence described by the general formula of a-b, where a is any integer between 1 to 1233 of SEQ ID NO:32, b is an integer of 15 to 1247, where both a and b correspond to the positions of nucleotide residues shown in SEQ ID NO:32, and where b is greater than or equal to a + 14.

15 **FEATURES OF PROTEIN ENCODED BY GENE NO: 23**

It has been discovered that this gene is expressed primarily in the following tissues: HeLa cell line, Human Whole Six Week Old Embryo, Human Cerebellum.

Preferred epitopes include those comprising a sequence shown in SEQ ID NO. 80 as residues: Asp-3 to Ser-11, Ser-14 to Ser-21, Pro-23 to Glu-37, Ser-85 to Asn-90.

Many polynucleotide sequences, such as EST sequences, are publicly available and accessible through sequence databases. Some of these sequences are related to SEQ ID NO:33 and may have been publicly available prior to conception of the present invention. Preferably, such related polynucleotides are specifically excluded from the scope of the present invention. To list every related sequence would be cumbersome. Accordingly, preferably excluded from the present invention are one or more polynucleotides comprising a nucleotide sequence described by the general formula of a-b, where a is any integer between 1 to 963 of SEQ ID NO:33, b is an integer of 15 to 977, where both a and b correspond to the positions of nucleotide residues shown in SEQ ID NO:33, and where b is greater than or equal to a + 14.

FEATURES OF PROTEIN ENCODED BY GENE NO: 24

It has been discovered that this gene is expressed primarily in Human Cerebellum..

5 Preferred epitopes include those comprising a sequence shown in SEQ ID NO. 81 as residues: Thr-31 to Pro-40.

Many polynucleotide sequences, such as EST sequences, are publicly available and accessible through sequence databases. Some of these sequences are related to SEQ ID NO:34 and may have been publicly available prior to conception of
10 the present invention. Preferably, such related polynucleotides are specifically excluded from the scope of the present invention. To list every related sequence would be cumbersome. Accordingly, preferably excluded from the present invention are one or more polynucleotides comprising a nucleotide sequence described by the general formula of a-b, where a is any integer between 1 to 1879 of SEQ ID NO:34, b
15 is an integer of 15 to 1893, where both a and b correspond to the positions of nucleotide residues shown in SEQ ID NO:34, and where b is greater than or equal to a + 14.

FEATURES OF PROTEIN ENCODED BY GENE NO: 25

20 It has been discovered that this gene is expressed primarily in the following tissues: Early Stage Human Brain, Human Cerebellum.

Many polynucleotide sequences, such as EST sequences, are publicly available and accessible through sequence databases. Some of these sequences are related to SEQ ID NO:35 and may have been publicly available prior to conception of
25 the present invention. Preferably, such related polynucleotides are specifically excluded from the scope of the present invention. To list every related sequence would be cumbersome. Accordingly, preferably excluded from the present invention are one or more polynucleotides comprising a nucleotide sequence described by the general formula of a-b, where a is any integer between 1 to 833 of SEQ ID NO:35, b
30 is an integer of 15 to 847, where both a and b correspond to the positions of

nucleotide residues shown in SEQ ID NO:35, and where b is greater than or equal to a + 14.

FEATURES OF PROTEIN ENCODED BY GENE NO: 26

5 It has been discovered that this gene is expressed primarily in the following tissues: Soares adult brain N2b5HB55Y and to a lesser extent in Early Stage Human Brain, Human Cerebellum, Human Cerebellum, subtracted, Human Hypothalamus, schizophrenia, re-excision, Human Whole Brain #2 - Oligo dT > 1.5Kb, Human Adrenal Gland Tumor, H. Frontal cortex, epileptic; re-excision, Soares
10 infant brain 1NIB.

Preferred epitopes include those comprising a sequence shown in SEQ ID NO. 83 as residues: Gln-6 to Leu-12, Pro-55 to Gly-63.

Many polynucleotide sequences, such as EST sequences, are publicly available and accessible through sequence databases. Some of these sequences are
15 related to SEQ ID NO:36 and may have been publicly available prior to conception of the present invention. Preferably, such related polynucleotides are specifically excluded from the scope of the present invention. To list every related sequence would be cumbersome. Accordingly, preferably excluded from the present invention are one or more polynucleotides comprising a nucleotide sequence described by the
20 general formula of a-b, where a is any integer between 1 to 1086 of SEQ ID NO:36, b is an integer of 15 to 1100, where both a and b correspond to the positions of nucleotide residues shown in SEQ ID NO:36, and where b is greater than or equal to a + 14.

25 FEATURES OF PROTEIN ENCODED BY GENE NO: 27

The computer algorithm BLASTX has been used to determine that the translation product of this gene shares sequence homology with, as a non-limiting example, the sequence accessible through the following database accession no. gil2773385 (all information available through the recited accession number is
30 incorporated herein by reference) which is described therein as "(AF041059) similar

to KIAA0291: D..." A partial alignment demonstrating the observed homology is shown immediately below.

```

5  >gi|2773385 (AF041059) similar to KIAA0291: DDBJ Accession Number
   AB006629,
      restin and CLIP-170 [Homo sapiens] >sp|O43611|O43611 SIMILAR
   TO
      KIAA0291: DDBJ ACCESSION NUMBER AB006629, RESTIN AND CLIP-170
10  (FRAGMENT). >sp|G1881850|G1881850 WSCR4=RESTIN HOMOLOG (REPEAT
   HOMOLOG      REGION 2). (SUB 180-299) >sp|G1881849|G1881849 WSCR4=RESTIN
      (REPEAT REGION 1). (SUB 40-114)
      Length = 419
15  Plus Strand HSPs:
      Score = 256 (90.1 bits), Expect = 3.8e-30, Sum P(3) = 3.8e-30
      Identities = 55/124 (44%), Positives = 80/124 (64%), Frame = +1
20  Query:  1165  GSLQQRDGAK-AEVDQVLVAGQKQGIVRFYGKTDFAPGYWYGIELDQPTGKHDGVSFVG
   1341
      GS+++  G K    +GD+VLV G K G+VR+ G+TDFA G W G+ELD+P GK+DG+V G
   Sbjct:  169  GSVKR--GEKDLRLGDRVLVGGTKTGVVRYVGETDFAKGEWCGVELDEPLGKNDGAVAGT
25  226
      Query:  1342  RYFTCPPRHGVFAPASRIQRIGGSTDSPGDSVGAKKVHQTMTQPKRTFTTVRTPKDIAS
   1521
      RYF CPP+ G+FAP  ++ RIG  + SP    AKK  ++ M    T +    +    I+S
30  Sbjct:  227  RYFQCCKPKFGLFAPIHKVIKVFSTSPAK---AKKTKRMAMGVSALTHSP---SSSSISS
   281
      Query:  1522  ENSIS 1536
      +S++
   Sbjct:  282  VSSVA 286
35

```

The segment of gi2773385 that is shown as "Sbjct" above is set out as sequence(s) SEQ ID NO. 133 Based on the structural similarity these homologous polypeptides are expected to share at least some biological activities. Such activities are known in the art and described elsewhere herein. Assays for determining such activities are also known in the art, some of which have been described elsewhere herein. Preferred polypeptides of the invention comprise polypeptides having the amino acid sequence(s) set out as SEQ ID NO. 134 which correspond to the Query sequence in the alignment shown above (gaps introduced in the sequence by the computer are, of course, removed).

It has been discovered that this gene is expressed primarily in the following tissues: Human Cerebellum and to a lesser extent in Soares breast 2NbHBst, Soares infant brain 1NIB, Soares adult brain N2b5HB55Y, Early Stage Human Brain, H.

- Frontal cortex,epileptic;re-excision, Soares adult brain N2b4HB55Y, Human Brain, Striatum, Soares breast 3NbHBst, Nine Week Old Early Stage Human, Human Hippocampus, subtracted, H. Striatum Depression, subt, Human Cerebellum, subtracted, H. cerebellum, Enzyme subtracted, H. Whole Brain #2, re-excision,
- 5 Human Fetal Dura Mater, Hemangiopericytoma, Soares melanocyte 2NbHM, Human Synovial Sarcoma, Endothelial cells-control, Human Amygdala, Human Brain, striatum, re-excision, brain stem, Human 8 Week Whole Embryo, subtracted, Human Fetal Brain, Human Adult Retina, Frontal lobe,dementia;re-excision, Stratagene neuroepithelium NT2RAMI 937234, Smooth Muscle- HASTE normalized, H. Kidney
- 10 Cortex, subtracted, Human Whole Brain #2 - Oligo dT > 1.5Kb, Synovial hypoxia, Jurkat T-cell G1 phase, H. Meningioma, M1, Human Manic Depression Tissue, Brain Frontal Cortex, re-excision, Spinal Cord, re-excision, human ovarian cancer, Human Hypothalamus,Schizophrenia, Liver, Hepatoma, Spinal cord, Synovial Fibroblasts (control), Human Fetal Brain, Human Fetal Heart, Endothelial-induced, Spleen,
- 15 Chronic lymphocytic leukemia, Human Testes, Osteoblasts, Human 8 Week Whole Embryo.

The tissue distribution in neural tissues suggests the protein product of this clone is useful for the detection, treatment, and/or prevention of neurodegenerative disease states, behavioral disorders, or inflammatory conditions which include, but

20 are not limited to Alzheimer's Disease, Parkinson's Disease, Huntington's Disease, Tourette Syndrome, meningitis, encephalitis, demyelinating diseases, peripheral neuropathies, neoplasia, trauma, congenital malformations, spinal cord injuries, ischemia and infarction, aneurysms, hemorrhages, schizophrenia, mania, dementia, paranoia, obsessive compulsive disorder, depression, panic disorder, learning

25 disabilities, ALS, psychoses, autism, and altered behaviors, including disorders in feeding, sleep patterns, balance, and perception. In addition, elevated expression of this gene product in regions of the brain suggests it plays a role in normal neural function. Potentially, this gene product is involved in synapse formation, neurotransmission, learning, cognition, homeostasis, or neuronal differentiation or

30 survival. Protein, as well as, antibodies directed against the protein may show utility as a tumor marker and/or immunotherapy targets for the above listed tissues.

Many polynucleotide sequences, such as EST sequences, are publicly available and accessible through sequence databases. Some of these sequences are related to SEQ ID NO:37 and may have been publicly available prior to conception of the present invention. Preferably, such related polynucleotides are specifically excluded from the scope of the present invention. To list every related sequence would be cumbersome. Accordingly, preferably excluded from the present invention are one or more polynucleotides comprising a nucleotide sequence described by the general formula of a-b, where a is any integer between 1 to 3131 of SEQ ID NO:37, b is an integer of 15 to 3145, where both a and b correspond to the positions of nucleotide residues shown in SEQ ID NO:37, and where b is greater than or equal to a + 14.

FEATURES OF PROTEIN ENCODED BY GENE NO: 28

It has been discovered that this gene is expressed primarily in the following tissues: Soares fetal liver spleen 1NFLS and to a lesser extent in Soares_testis_NHT, Soares placenta Nb2HP, Soares_NFL_T_GBC_S1, NCI_CGAP_Co3, Human Epididymus, Human Fetal Brain, Human Synovial Sarcoma, Soares_fetal_heart_NbHH19W, Primary Dendritic Cells, lib 1, Soares_senescent_fibroblasts_NbHSF, Human Adult Retina, NCI_CGAP_GC4, NCI_CGAP_Kid5, Stratagene muscle 937209, Stratagene lung (#937210), Soares_testis_NHT, NCI_CGAP_GCB1, Human 8 Week Whole Embryo, Human Cerebellum, Stratagene colon (#937204), Sinus piniformis Tumour, K562 + PMA (36 hrs), HPAS (human pancreas, subtracted), Testes, LNCAP + 0.3nM R1881, H Umbilical Vein Endothelial Cells, frac A, re-excision, Human Colon Cancer, subtracted, Human Fetal Brain, random primed, Human Fetal Bone, Breast Lymph node cDNA library, Resting T-Cell; re-excision, NTERA2 teratocarcinoma cell line+retinoic acid (14 days), human corpus colosum, Human Whole Brain #2 - Oligo dT > 1.5Kb, Human endometrial stromal cells, Jurkat T-cell G1 phase, Human Thymus, 22 week old human fetal liver cDNA library, HM1, NCI_CGAP_Br2, NCI_CGAP_Co8, NCI_CGAP_GCB1, Monocyte activated; re-excision, Human Osteoblasts II, 12 Week Old Early Stage Human, II, Soares_NhHMPu_S1, Merkel

Cells, Ulcerative Colitis, HM3, Infant brain, Bento Soares,
 Soares_pineal_gland_N3HPG, Soares_fetal_heart_NbHH19W,
 Soares_parathyroid_tumor_NbHPA, Human Testes Tumor, re-excision, Human
 Adrenal Gland Tumor, Rejected Kidney, lib 4, Macrophage-oxLDL; re-excision,
 5 NCI_CGAP_AA1, Colon Normal II, Human Testes Tumor, H Macrophage (GM-CSF
 treated), re-excision, Human Neutrophil, Activated, Activated T-Cell
 (12hs)/Thiouridine labelledEco, Human Microvascular Endothelial Cells, fract. A,
 Smooth muscle,control, Monocyte activated, Nine Week Old Early Stage Human,
 Soares infant brain 1NIB.

10 Preferred epitopes include those comprising a sequence shown in SEQ ID NO.
 85 as residues: His-4 to Thr-9, Ala-31 to Pro-36.

Many polynucleotide sequences, such as EST sequences, are publicly
 available and accessible through sequence databases. Some of these sequences are
 related to SEQ ID NO:38 and may have been publicly available prior to conception of
 15 the present invention. Preferably, such related polynucleotides are specifically
 excluded from the scope of the present invention. To list every related sequence
 would be cumbersome. Accordingly, preferably excluded from the present invention
 are one or more polynucleotides comprising a nucleotide sequence described by the
 general formula of a-b, where a is any integer between 1 to 826 of SEQ ID NO:38, b
 20 is an integer of 15 to 840, where both a and b correspond to the positions of
 nucleotide residues shown in SEQ ID NO:38, and where b is greater than or equal to a
 + 14.

FEATURES OF PROTEIN ENCODED BY GENE NO: 29

25 It has been discovered that this gene is expressed primarily in the following
 tissues: Spleen, Chronic lymphocytic leukemia, Human Cerebellum.

Preferred epitopes include those comprising a sequence shown in SEQ ID NO.
 86 as residues: Tyr-21 to Lys-30.

Many polynucleotide sequences, such as EST sequences, are publicly
 30 available and accessible through sequence databases. Some of these sequences are
 related to SEQ ID NO:39 and may have been publicly available prior to conception of

the present invention. Preferably, such related polynucleotides are specifically excluded from the scope of the present invention. To list every related sequence would be cumbersome. Accordingly, preferably excluded from the present invention are one or more polynucleotides comprising a nucleotide sequence described by the
5 general formula of a-b, where a is any integer between 1 to 1718 of SEQ ID NO:39, b is an integer of 15 to 1732, where both a and b correspond to the positions of nucleotide residues shown in SEQ ID NO:39, and where b is greater than or equal to a + 14.

10 **FEATURES OF PROTEIN ENCODED BY GENE NO: 30**

It has been discovered that this gene is expressed primarily in the following tissues: HUMAN STOMACH, Human Cerebellum.

Many polynucleotide sequences, such as EST sequences, are publicly available and accessible through sequence databases. Some of these sequences are
15 related to SEQ ID NO:40 and may have been publicly available prior to conception of the present invention. Preferably, such related polynucleotides are specifically excluded from the scope of the present invention. To list every related sequence would be cumbersome. Accordingly, preferably excluded from the present invention are one or more polynucleotides comprising a nucleotide sequence described by the
20 general formula of a-b, where a is any integer between 1 to 2193 of SEQ ID NO:40, b is an integer of 15 to 2207, where both a and b correspond to the positions of nucleotide residues shown in SEQ ID NO:40, and where b is greater than or equal to a + 14.

25 **FEATURES OF PROTEIN ENCODED BY GENE NO: 31**

It has been discovered that this gene is expressed primarily in the following tissues: Soares fetal liver spleen 1NFLS and to a lesser extent in Soares placenta Nb2HP, Soares_fetal_heart_NbHH19W, Soares_fetal_liver_spleen_1NFLS_S1, Soares_pregnant_uterus_NbHPU, Soares_NFL_T_GBC_S1, Stratagene placenta
30 (#937225), Soares_fetal_heart_NbHH19W, Soares_pregnant_uterus_NbHPU, Soares_NhHMPu_S1, Human Fetal Kidney; Reexcision, Human Synovial Sarcoma,

- NCI_CGAP_GCB1, Soares_senescent_fibroblasts_NbHSF, Soares_testis_NHT,
 Soares_fetal_heart_NbHH19W, Soares_total_fetus_Nb2HF8_9w, Soares ovary tumor
 NbHOT, Soares retina N2b4HR, Soares adult brain N2b4HB55Y,
 Soares_total_fetus_Nb2HF8_9w, Soares_fetal_liver_spleen_1NFLS_S1,
- 5 Hemangiopericytoma, Human Adrenal Gland Tumor, Soares breast 3NbHBst, Human
 fetal heart, Lambda ZAP Express, Soares_parathyroid_tumor_NbHPA,
 NCI_CGAP_Kid3, Activated T-cell(12h)/Thiouridine-re-excision, Stratagene colon
 (#937204), Primary Dendritic Cells, lib 1, Human Umbilical Vein Endothelial Cells,
 fract. B, Human Lung Cancer, subtracted, WI 38 cells, prostate-edited, Human Fetal
- 10 Spleen, Human Fetal Bone, Human T-cell lymphoma;re-excision, NTERA2
 teratocarcinoma cell line+retinoic acid (14 days), Human Epididymus, Smooth
 muscle, IL1b induced, Breast Cancer cell line, MDA 36, NTERA2 + retinoic acid, 14
 days, Soares_multiple_sclerosis_2NbHMSP, Human Fetal Epithelium (Skin),
 Soares_fetal_lung_NbHL19W, H. Kidney Cortex, subtracted, Prostate BPH, Human
- 15 Fetal Dura Mater, Human Uterine Cancer, T-Cell PHA 24 hrs, Liver HepG2 cell line.,
 Human Activated T-Cells, re-excision, Ulcerative Colitis, Human Thymus,
 Macrophage (GM-CSF treated), Pancreas Islet Cell Tumor, PC3 Prostate cell line,
 Fetal Heart, Gessler Wilms tumor, NCI_CGAP_Lu5, Soares_NbHFB,
 Soares_NSF_F8_9W_OT_PA_P_S1, Soares_fetal_liver_spleen_1NFLS_S1, Primary
- 20 Dendritic cells,frac 2, Human Fetal Lung III, Human Placenta, Human Fetal Heart,
 NCI_CGAP_Ew1, NCI_CGAP_Lu5, NCI_CGAP_Pr22, NCI_CGAP_Sch1,
 Keratinocyte, Human 8 Week Whole Embryo, Nine Week Old Early Stage Human,
 Human Cerebellum, Soares_senescent_fibroblasts_NbHSF.

Preferred epitopes include those comprising a sequence shown in SEQ ID NO.

- 25 88 as residues: Pro-31 to Pro-37.

Many polynucleotide sequences, such as EST sequences, are publicly
 available and accessible through sequence databases. Some of these sequences are
 related to SEQ ID NO:41 and may have been publicly available prior to conception of
 the present invention. Preferably, such related polynucleotides are specifically

30 excluded from the scope of the present invention. To list every related sequence
 would be cumbersome. Accordingly, preferably excluded from the present invention

are one or more polynucleotides comprising a nucleotide sequence described by the general formula of a-b, where a is any integer between 1 to 1422 of SEQ ID NO:41, b is an integer of 15 to 1436, where both a and b correspond to the positions of nucleotide residues shown in SEQ ID NO:41, and where b is greater than or equal to a + 14.

FEATURES OF PROTEIN ENCODED BY GENE NO: 32

It has been discovered that this gene is expressed primarily in Human Cerebellum.

Preferred epitopes include those comprising a sequence shown in SEQ ID NO. 89 as residues: Lys-36 to Glu-44.

Many polynucleotide sequences, such as EST sequences, are publicly available and accessible through sequence databases. Some of these sequences are related to SEQ ID NO:42 and may have been publicly available prior to conception of the present invention. Preferably, such related polynucleotides are specifically excluded from the scope of the present invention. To list every related sequence would be cumbersome. Accordingly, preferably excluded from the present invention are one or more polynucleotides comprising a nucleotide sequence described by the general formula of a-b, where a is any integer between 1 to 1405 of SEQ ID NO:42, b is an integer of 15 to 1419, where both a and b correspond to the positions of nucleotide residues shown in SEQ ID NO:42, and where b is greater than or equal to a + 14.

FEATURES OF PROTEIN ENCODED BY GENE NO: 33

It has been discovered that this gene is expressed primarily in the following tissues: Soares infant brain 1NIB, Soares_NhHMPu_S1 and to a lesser extent in Soares_fetal_heart_NbHH19W, Soares_multiple_sclerosis_2NbHMSP, Glioblastoma, Human Cerebellum.

Many polynucleotide sequences, such as EST sequences, are publicly available and accessible through sequence databases. Some of these sequences are related to SEQ ID NO:43 and may have been publicly available prior to conception of

the present invention. Preferably, such related polynucleotides are specifically excluded from the scope of the present invention. To list every related sequence would be cumbersome. Accordingly, preferably excluded from the present invention are one or more polynucleotides comprising a nucleotide sequence described by the
 5 general formula of a-b, where a is any integer between 1 to 2382 of SEQ ID NO:43, b is an integer of 15 to 2396, where both a and b correspond to the positions of nucleotide residues shown in SEQ ID NO:43, and where b is greater than or equal to a + 14.

10 **FEATURES OF PROTEIN ENCODED BY GENE NO: 34**

It has been discovered that this gene is expressed primarily in the following tissues: Soares infant brain 1NIB and to a lesser extent in Soares melanocyte 2NbHM, Soares adult brain N2b5HB55Y, Human Osteoclastoma Stromal Cells - unamplified, Colon Carcinoma, Synovial Fibroblasts (II1/TNF), subt, Stratagene NT2 neuronal
 15 precursor 937230, Soares fetal liver spleen 1NFLS, Human Tonsils, Lib 2, Human Osteoclastoma, re-excision, Fetal Heart, Early Stage Human Brain, Human Placenta, Activated T-Cell (12hs)/Thiouridine labelledEco, Osteoblasts, Soares placenta Nb2HP, Human Adult Heart, subtracted, Stratagene neuroepithelium (#937231), Stratagene colon (#937204), Stratagene hNT neuron (#937233). Human Manic
 20 Depression Tissue, Spinal Cord, re-excision, Human Bone Marrow, re-excision, Human Pancreas Tumor, Human Hippocampus, Clontech human aorta polyA+ mRNA (#6572), Human placenta cDNA (TFujiwara), Stratagene hNT neuron (#937233), NCI_CGAP_Kid3, Soares_multiple_sclerosis_2NbHMSP, Primary Dendritic cells,frac 2, Human Osteoclastoma, Nine Week Old Early Stage Human,
 25 Soares ovary tumor NbHOT, Human Gastrocnemius, Human 8 Week Whole Embryo, subtracted, HUMAN TONSILS, FRACTION 2, Human Cerebellum, subtracted, Human Adult Pulmonary, eosinophil-IL5 induced, Human Liver, H. Epididymus, cauda, Salivary Gland, Human Adult Heart;re-excision, Human adult small intestine,re-excision, Messangial cell, frac 2, Human Hypothalamus,schizophrenia,
 30 re-excision, Human Prostate Cancer, Stage C fraction, Soares_parathyroid_tumor_NbHPA, Synovial IL-1/TNF stimulated, Stratagene

- placenta (#937225), Glioblastoma, Human Whole Brain #2 - Oligo dT > 1.5Kb, Human Stomach;re-excision, Human endometrial stromal cells-treated with progesterone, Salivary Gland, Lib 2, Human Colon; re-excision, Myeloid Progenitor Cell Line, Brain Frontal Cortex, re-excision, Human Chronic Synovitis, Human
- 5 Prostate, Human Brain, Striatum, L428, Apoptotic T-cell, T-Cell PHA 24 hrs, Stromal cell TF274, Human Ovarian Cancer Reexcision, Human Hypothalamus, Schizophrenia, Stratagene colon (#937204), Human Adipose, Spinal cord, Synovial Fibroblasts (control), Stratagene neuroepithelium NT2RAMI 937234, Ulcerative Colitis, Human Thymus, Hepatocellular Tumor; re-excision, Pancreas Islet
- 10 Cell Tumor, Human fetal brain (TFujiwara), NCI_CGAP_Lu5, NCI_CGAP_Kid3, PC3 Prostate cell line, Human Substantia Nigra, Colon Normal II, NCI_CGAP_Co8, Colon Tumor II, Human Testes, Reexcision, Bone marrow, Human Adult Pulmonary;re-excision, Endothelial cells-control, Colon Normal III, NCI_CGAP_Co3, NCI_CGAP_Ew1, NCI_CGAP_Co10, NCI_CGAP_Co12,
- 15 NCI_CGAP_GCB1, NCI_CGAP_Lar1, NCI_CGAP_Br1.1, Human Amygdala, Human Microvascular Endothelial Cells, fract. A, HUMAN B CELL LYMPHOMA, NCI_CGAP_Co3, NCI_CGAP_GCB1, NCI_CGAP_Pr12, Activated T-cell(12h)/Thiouridine-re-excision, Human Testes, Human Endometrial Tumor, Stratagene pancreas (#937208), Hodgkin's Lymphoma II, NCI_CGAP_GCB1,
- 20 NCI_CGAP_Kid5, Keratinocyte, Human 8 Week Whole Embryo, Human Cerebellum, Primary Dendritic Cells, lib 1.

Many polynucleotide sequences, such as EST sequences, are publicly available and accessible through sequence databases. Some of these sequences are related to SEQ ID NO:44 and may have been publicly available prior to conception of

25 the present invention. Preferably, such related polynucleotides are specifically excluded from the scope of the present invention. To list every related sequence would be cumbersome. Accordingly, preferably excluded from the present invention are one or more polynucleotides comprising a nucleotide sequence described by the general formula of a-b, where a is any integer between 1 to 1970 of SEQ ID NO:44, b

30 is an integer of 15 to 1984, where both a and b correspond to the positions of

nucleotide residues shown in SEQ ID NO:44, and where b is greater than or equal to a + 14.

FEATURES OF PROTEIN ENCODED BY GENE NO: 35

5 It has been discovered that this gene is expressed primarily in the following tissues: Soares infant brain 1NIB, Soares placenta Nb2HP and to a lesser extent in T-Cell PHA 16 hrs, Human Brain, striatum, re-excision, Soares ovary tumor NbHOT, Healing Abdomen wound;70&90 min post incision, Human Whole Brain, re-excision, Soares adult brain N2b4HB55Y, Human Infant Brain, Human Prostate, Human
10 Hypothalamus, Schizophrenia, Human Bone Marrow, treated, Human Cerebellum.

Many polynucleotide sequences, such as EST sequences, are publicly available and accessible through sequence databases. Some of these sequences are related to SEQ ID NO:45 and may have been publicly available prior to conception of the present invention. Preferably, such related polynucleotides are specifically
15 excluded from the scope of the present invention. To list every related sequence would be cumbersome. Accordingly, preferably excluded from the present invention are one or more polynucleotides comprising a nucleotide sequence described by the general formula of a-b, where a is any integer between 1 to 1939 of SEQ ID NO:45, b is an integer of 15 to 1953, where both a and b correspond to the positions of
20 nucleotide residues shown in SEQ ID NO:45, and where b is greater than or equal to a + 14.

FEATURES OF PROTEIN ENCODED BY GENE NO: 36

It has been discovered that this gene is expressed primarily in Human
25 Cerebellum.

Preferred epitopes include those comprising a sequence shown in SEQ ID NO. 93 as residues: Gly-35 to Pro-41.

Many polynucleotide sequences, such as EST sequences, are publicly available and accessible through sequence databases. Some of these sequences are
30 related to SEQ ID NO:46 and may have been publicly available prior to conception of the present invention. Preferably, such related polynucleotides are specifically

excluded from the scope of the present invention. To list every related sequence would be cumbersome. Accordingly, preferably excluded from the present invention are one or more polynucleotides comprising a nucleotide sequence described by the general formula of a-b, where a is any integer between 1 to 1038 of SEQ ID NO:46, b is an integer of 15 to 1052, where both a and b correspond to the positions of nucleotide residues shown in SEQ ID NO:46, and where b is greater than or equal to a + 14.

FEATURES OF PROTEIN ENCODED BY GENE NO: 37

It has been discovered that this gene is expressed primarily in the following tissues: LNCAP + 30nM R1881, LNCAP + 0.3nM R1881, Human T-cell lymphoma;re-excision, Human Cerebellum.

Many polynucleotide sequences, such as EST sequences, are publicly available and accessible through sequence databases. Some of these sequences are related to SEQ ID NO:47 and may have been publicly available prior to conception of the present invention. Preferably, such related polynucleotides are specifically excluded from the scope of the present invention. To list every related sequence would be cumbersome. Accordingly, preferably excluded from the present invention are one or more polynucleotides comprising a nucleotide sequence described by the general formula of a-b, where a is any integer between 1 to 1367 of SEQ ID NO:47, b is an integer of 15 to 1381, where both a and b correspond to the positions of nucleotide residues shown in SEQ ID NO:47, and where b is greater than or equal to a + 14.

FEATURES OF PROTEIN ENCODED BY GENE NO: 38

The computer algorithm BLASTX has been used to determine that the translation product of this gene shares sequence homology with, as a non-limiting example, the sequence accessible through the following database accession no. gil1871170 (all information available through the recited accession number is incorporated herein by reference) which is described therein as "sodium channel 2

[Homo sapiens]". A partial alignment demonstrating the observed homology is shown immediately below.

```

5      >gi|1871170 sodium channel 2 [Homo sapiens]
          Length = 528

          Plus Strand HSPs:

10      Score = 223 (78.5 bits), Expect = 1.5e-26, Sum P(2) = 1.5e-26
          Identities = 39/41 (95%), Positives = 40/41 (97%), Frame = +2

          Query:   311 QQLIYLPPPWTCKAVTMSDLDFDYSITXCRIDCETRY 433
                   Q+LIYLPPPWTCKAVTMSDLDFDYSIT CRIDCETRY
          Sbjct:   278 QRLIYLPPPWTCKAVTMSDLDFDYSITACRIDCETRY 318

15      Score = 112 (39.4 bits), Expect = 1.5e-26, Sum P(2) = 1.5e-26
          Identities = 19/19 (100%), Positives = 19/19 (100%), Frame = +1

          Query:   658 GDAPYCTPEQYKECADPAL 714
                   GDAPYCTPEQYKECADPAL
          Sbjct:   332 GDAPYCTPEQYKECADPAL 350

20      Score = 87 (30.6 bits), Expect = 6.3e-24, Sum P(2) = 6.3e-24
          Identities = 15/16 (93%), Positives = 15/16 (93%), Frame = +1

25      Query:   433 LVENCNCRMVHMPGQA 480
                   LVENCNCRMVHMPG A
          Sbjct:   319 LVENCNCRMVHMPGDA 334

```

30 The segment of gi|1871170 that is shown as "Sbjct" above is set out as sequence(s) SEQ ID NO. 135, SEQ ID NO. 137, and SEQ ID NO. 139. Based on the structural similarity these homologous polypeptides are expected to share at least some biological activities. Such activities are known in the art and described elsewhere herein. Assays for determining such activities are also known in the art,

35 some of which have been described elsewhere herein. Preferred polypeptides of the invention comprise polypeptides having the amino acid sequence(s) set out as SEQ ID NO. 136, SEQ ID NO. 138, and SEQ ID NO. 140 which correspond to the Query sequence in the alignment shown above (gaps introduced in the sequence by the computer are, of course, removed).

40 It has been discovered that this gene is expressed primarily in the following tissues: Nine Week Old Early Stage Human, Human Cerebellum.

Preferred epitopes include those comprising a sequence shown in SEQ ID NO. 95 as residues: Met-14 to Asn-19.

The tissue distribution suggests the protein product of this clone is useful for the detection, treatment, and/or prevention of neurodegenerative disease states, behavioral disorders, or inflammatory conditions which include, but are not limited to Alzheimer's Disease, Parkinson's Disease, Huntington's Disease, Tourette Syndrome, 5 meningitis, encephalitis, demyelinating diseases, peripheral neuropathies, neoplasia, trauma, congenital malformations, spinal cord injuries, ischemia and infarction, aneurysms, hemorrhages, schizophrenia, mania, dementia, paranoia, obsessive compulsive disorder, depression, panic disorder, learning disabilities, ALS, psychoses, autism, and altered behaviors, including disorders in feeding, sleep 10 patterns, balance, and perception. In addition, elevated expression of this gene product in regions of the brain suggests it plays a role in normal neural function. Potentially, this gene product is involved in synapse formation, neurotransmission, learning, cognition, homeostasis, or neuronal differentiation or survival.

Moreover, the expression within embryonic tissue suggests this protein may 15 play a role in the regulation of cellular division, and may show utility in the diagnosis and treatment of cancer and other proliferative disorders. Similarly, developmental tissues rely on decisions involving cell differentiation and/or apoptosis in pattern formation. Dysregulation of apoptosis can result in inappropriate suppression of cell death, as occurs in the development of some cancers, or in failure to control the extent 20 of cell death, as is believed to occur in acquired immunodeficiency and certain neurodegenerative disorders, such as spinal muscular atrophy (SMA). Therefore, the polynucleotides and polypeptides of the present invention are useful in treating, detecting, and/or preventing said disorders and conditions, in addition to other types of degenerative conditions. Thus this protein may modulate apoptosis or tissue 25 differentiation and would be useful in the detection, treatment, and/or prevention of degenerative or proliferative conditions and diseases. Protein, as well as, antibodies directed against the protein may show utility as a tumor marker and/or immunotherapy targets for the above listed tissues.

Many polynucleotide sequences, such as EST sequences, are publicly 30 available and accessible through sequence databases. Some of these sequences are related to SEQ ID NO:48 and may have been publicly available prior to conception of

the present invention. Preferably, such related polynucleotides are specifically excluded from the scope of the present invention. To list every related sequence would be cumbersome. Accordingly, preferably excluded from the present invention are one or more polynucleotides comprising a nucleotide sequence described by the
 5 general formula of a-b, where a is any integer between 1 to 978 of SEQ ID NO:48, b is an integer of 15 to 992, where both a and b correspond to the positions of nucleotide residues shown in SEQ ID NO:48, and where b is greater than or equal to a + 14.

10 **FEATURES OF PROTEIN ENCODED BY GENE NO: 39**

It has been discovered that this gene is expressed primarily in the following tissues: Soares melanocyte 2NbHM and to a lesser extent in Soares_multiple_sclerosis_2NbHMSP, Soares adult brain N2b4HB55Y, Human 8 Week Whole Embryo, Soares fetal liver spleen 1NFLS, Stratagene hNT neuron
 15 (#937233), H. Frontal cortex,epileptic;re-excision, Human Amygdala, Soares ovary tumor NbHOT, Stratagene ovarian cancer (#937219), Human Hypothalamus,Schizophrenia, Soares_fetal_liver_spleen_1NFLS_S1, Stratagene colon HT29 (#937221), Human Endometrial Tumor, Nine Week Old Early Stage Human, Human Cerebellum, Human Thymus Tumor, Early Stage Human Brain, random
 20 primed, Human Amygdala Depression, re-excision, HPAS (human pancreas, subtracted), H. Frontal Cortex, Epileptic, Human Hippocampus, subtracted, HUMAN TONSILS, FRACTION 2, Human Adult Retina, Human Pancreatic Carcinoma, Frontal lobe,dementia;re-excision, Human adult small intestine,re-excision, Human Lung Cancer;re-excision, Human Epididymus, Human Umbilical Vein, Endo.
 25 remake, Synovial hypoxia-RSF subtracted, Stratagene pancreas (#937208), Human Adipose, Human Whole Six Week Old Embryo, Human Substantia Nigra, Brain frontal cortex, Soares breast 3NbHBst, Soares_fetal_heart_NbHH19W, Bone marrow, Human Fetal Heart, Endothelial cells-control, Bone Marrow Cell Line (RS4;11).

Many polynucleotide sequences, such as EST sequences, are publicly
 30 available and accessible through sequence databases. Some of these sequences are related to SEQ ID NO:49 and may have been publicly available prior to conception of

the present invention. Preferably, such related polynucleotides are specifically excluded from the scope of the present invention. To list every related sequence would be cumbersome. Accordingly, preferably excluded from the present invention are one or more polynucleotides comprising a nucleotide sequence described by the
5 general formula of a-b, where a is any integer between 1 to 2115 of SEQ ID NO:49, b is an integer of 15 to 2129, where both a and b correspond to the positions of nucleotide residues shown in SEQ ID NO:49, and where b is greater than or equal to a + 14.

10 **FEATURES OF PROTEIN ENCODED BY GENE NO: 40**

It has been discovered that this gene is expressed primarily in the following tissues: Human Prostate Cancer, Stage C fraction, Human Cerebellum.

Many polynucleotide sequences, such as EST sequences, are publicly available and accessible through sequence databases. Some of these sequences are
15 related to SEQ ID NO:50 and may have been publicly available prior to conception of the present invention. Preferably, such related polynucleotides are specifically excluded from the scope of the present invention. To list every related sequence would be cumbersome. Accordingly, preferably excluded from the present invention are one or more polynucleotides comprising a nucleotide sequence described by the
20 general formula of a-b, where a is any integer between 1 to 1460 of SEQ ID NO:50, b is an integer of 15 to 1474, where both a and b correspond to the positions of nucleotide residues shown in SEQ ID NO:50, and where b is greater than or equal to a + 14.

25 **FEATURES OF PROTEIN ENCODED BY GENE NO: 41**

It has been discovered that this gene is expressed primarily in the following tissues: Human Pineal Gland, Human Cerebellum.

Many polynucleotide sequences, such as EST sequences, are publicly available and accessible through sequence databases. Some of these sequences are
30 related to SEQ ID NO:51 and may have been publicly available prior to conception of the present invention. Preferably, such related polynucleotides are specifically

excluded from the scope of the present invention. To list every related sequence would be cumbersome. Accordingly, preferably excluded from the present invention are one or more polynucleotides comprising a nucleotide sequence described by the general formula of a-b, where a is any integer between 1 to 2086 of SEQ ID NO:51, b
5 is an integer of 15 to 2100, where both a and b correspond to the positions of nucleotide residues shown in SEQ ID NO:51, and where b is greater than or equal to a + 14.

FEATURES OF PROTEIN ENCODED BY GENE NO: 42

10 It has been discovered that this gene is expressed primarily in Human Cerebellum..

Many polynucleotide sequences, such as EST sequences, are publicly available and accessible through sequence databases. Some of these sequences are related to SEQ ID NO:52 and may have been publicly available prior to conception of
15 the present invention. Preferably, such related polynucleotides are specifically excluded from the scope of the present invention. To list every related sequence would be cumbersome. Accordingly, preferably excluded from the present invention are one or more polynucleotides comprising a nucleotide sequence described by the general formula of a-b, where a is any integer between 1 to 1593 of SEQ ID NO:52, b
20 is an integer of 15 to 1607, where both a and b correspond to the positions of nucleotide residues shown in SEQ ID NO:52, and where b is greater than or equal to a + 14.

FEATURES OF PROTEIN ENCODED BY GENE NO: 43

25 It has been discovered that this gene is expressed primarily in the following tissues: Human Osteoclastoma and to a lesser extent in Soares retina N2b4HR, normalized infant brain cDNA, NCI_CGAP_GCB1, Human Osteoclastoma, re-excision, Soares placenta Nb2HP, Soares infant brain 1NIB, NCI_CGAP_GCB1, Soares retina N2b5HR, Human Activated T-Cells, re-excision, Stratagene hNT
30 neuron (#937233), NTERA2, control, Soares_total_fetus_Nb2HF8_9w, Nine Week Old Early Stage Human, Human Fetal Brain, normalized 50021F, H. Striatum

Depression, subt II, Brain; normal, Larynx carcinoma IV, Osteoclastoma-normalized A, Human Adult Liver, subtracted, Soares_multiple_sclerosis_2NbHMSP, Human T-cell lymphoma; re-excision, Jurkat T-cell G1 phase, Human Adult Small Intestine, Human Prostate, Soares_multiple_sclerosis_2NbHMSP, H. Kidney Medulla, re-excision, Stromal cell TF274, Human Testes Tumor, re-excision, Rejected Kidney, lib 4, Hepatocellular Tumor; re-excision, Soares_pregnant_uterus_NbHPU, Human Liver, normal, Human Gall Bladder, Colon Carcinoma, Human Substantia Nigra, Human Placenta, NCI_CGAP_Lu5, NCI_CGAP_Kid5, Soares_pregnant_uterus_NbHPU, Soares_parathyroid_tumor_NbHPA, HUMAN B
 10 CELL LYMPHOMA, T cell helper II, Human 8 Week Whole Embryo, Human Cerebellum.

Preferred epitopes include those comprising a sequence shown in SEQ ID NO. 100 as residues: Thr-33 to Trp-38.

Many polynucleotide sequences, such as EST sequences, are publicly
 15 available and accessible through sequence databases. Some of these sequences are related to SEQ ID NO:53 and may have been publicly available prior to conception of the present invention. Preferably, such related polynucleotides are specifically excluded from the scope of the present invention. To list every related sequence would be cumbersome. Accordingly, preferably excluded from the present invention
 20 are one or more polynucleotides comprising a nucleotide sequence described by the general formula of a-b, where a is any integer between 1 to 1418 of SEQ ID NO:53, b is an integer of 15 to 1432, where both a and b correspond to the positions of nucleotide residues shown in SEQ ID NO:53, and where b is greater than or equal to a + 14.

25

FEATURES OF PROTEIN ENCODED BY GENE NO: 44

The computer algorithm BLASTX has been used to determine that the translation product of this gene shares sequence homology with, as a non-limiting example, the sequence accessible through the following database accession no.
 30 dbj|AB006085_1 (all information available through the recited accession number is incorporated herein by reference) which is described therein as "MINDIN2 [Danio

rerio] ". A partial alignment demonstrating the observed homology is shown immediately below.

```

5      >dbj||AB006085_1 (AB006085) MINDIN2 [Danio rerio]
          Length = 331

          Plus Strand HSPs:

10      Score = 261 (91.9 bits), Expect = 8.5e-21, P = 8.5e-21
          Identities = 42/63 (66%), Positives = 49/63 (77%), Frame = +3

          Query:  1497 GPVPETPLDCEVSLWSSWGLCGGHCGRGKSRTRYVVRVQPANNGSPCPELEEEAECVPD
          1676
          GP+  TPLDCEVS+WS WGLC G CG  G K RTRY+ + PANNG+PCP LEE+  C+PD
15      Sbjct:  269 GPLINTPLDCEVSVWSPWGLCKGQCGEKGVKHRTRYIHMHPANNGAPCPSLEEKRLCIPD
          328

          Query:  1677 NCV 1685
          NCV
20      Sbjct:  329 NCV 331

```

The segment of dbj||AB006085_1 that is shown as "Sbjct" above is set out as sequence(s) SEQ ID NO. 141. Based on the structural similarity these homologous polypeptides are expected to share at least some biological activities. Such activities are known in the art and described elsewhere herein. Assays for determining such activities are also known in the art some of which have been described elsewhere herein. Preferred polypeptides of the invention comprise polypeptides having the amino acid sequence(s) set out as SEQ ID NO. 142 which correspond to the Query sequence in the alignment shown above (gaps introduced in the sequence by the computer are, of course, removed).

Additionally, the translation product of this gene shares sequence homology with mindin, F-spondin, and neuronal attachment factor-1 (see, e.g., Genseq accession number W23663 and WO 9729189; which is hereby incorporated by reference herein.) which is thought to be important in embryonic development, neural cell pattern formation, axonal growth. F-spondin (FSP) is a gene that is predominantly expressed during the early development of the vertebrate nervous system. The main function is thought to be in neural cell pattern formation and axonal growth. It was found in a subtractive hybridization screen designed to isolate floor-plate specific genes. The floor-plate provides diffusible signals that act on the neurons that extend from the developing spinal cord. These signals can lead to chemoattraction and

fasciculation of commissural axons in the ventral midline. F-spondin mRNA is expressed at high levels in the developing neural tube at the ventral midline even before cell differentiation markers can detect the floor-plate. F-spondin is not detectable in other regions of the spinal cord until later in embryonic life. There is also transient F-spondin expression early in peripheral nerve development which diminishes to undetectable levels following birth. The adult central nervous system contains F-spondin while the peripheral nerve (sciatic nerve) does not. Outside the adult nervous system, organs such as the lung and kidney also express F-spondin. Based on the sequence similarity, the translation product of this clone is expected to share at least some biological activities with F-spondin and/or neuronal attachment factor proteins. Such activities are known in the art, some of which are described elsewhere herein.

It has been discovered that this gene is expressed primarily in the following tissues: Human endometrial stromal cells-treated with progesterone and to a lesser extent in Human endometrial stromal cells, Human endometrial stromal cells-treated with estradiol, NCI_CGAP_Pr2, Soares_pregnant_uterus_NbHPU, Soares_pregnant_uterus_NbHPU, Human Adrenal Gland Tumor, NCI_CGAP_HSC1, Colon Carcinoma, Soares_fetal_heart_NbHH19W, Soares ovary tumor NbHOT, Larynx Normal, Colorectal Tumor, Human epithelioid sarcoma, SKIN, H. Atrophic Endometrium, HSA 172 Cells, Hepatocellular Tumor, Human Prostate Cancer, Stage C fraction, Human Pituitary, subt IX, wilm's tumor, Human Prostate, human ovarian cancer, Human Pancreas Tumor, Synovial Fibroblasts (control), NCI_CGAP_Br2, NCI_CGAP_Kid6, NCI_CGAP_Pr12, NCI_CGAP_Pr21, NCI_CGAP_Pr23, NCI_CGAP_Sch1, Human Synovial Sarcoma, CD34 depleted Buffy Coat (Cord Blood), re-excision, Human Cerebellum, Soares fetal liver spleen 1NFLS. Polynucleotides and polypeptides of the invention are useful as reagents for differential identification of the tissue(s) or cell type(s) present in a biological sample and for diagnosis of diseases and conditions which include but are not limited to: cancer, disorders and/or diseases of the central nervous system (CNS), developmental and reproductive disorders, microvascular disease, vascular leak syndrome, aneurysm, stroke, embolism, thrombosis, coronary artery disease and/or

arteriosclerosis. Similarly, polypeptides and antibodies directed to these polypeptides are useful in providing immunological probes for differential identification of the tissue(s) or cell type(s). For a number of disorders of the above tissues or cells, particularly of the CNS, reproductive system and the vasculature expression of this
5 gene at significantly higher or lower levels may be routinely detected in certain tissues or cell types (e.g., cancerous and wounded tissues) or bodily fluids (e.g., serum, plasma, urine, synovial fluid and spinal fluid) or another tissue or sample taken from an individual having such a disorder, relative to the standard gene expression level, i.e., the expression level in healthy tissue or bodily fluid from an individual not
10 having the disorder.

Preferred epitopes include those comprising a sequence shown in SEQ ID NO. 101 as residues: Pro-6 to Gln-11, Pro-32 to Arg-37, His-41 to Trp-46, Pro-49 to Gly-54.

The tissue distribution in endometrial cells and tissues suggests that the
15 protein is useful in the detection, treatment, and/or prevention of developmental and reproductive disorders. Moreover, the protein is useful in the detection, treatment, and/or prevention of a variety of vascular disorders and conditions, which include, but are not limited to microvascular disease, vascular leak syndrome, aneurysm, stroke, embolism, thrombosis, coronary artery disease, arteriosclerosis, and/or
20 atherosclerosis. Protein, as well as, antibodies directed against the protein may show utility as a tumor marker and/or immunotherapy targets for the above listed tissues.

Moreover, the expression within embryonic tissue, endometrium and other cellular sources marked by proliferating cells indicates this protein may play a role in the regulation of cellular division, and may show utility in the diagnosis, treatment,
25 and/or prevention of developmental diseases and disorders, including cancer, and other proliferative conditions. Representative uses are described in the "Hyperproliferative Disorders" and "Regeneration" sections below and elsewhere herein. Briefly, developmental tissues rely on decisions involving cell differentiation and/or apoptosis in pattern formation. Dysregulation of apoptosis can result in
30 inappropriate suppression of cell death, as occurs in the development of some cancers, or in failure to control the extent of cell death, as is believed to occur in acquired

immunodeficiency and certain degenerative disorders, such as spinal muscular atrophy (SMA). Alternatively, this gene product may be involved in the pattern of cellular proliferation that accompanies early embryogenesis. Thus, aberrant expression of this gene product in tissues - particularly adult tissues - may correlate with patterns of abnormal cellular proliferation, such as found in various cancers. Because of potential roles in proliferation and differentiation, this gene product may have applications in the adult for tissue regeneration and the treatment of cancers. It may also act as a morphogen to control cell and tissue type specification. Therefore, the polynucleotides and polypeptides of the present invention are useful in treating, detecting, and/or preventing said disorders and conditions, in addition to other types of degenerative conditions. Thus this protein may modulate apoptosis or tissue differentiation and would be useful in the detection, treatment, and/or prevention of degenerative or proliferative conditions and diseases. The protein would be useful in modulating the immune response to aberrant polypeptides, as may exist in proliferating and cancerous cells and tissues. The protein can also be used to gain new insight into the regulation of cellular growth and proliferation. Furthermore, the protein may also be used to determine biological activity, to raise antibodies, as tissue markers, to isolate cognate ligands or receptors, to identify agents that modulate their interactions, in addition to its use as a nutritional supplement. Protein, as well as, antibodies directed against the protein may show utility as a tumor marker and/or immunotherapy targets for the above listed tissues.

Many polynucleotide sequences, such as EST sequences, are publicly available and accessible through sequence databases. Some of these sequences are related to SEQ ID NO:54 and may have been publicly available prior to conception of the present invention. Preferably, such related polynucleotides are specifically excluded from the scope of the present invention. To list every related sequence would be cumbersome. Accordingly, preferably excluded from the present invention are one or more polynucleotides comprising a nucleotide sequence described by the general formula of a-b, where a is any integer between 1 to 2247 of SEQ ID NO:54, b is an integer of 15 to 2261, where both a and b correspond to the positions of

nucleotide residues shown in SEQ ID NO:54, and where b is greater than or equal to a + 14.

FEATURES OF PROTEIN ENCODED BY GENE NO: 45

5 It has been discovered that this gene is expressed primarily in Human Chondrosarcoma..

Preferred epitopes include those comprising a sequence shown in SEQ ID NO. 102 as residues: Thr-26 to Tyr-38.

Many polynucleotide sequences, such as EST sequences, are publicly
10 available and accessible through sequence databases. Some of these sequences are related to SEQ ID NO:55 and may have been publicly available prior to conception of the present invention. Preferably, such related polynucleotides are specifically excluded from the scope of the present invention. To list every related sequence would be cumbersome. Accordingly, preferably excluded from the present invention
15 are one or more polynucleotides comprising a nucleotide sequence described by the general formula of a-b, where a is any integer between 1 to 799 of SEQ ID NO:55, b is an integer of 15 to 813, where both a and b correspond to the positions of nucleotide residues shown in SEQ ID NO:55, and where b is greater than or equal to a + 14.

20

FEATURES OF PROTEIN ENCODED BY GENE NO: 46

The computer algorithm BLASTX has been used to determine that the translation product of this gene shares sequence homology with, as a non-limiting example, the sequence accessible through the following database accession no.
25 gill151254 (all information available through the recited accession number is incorporated herein by reference) which is described therein as "transmembrane receptor [Mus musculus]". A partial alignment demonstrating the observed homology is shown immediately below.

30

```
>gi|1151254 transmembrane receptor [Mus musculus] >sp|Q61088|Q61088
FRIZZLED
HOMOLOG 4 (TRANSMEMBRANE RECEPTOR).
Length = 537
```

Plus Strand HSPs:

5 Score = 942 (331.6 bits), Expect = 1.0e-166, Sum P(2) = 1.0e-166
 Identities = 166/171 (97%), Positives = 170/171 (99%), Frame = +3

Query: 354 CSVYVPMCTEKINIPIGPCGGMCLSVKRRCEPVLKEFGFAWPESLNC SKFPQNDHNHMC
 533

10 Sbjct: 99 CSVYVPMCTEKINIPIGPCGGMCLSVKRRCEPVL+EFGFAWP++LNC SKFPQNDHNHMC
 158

Query: 534 MEGPGDEEVPLPHKTPIQPGEECHSVGTNSDQYIWKRLNCVLKCGYDAGLYSRSAKEF
 713

15 Sbjct: 159 MEGPGDEEVPLPHKTPIQPGEECHSVG+NSDQYIWKRLNCVLKCGYDAGLYSRSAKEF
 218

Query: 714 TDIWMAVWASLCFISTFTVLTLFLIDSSRFSYPERPIIFLSMCYNIYSIAY 866
 TDIWMAVWASLCFIST FTVLTLFLIDSSRFSYPERPIIFLSMCYNIYSIAY
 20 Sbjct: 219 TDIWMAVWASLCFISTFTVLTLFLIDSSRFSYPERPIIFLSMCYNIYSIAY 269

Score = 708 (249.2 bits), Expect = 1.0e-166, Sum P(2) = 1.0e-166
 Identities = 140/177 (79%), Positives = 146/177 (82%), Frame = +1

25 Query: 826 YFSVCAIIFIALLVRLTVGRERISCDFEAAEPVLIQEGLKNTGCAIIFLLMYFFGMAS
 1005

30 Sbjct: 257 FLSCYNIYSIAYIVRLTVGRERISCDFEAAEPVLIQEGLKNTGCAIIFLLMYFFGMAS
 316

Query: 1006 SIWWVILTLTWFLAAGLKWGHEAIEMHSSYFHIAAWAIPAVKTIVILIMRLVDADEL TGL
 1185

35 Sbjct: 317 SIWWVILTLTWFLAAGLKWGHEAIEMHSSYFHIAAWAIPAVKTIVILIMRLVDADEL TGL
 376

Query: 1186 CYVGNQNLDA LTGFV VAPLFTYLAL EXXXXXXXXXXXXXXXXXKIFKXDGTKDQTKLEK
 1356

40 Sbjct: 377 CYVGNQNLDA LTGFV VAPLFTYL + DGTK KLE+
 432

The segment of gill151254 that is shown as "Sbjct" above is set out as
 sequence(s) SEQ ID NO. 143 and SEQ ID NO. 145. Based on the structural similarity
 45 these homologous polypeptides are expected to share at least some biological
 activities. Such activities are known in the art and described elsewhere herein. Assays
 for determining such activities are also known in the art, some of which have been
 described elsewhere herein. Preferred polypeptides of the invention comprise
 polypeptides having the amino acid sequence(s) set out as SEQ ID NO. 144 and SEQ
 50 ID NO. 146 which correspond to the Query sequence in the alignment shown above
 (gaps introduced in the sequence by the computer are, of course, removed).

It has been discovered that this gene is expressed primarily in the following tissues: Soares placenta Nb2HP and to a lesser extent in Synovial hypoxia-RSF subtracted, Synovial hypoxia, Human Chondrosarcoma, Human Testes, Reexcision, Soares_fetal_heart_NbHH19W, Endothelial cells-control, NCI_CGAP_Pr22,
5 Hodgkin's Lymphoma II.

Preferred epitopes include those comprising a sequence shown in SEQ ID NO. 103 as residues: Pro-20 to Phe-25.

In *Drosophila melanogaster*, the frizzled gene plays an essential role in the development of tissue polarity as assessed by the orientation of cuticular structures.
10 Through a combination of random cDNA sequencing, degenerate polymerase chain reaction amplification, and low stringency hybridization, six novel frizzled homologues have been identified from mammals, at least 11 from zebrafish, several from chicken and sea urchin, and one from *Caenorhabditis elegans*. The complete deduced amino acid sequences of the mammalian and nematode homologues share
15 with the *Drosophila* frizzled protein a conserved amino-terminal cysteine-rich domain and seven putative transmembrane segments. Each of the mammalian homologues is expressed in a distinctive set of tissues in the adult, and at least three are expressed during embryogenesis. As hypothesized for the *Drosophila* frizzled protein, the frizzled homologues are likely to act as transmembrane receptors for as yet
20 unidentified ligands. These observations predict the existence of a family of signal transduction pathways that are homologous to the pathway that determines tissue polarity in *Drosophila*.

The expression of the present invention within fetal tissue and other cellular sources marked by proliferating cells suggests this protein may play a role in the
25 regulation of cellular division, and may show utility in the diagnosis and treatment of cancer and other proliferative disorders. Similarly, developmental tissues rely on decisions involving cell differentiation and/or apoptosis in pattern formation. Dysregulation of apoptosis can result in inappropriate suppression of cell death, as occurs in the development of some cancers, or in failure to control the extent of cell
30 death, as is believed to occur in acquired immunodeficiency and certain neurodegenerative disorders, such as spinal muscular atrophy (SMA). Therefore, the

polynucleotides and polypeptides of the present invention are useful in treating, detecting, and/or preventing said disorders and conditions, in addition to other types of degenerative conditions. Thus this protein may modulate apoptosis or tissue differentiation and would be useful in the detection, treatment, and/or prevention of degenerative or proliferative conditions and diseases. Protein, as well as, antibodies directed against the protein may show utility as a tumor marker and/or immunotherapy targets for the above listed tissues.

Many polynucleotide sequences, such as EST sequences, are publicly available and accessible through sequence databases. Some of these sequences are related to SEQ ID NO:56 and may have been publicly available prior to conception of the present invention. Preferably, such related polynucleotides are specifically excluded from the scope of the present invention. To list every related sequence would be cumbersome. Accordingly, preferably excluded from the present invention are one or more polynucleotides comprising a nucleotide sequence described by the general formula of a-b, where a is any integer between 1 to 1378 of SEQ ID NO:56, b is an integer of 15 to 1392, where both a and b correspond to the positions of nucleotide residues shown in SEQ ID NO:56, and where b is greater than or equal to a + 14.

20 FEATURES OF PROTEIN ENCODED BY GENE NO: 47

The computer algorithm BLASTX has been used to determine that the translation product of this gene shares sequence homology with, as a non-limiting example, the sequence accessible through the following database accession no. gi|2738927 (all information available through the recited accession number is incorporated herein by reference) which is described therein as "(AF020762) unknown protein [Homo sapiens]". A partial alignment demonstrating the observed homology is shown immediately below.

30 >gi|2738927 (AF020762) unknown protein [Homo sapiens] >sp|O43466|O43466
HYPOTHETICAL 31.3 KD PROTEIN (FRAGMENT).
Length = 267

Plus Strand HSPs:

Score = 434 (152.8 bits), Expect = 3.8e-40, P = 3.8e-40
 Identities = 115/255 (45%), Positives = 128/255 (50%), Frame = +1

```

5   Query:   217 VL*EPVVTVSSNLKSIS*QVYIRCSWL*LKAIFKT*KL-NTSQIIIEKEMASV*SRSE*P
393
      +L EPVVTVSSNLKSIS QVYIR SWL LKAIFKT K  + +  ++  S+ S
      Sbjct:   3 LLSEPVVTVSSNLKSISYQVYIRYSWLSLKAIFKTSKTQHLTDYNRKRNGLSLISFRMTQ
62
10  Query:   394 RLFLWVQLFSSSELYRELFSXXXXXXXXXXXXXXXXXSRVVEHLRDFSSGTSTNFESLYL
573
      +F                               F                               SRYVEHLRDFSSGTSTNF  L
      Sbjct:   63 IVFCFGCSCLVQSYITENYFLRNLKTRMFKTNSIIEVSRYVEHLRDFSSGTSTNFRILVS
15  122
      Query:   574 YLVFILLVAKYRFF--VLFCFGFIEYLKCLKLFCCTKNKIKDFIXXXXXXXXXXXXXXXXXXII
747
      K + F  VLFCFGFIEYLKCLKLFC KNKIKDFI
      Sbjct:   123 LFGIHTTSRKIQVFCFVLFCFGFIEYLKCLKLFCPKNKIKDFIKLKLAPTELKYIKCIKYN
20  182
      Query:   748 QHMGSNIALIWKSLPETVYFLLCYLVLTNPTLFGFGITY*GELEIHMPMLKFYSFNLCY
927
      GSNIALIWKSLPETVYFLLCYLVLT P          +          +  P L F +
      Sbjct:   183 SAYGSNIALIWKSLPETVYFLLCYLVLTTPHYSSLELLIKENWKYICPLNFIALICVI
25  242
      Query:   928 FFIDGKRYIFFSLLK 972
      + G+  FF  K
      Sbjct:   243 SLLTGRGTSFFPYSK 257
30

```

The segment of gil2738927 that is shown as "Sbjct" above is set out as sequence(s) SEQ ID NO. 147. Based on the structural similarity these homologous polypeptides are expected to share at least some biological activities. Such activities are known in the art and described elsewhere herein. Assays for determining such activities are also known in the art, some of which have been described elsewhere herein. Preferred polypeptides of the invention comprise polypeptides having the amino acid sequence(s) set out as SEQ ID NO. 148 which correspond to the Query sequence in the alignment shown above (gaps introduced in the sequence by the computer are, of course, removed).

It has been discovered that this gene is expressed primarily in the following tissues: Soares_multiple_sclerosis_2NbHMSP and to a lesser extent in Soares_parathyroid_tumor_NbHPA, Soares melanocyte 2NbHM, Human Cerebellum, Human Colon Cancer;re-excision, NCI_CGAP_GCB1, Human adult small intestine, re-excision, NTERA2 + retinoic acid, 14 days, Glioblastoma, Human Stomach;re-excision, Stratagene neuroepithelium NT2RAMI 937234, Human

endometrial stromal cells, Fetal Liver, subtraction II, L428, 12 Week Old Early Stage Human, II, Human fetal heart, Lambda ZAP Express, NCI_CGAP_Br2, NCI_CGAP_Ov8, NCI_CGAP_GCB1, NCI_CGAP_Kid6, Human Umbilical Vein Endothelial Cells, uninduced, T-Cell PHA 24 hrs, Stromal cell TF274, Ulcerative
5 Colitis, Bone Marrow Stromal Cell, untreated, Human Adrenal Gland Tumor, NTERA2, control, PC3 Prostate cell line, Adipocytes, Human Testes Tumor, Colon Tumor II, Human Adult Pulmonary;re-excision, NCI_CGAP_GCB1, Human Osteoclastoma, Spleen, Chronic lymphocytic leukemia, Human Testes, Soares fetal liver spleen 1NFLS.

10 The tissue distribution in multiple sclerotic tissue suggests the protein product of this clone is useful for the detection, treatment, and/or prevention of various muscle disorders, such as muscular dystrophy, cardiomyopathy, fibroids, myomas, rhabdomyosarcomas, and musculo-degenerative conditions and/or diseases. Protein, as well as, antibodies directed against the protein may show utility as a tumor marker
15 and/or immunotherapy targets for the above listed tissues.

Many polynucleotide sequences, such as EST sequences, are publicly available and accessible through sequence databases. Some of these sequences are related to SEQ ID NO:57 and may have been publicly available prior to conception of the present invention. Preferably, such related polynucleotides are specifically
20 excluded from the scope of the present invention. To list every related sequence would be cumbersome. Accordingly, preferably excluded from the present invention are one or more polynucleotides comprising a nucleotide sequence described by the general formula of a-b, where a is any integer between 1 to 1793 of SEQ ID NO:57, b is an integer of 15 to 1807, where both a and b correspond to the positions of
25 nucleotide residues shown in SEQ ID NO:57, and where b is greater than or equal to a + 14.

Table 1

Gene No.	cDNA Clone ID	ATCC Deposit No:Z and Date	Vector	NT SEQ ID NO: X	Total NT Seq.	5' NT of Clone Seq.	3' NT of Clone Seq.	5' NT of Start Codon	5' NT of AA of Signal Pep	AA SEQ ID NO: Y	First AA of Sig Pep	Last AA of Sig Pep	First AA of Secreted Portion	Last AA of ORF
1	HCEEU18	203917 4/8/99	Uni-ZAP XR	11	1229	1	1229	209	209	58	1	30	31	43
2	HCDCB03	203917 4/8/99	Uni-ZAP XR	12	1538	1	1538	32	32	59	1	35	36	139
3	HCDES69	203917 4/8/99	Uni-ZAP XR	13	1032	1	1032		271	60	1			2
4	HCE1G78	203917 4/8/99	Uni-ZAP XR	14	1896	1	1896	77	77	61	1	18	19	254
5	HCE3C46	203917 4/8/99	Uni-ZAP XR	15	1230	1	1230		290	62	1			6
6	HCDEB19	203917 4/8/99	Uni-ZAP XR	16	1744	1	1744		266	63	1	32	33	48
7	HCE5B62	203917 4/8/99	Uni-ZAP XR	17	1714	1	1714	204	204	64	1	23	24	29
8	HCEDR26	203917 4/8/99	Uni-ZAP XR	18	1419	1	1419	177	177	65	1	26	27	55
9	HCEEM33	203917 4/8/99	Uni-ZAP XR	19	1165	1	1165	244	244	66	1	26	27	35
10	HCEFG93	203917 4/8/99	Uni-ZAP XR	20	2280	1	2280		166	67	1			14

Gene No.	cDNA Clone ID	ATCC Deposit No:Z and Date	Vector	NT SEQ ID NO: X	Total NT Seq.	5' NT of Clone Seq.	3' NT of Clone Seq.	5' NT of Start Codon	5' NT of First AA of Signal Pep	AA SEQ ID NO: Y	First AA of Sig Pep	Last AA of Sig Pep	First AA of Secreted Portion	Last AA of ORF
11	HCDBW67	203917 4/8/99	Uni-ZAP XR	21	677	409	677		498	68	1	20	21	44
12	HCDBZ31	203917 4/8/99	Uni-ZAP XR	22	858	21	858		565	69	1	11	12	14
13	HCDC51	203917 4/8/99	Uni-ZAP XR	23	1085	1	1085	46	46	70	1			2
14	HCDDDB15	203917 4/8/99	Uni-ZAP XR	24	1422	514	1202	668	668	71	1			12
15	HCDDX81	203917 4/8/99	Uni-ZAP XR	25	419	1	419	274	274	72	1	16	17	48
16	HBZAI75	203917 4/8/99	pSport1	26	1531	1	1531		260	73	1	25	26	48
17	HCDEN46	203917 4/8/99	Uni-ZAP XR	27	957	1	957	83	83	74	1	33	34	59
18	HCEID45	203917 4/8/99	Uni-ZAP XR	28	1390	1	1390	101	101	75	1	28	29	44
19	HCEIN56	203917 4/8/99	Uni-ZAP XR	29	1590	1	1587	106	106	76	1	37	38	38
20	HCEIY27	203917 4/8/99	Uni-ZAP XR	30	1651	1	1651	374	374	77	1	20	21	133

Gene No.	cDNA Clone ID	ATCC Deposit No:Z and Date	Vector	NT SEQ ID NO: X	Total NT Seq.	5' NT of Clone Seq.	3' NT of Clone Seq.	5' NT of Start Codon	5' NT of First AA of Signal Pep	AA SEQ ID NO: Y	First AA of Sig Pep	Last AA of Sig Pep	First AA of Secreted Portion	Last AA of ORF
21	HCE1Y34	203917 4/8/99	Uni-ZAP XR	31	1720	1	1720		757	78	1			8
22	HCE2B57	203917 4/8/99	Uni-ZAP XR	32	1247	1	1247		279	79	1			9
23	HCE2I23	203917 4/8/99	Uni-ZAP XR	33	977	1	977	56	56	80	1	33	34	90
24	HCE2P90	203917 4/8/99	Uni-ZAP XR	34	1893	1	1893		142	81	1	25	26	42
25	HCE3A54	203917 4/8/99	Uni-ZAP XR	35	847	1	847	51	51	82	1	18	19	42
26	HCE3D89	203917 4/8/99	Uni-ZAP XR	36	1100	17	1100	73	73	83	1	33	34	87
27	HCE3N23	203917 4/8/99	Uni-ZAP XR	37	3145	595	3114	789	789	84	1	24	25	51
28	HCE4T64	203917 4/8/99	Uni-ZAP XR	38	840	1	840		91	85	1	33	34	76
29	HCE5F78	203917 4/8/99	Uni-ZAP XR	39	1732	282	1732		566	86	1	14	15	32
30	HCE5J64	203917 4/8/99	Uni-ZAP XR	40	2207	1	2207	114	114	87	1	31	32	39

Gene No.	cDNA Clone ID	ATCC Deposit No:Z and Date	Vector	NT SEQ ID NO: X	Total NT Seq.	5' NT of Clone Seq.	3' NT of Clone Seq.	5' NT of Start Codon	5' NT of First AA of Signal Pep	AA SEQ ID NO: Y	First AA of Sig Pep	Last AA of Sig Pep	First AA of Secreted Portion	Last AA of ORF
31	HCECO77	203917 4/8/99	Uni-ZAP XR	41	1436	1	1436	124	124	88	1	28	29	66
32	HCEDH42	203917 4/8/99	Uni-ZAP XR	42	1419	1	1419	71	71	89	1	20	21	55
33	HCEDI05	203917 4/8/99	Uni-ZAP XR	43	2396	1	2396	67	67	90	1	23	24	46
34	HCEDI26	203917 4/8/99	Uni-ZAP XR	44	1984	34	1984		981	91	1			2
35	HCEDO17	203917 4/8/99	Uni-ZAP XR	45	1953	255	1953		608	92	1			34
36	HCEEE79	203917 4/8/99	Uni-ZAP XR	46	1052	1	1052	131	131	93	1	17	18	55
37	HCEEP16	203917 4/8/99	Uni-ZAP XR	47	1381	1	1381	276	276	94	1			1
38	HCEEQ25	203917 4/8/99	Uni-ZAP XR	48	992	1	992		111	95	1	15	16	23
39	HCE3J43	203917 4/8/99	Uni-ZAP XR	49	2129	836	2087	1075	1075	96	1	22	23	22
40	HCEFA10	203917 4/8/99	Uni-ZAP XR	50	1474	1	1474		282	97	1			4

Gene No.	cDNA Clone ID	ATCC Deposit No.:Z and Date	Vector	NT SEQ ID NO: X	Total NT Seq.	5' NT of Clone Seq.	3' NT of Clone Seq.	5' NT of Start Codon	5' NT of First AA of Signal Pep	AA SEQ ID NO: Y	First AA of Sig Pep	Last AA of Sig Pep	First AA of Secreted Portion	Last AA of ORF
41	HCEFA50	203917 4/8/99	Uni-ZAP XR	51	2100	1	2100	223	223	98	1			1
42	HCEFA94	203917 4/8/99	Uni-ZAP XR	52	1607	1	1607	232	232	99	1			17
43	HCEFC27	203917 4/8/99	Uni-ZAP XR	53	1432	1	1432	224	224	100	1	22	23	38
44	HCEFH31	203917 4/8/99	Uni-ZAP XR	54	2261	99	2241	260	260	101	1	33	34	59
45	HCDDL48	203917 4/8/99	Uni-ZAP XR	55	813	1	813	333	333	102	1	33	34	40
46	HCDCY76	203917 4/8/99	Uni-ZAP XR	56	1392	628	1392		860	103	1	20	21	35
47	HCE3B04	203917 4/8/99	Uni-ZAP XR	57	1807	1347	1806		1588	104	1	14	15	32

Table 1 summarizes the information corresponding to each "Gene No." described above. The nucleotide sequence identified as "NT SEQ ID NO:X" was assembled from partially homologous ("overlapping") sequences obtained from the "cDNA clone ID" identified in Table 1 and, in some cases, from additional related DNA clones. The overlapping sequences were assembled into a single contiguous sequence of high redundancy (usually three to five overlapping sequences at each nucleotide position), resulting in a final sequence identified as SEQ ID NO:X.

The cDNA Clone ID was deposited on the date and given the corresponding deposit number listed in "ATCC Deposit No:Z and Date." Some of the deposits contain multiple different clones corresponding to the same gene. "Vector" refers to the type of vector contained in the cDNA Clone ID.

"Total NT Seq." refers to the total number of nucleotides in the contig identified by "Gene No." The deposited clone may contain all or most of these sequences, reflected by the nucleotide position indicated as "5' NT of Clone Seq." and the "3' NT of Clone Seq." of SEQ ID NO:X. The nucleotide position of SEQ ID NO:X of the putative start codon (methionine) is identified as "5' NT of Start Codon." Similarly, the nucleotide position of SEQ ID NO:X of the predicted signal sequence is identified as "5' NT of First AA of Signal Pep."

The translated amino acid sequence, beginning with the methionine, is identified as "AA SEQ ID NO:Y," although other reading frames can also be easily translated using known molecular biology techniques. The polypeptides produced by these alternative open reading frames are specifically contemplated by the present invention.

The first and last amino acid position of SEQ ID NO:Y of the predicted signal peptide is identified as "First AA of Sig Pep" and "Last AA of Sig Pep." The predicted first amino acid position of SEQ ID NO:Y of the secreted portion is identified as "Predicted First AA of Secreted Portion." Finally, the amino acid position of SEQ ID NO:Y of the last amino acid in the open reading frame is identified as "Last AA of ORF."

SEQ ID NO:X (where X may be any of the polynucleotide sequences disclosed in the sequence listing) and the translated SEQ ID NO:Y (where Y may be

any of the polypeptide sequences disclosed in the sequence listing) are sufficiently accurate and otherwise suitable for a variety of uses well known in the art and described further below. For instance, SEQ ID NO:X is useful for designing nucleic acid hybridization probes that will detect nucleic acid sequences contained in SEQ ID
5 NO:X or the cDNA contained in the deposited clone. These probes will also hybridize to nucleic acid molecules in biological samples, thereby enabling a variety of forensic and diagnostic methods of the invention. Similarly, polypeptides identified from SEQ ID NO:Y may be used, for example, to generate antibodies which bind specifically to proteins containing the polypeptides and the secreted
10 proteins encoded by the cDNA clones identified in Table 1.

Nevertheless, DNA sequences generated by sequencing reactions can contain sequencing errors. The errors exist as misidentified nucleotides, or as insertions or deletions of nucleotides in the generated DNA sequence. The erroneously inserted or deleted nucleotides cause frame shifts in the reading frames of the predicted amino
15 acid sequence. In these cases, the predicted amino acid sequence diverges from the actual amino acid sequence, even though the generated DNA sequence may be greater than 99.9% identical to the actual DNA sequence (for example, one base insertion or deletion in an open reading frame of over 1000 bases).

Accordingly, for those applications requiring precision in the nucleotide
20 sequence or the amino acid sequence, the present invention provides not only the generated nucleotide sequence identified as SEQ ID NO:X and the predicted translated amino acid sequence identified as SEQ ID NO:Y, but also a sample of plasmid DNA containing a human cDNA of the invention deposited with the ATCC, as set forth in Table 1. The nucleotide sequence of each deposited clone can readily
25 be determined by sequencing the deposited clone in accordance with known methods. The predicted amino acid sequence can then be verified from such deposits. Moreover, the amino acid sequence of the protein encoded by a particular clone can also be directly determined by peptide sequencing or by expressing the protein in a suitable host cell containing the deposited human cDNA, collecting the protein, and
30 determining its sequence.

The present invention also relates to the genes corresponding to SEQ ID NO:X, SEQ ID NO:Y, or the deposited clone. The corresponding gene can be isolated in accordance with known methods using the sequence information disclosed herein. Such methods include preparing probes or primers from the disclosed
5 sequence and identifying or amplifying the corresponding gene from appropriate sources of genomic material.

Also provided in the present invention are allelic variants, orthologs, and/or species homologs. Procedures known in the art can be used to obtain full-length genes, allelic variants, splice variants, full-length coding portions, orthologs, and/or
10 species homologs of genes corresponding to SEQ ID NO:X, SEQ ID NO:Y, or a deposited clone, using information from the sequences disclosed herein or the clones deposited with the ATCC. For example, allelic variants and/or species homologs may be isolated and identified by making suitable probes or primers from the sequences provided herein and screening a suitable nucleic acid source for allelic variants and/or
15 the desired homologue.

The polypeptides of the invention can be prepared in any suitable manner. Such polypeptides include isolated naturally occurring polypeptides, recombinantly produced polypeptides, synthetically produced polypeptides, or polypeptides produced by a combination of these methods. Means for preparing such polypeptides
20 are well understood in the art.

The polypeptides may be in the form of the secreted protein, including the mature form, or may be a part of a larger protein, such as a fusion protein (see below). It is often advantageous to include an additional amino acid sequence which contains secretory or leader sequences, pro-sequences, sequences which aid in purification ,
25 such as multiple histidine residues, or an additional sequence for stability during recombinant production.

The polypeptides of the present invention are preferably provided in an isolated form, and preferably are substantially purified. A recombinantly produced version of a polypeptide, including the secreted polypeptide, can be substantially
30 purified using techniques described herein or otherwise known in the art, such as, for example, by the one-step method described in Smith and Johnson, Gene 67:31-40

(1988). Polypeptides of the invention also can be purified from natural, synthetic or recombinant sources using techniques described herein or otherwise known in the art, such as, for example, antibodies of the invention raised against the secreted protein.

The present invention provides a polynucleotide comprising, or alternatively
5 consisting of, the nucleic acid sequence of SEQ ID NO:X, and/or a cDNA contained in ATCC deposit Z. The present invention also provides a polypeptide comprising, or alternatively, consisting of, the polypeptide sequence of SEQ ID NO:Y and/or a polypeptide encoded by the cDNA contained in ATCC deposit Z. Polynucleotides encoding a polypeptide comprising, or alternatively consisting of the polypeptide
10 sequence of SEQ ID NO:Y and/or a polypeptide sequence encoded by the cDNA contained in ATCC deposit Z are also encompassed by the invention.

Signal Sequences

The present invention also encompasses mature forms of the polypeptide having the polypeptide sequence of SEQ ID NO:Y and/or the polypeptide sequence
15 encoded by the cDNA in a deposited clone. Polynucleotides encoding the mature forms (such as, for example, the polynucleotide sequence in SEQ ID NO:X and/or the polynucleotide sequence contained in the cDNA of a deposited clone) are also encompassed by the invention. According to the signal hypothesis, proteins secreted by mammalian cells have a signal or secretory leader sequence which is cleaved from
20 the mature protein once export of the growing protein chain across the rough endoplasmic reticulum has been initiated. Most mammalian cells and even insect cells cleave secreted proteins with the same specificity. However, in some cases, cleavage of a secreted protein is not entirely uniform, which results in two or more mature species of the protein. Further, it has long been known that cleavage
25 specificity of a secreted protein is ultimately determined by the primary structure of the complete protein, that is, it is inherent in the amino acid sequence of the polypeptide.

Methods for predicting whether a protein has a signal sequence, as well as the cleavage point for that sequence, are available. For instance, the method of
30 McGeoch, Virus Res. 3:271-286 (1985), uses the information from a short N-terminal charged region and a subsequent uncharged region of the complete (uncleaved)

protein. The method of von Heinje, *Nucleic Acids Res.* 14:4683-4690 (1986) uses the information from the residues surrounding the cleavage site, typically residues -13 to +2, where +1 indicates the amino terminus of the secreted protein. The accuracy of predicting the cleavage points of known mammalian secretory proteins for each of these methods is in the range of 75-80%. (von Heinje, *supra*.) However, the two methods do not always produce the same predicted cleavage point(s) for a given protein.

In the present case, the deduced amino acid sequence of the secreted polypeptide was analyzed by a computer program called SignalP (Henrik Nielsen et al., *Protein Engineering* 10:1-6 (1997)), which predicts the cellular location of a protein based on the amino acid sequence. As part of this computational prediction of localization, the methods of McGeoch and von Heinje are incorporated. The analysis of the amino acid sequences of the secreted proteins described herein by this program provided the results shown in Table 1.

As one of ordinary skill would appreciate, however, cleavage sites sometimes vary from organism to organism and cannot be predicted with absolute certainty. Accordingly, the present invention provides secreted polypeptides having a sequence shown in SEQ ID NO:Y which have an N-terminus beginning within 5 residues (i.e., + or - 5 residues) of the predicted cleavage point. Similarly, it is also recognized that in some cases, cleavage of the signal sequence from a secreted protein is not entirely uniform, resulting in more than one secreted species. These polypeptides, and the polynucleotides encoding such polypeptides, are contemplated by the present invention.

Moreover, the signal sequence identified by the above analysis may not necessarily predict the naturally occurring signal sequence. For example, the naturally occurring signal sequence may be further upstream from the predicted signal sequence. However, it is likely that the predicted signal sequence will be capable of directing the secreted protein to the ER. Nonetheless, the present invention provides the mature protein produced by expression of the polynucleotide sequence of SEQ ID NO:X and/or the polynucleotide sequence contained in the cDNA of a deposited clone, in a mammalian cell (e.g., COS cells, as described below). These polypeptides,

and the polynucleotides encoding such polypeptides, are contemplated by the present invention.

Polynucleotide and Polypeptide Variants

5 The present invention is directed to variants of the polynucleotide sequence disclosed in SEQ ID NO:X, the complementary strand thereto, and/or the cDNA sequence contained in a deposited clone.

 The present invention also encompasses variants of the polypeptide sequence disclosed in SEQ ID NO:Y and/or encoded by a deposited clone.

10 "Variant" refers to a polynucleotide or polypeptide differing from the polynucleotide or polypeptide of the present invention, but retaining essential properties thereof. Generally, variants are overall closely similar, and, in many regions, identical to the polynucleotide or polypeptide of the present invention.

 The present invention is also directed to nucleic acid molecules which
15 comprise, or alternatively consist of, a nucleotide sequence which is at least 80%, 85%, 90%, 95%, 96%, 97%, 98% or 99% identical to, for example, the nucleotide coding sequence in SEQ ID NO:X or the complementary strand thereto, the nucleotide coding sequence contained in a deposited cDNA clone or the complementary strand thereto, a nucleotide sequence encoding the polypeptide of
20 SEQ ID NO:Y, a nucleotide sequence encoding the polypeptide encoded by the cDNA contained in a deposited clone, and/or polynucleotide fragments of any of these nucleic acid molecules (e.g., those fragments described herein).

 Polynucleotides which hybridize to these nucleic acid molecules under stringent hybridization conditions or lower stringency conditions are also encompassed by the
25 invention, as are polypeptides encoded by these polynucleotides.

 The present invention is also directed to polypeptides which comprise, or alternatively consist of, an amino acid sequence which is at least 80%, 85%, 90%, 95%, 96%, 97%, 98%, 99% identical to, for example, the polypeptide sequence shown in SEQ ID NO:Y, the polypeptide sequence encoded by the cDNA contained
30 in a deposited clone, and/or polypeptide fragments of any of these polypeptides (e.g., those fragments described herein).

By a nucleic acid having a nucleotide sequence at least, for example, 95% "identical" to a reference nucleotide sequence of the present invention, it is intended that the nucleotide sequence of the nucleic acid is identical to the reference sequence except that the nucleotide sequence may include up to five point mutations per each 100 nucleotides of the reference nucleotide sequence encoding the polypeptide. In other words, to obtain a nucleic acid having a nucleotide sequence at least 95% identical to a reference nucleotide sequence, up to 5% of the nucleotides in the reference sequence may be deleted or substituted with another nucleotide, or a number of nucleotides up to 5% of the total nucleotides in the reference sequence may be inserted into the reference sequence. The query sequence may be an entire sequence shown in Table 1, the ORF (open reading frame), or any fragment specified as described herein.

As a practical matter, whether any particular nucleic acid molecule or polypeptide is at least 80%, 85%, 90%, 95%, 96%, 97%, 98% or 99% identical to a nucleotide sequence of the present invention can be determined conventionally using known computer programs. A preferred method for determining the best overall match between a query sequence (a sequence of the present invention) and a subject sequence, also referred to as a global sequence alignment, can be determined using the FASTDB computer program based on the algorithm of Brutlag et al. (Comp. App. Biosci. 6:237-245(1990)). In a sequence alignment the query and subject sequences are both DNA sequences. An RNA sequence can be compared by converting U's to T's. The result of said global sequence alignment is in percent identity. Preferred parameters used in a FASTDB alignment of DNA sequences to calculate percent identity are: Matrix=Unitary, k-tuple=4, Mismatch Penalty=1, Joining Penalty=30, Randomization Group Length=0, Cutoff Score=1, Gap Penalty=5, Gap Size Penalty 0.05, Window Size=500 or the length of the subject nucleotide sequence, whichever is shorter.

If the subject sequence is shorter than the query sequence because of 5' or 3' deletions, not because of internal deletions, a manual correction must be made to the results. This is because the FASTDB program does not account for 5' and 3' truncations of the subject sequence when calculating percent identity. For subject

sequences truncated at the 5' or 3' ends, relative to the query sequence, the percent identity is corrected by calculating the number of bases of the query sequence that are 5' and 3' of the subject sequence, which are not matched/aligned, as a percent of the total bases of the query sequence. Whether a nucleotide is matched/aligned is
5 determined by results of the FASTDB sequence alignment. This percentage is then subtracted from the percent identity, calculated by the above FASTDB program using the specified parameters, to arrive at a final percent identity score. This corrected score is what is used for the purposes of the present invention. Only bases outside the 5' and 3' bases of the subject sequence, as displayed by the FASTDB alignment,
10 which are not matched/aligned with the query sequence, are calculated for the purposes of manually adjusting the percent identity score.

For example, a 90 base subject sequence is aligned to a 100 base query sequence to determine percent identity. The deletions occur at the 5' end of the subject sequence and therefore, the FASTDB alignment does not show a
15 matched/alignment of the first 10 bases at 5' end. The 10 unpaired bases represent 10% of the sequence (number of bases at the 5' and 3' ends not matched/total number of bases in the query sequence) so 10% is subtracted from the percent identity score calculated by the FASTDB program. If the remaining 90 bases were perfectly matched the final percent identity would be 90%. In another example, a 90 base
20 subject sequence is compared with a 100 base query sequence. This time the deletions are internal deletions so that there are no bases on the 5' or 3' of the subject sequence which are not matched/aligned with the query. In this case the percent identity calculated by FASTDB is not manually corrected. Once again, only bases 5' and 3' of the subject sequence which are not matched/aligned with the query sequence
25 are manually corrected for. No other manual corrections are to made for the purposes of the present invention.

By a polypeptide having an amino acid sequence at least, for example, 95% "identical" to a query amino acid sequence of the present invention, it is intended that the amino acid sequence of the subject polypeptide is identical to the query sequence
30 except that the subject polypeptide sequence may include up to five amino acid alterations per each 100 amino acids of the query amino acid sequence. In other

words, to obtain a polypeptide having an amino acid sequence at least 95% identical to a query amino acid sequence, up to 5% of the amino acid residues in the subject sequence may be inserted, deleted, (indels) or substituted with another amino acid.

These alterations of the reference sequence may occur at the amino or carboxy
5 terminal positions of the reference amino acid sequence or anywhere between those terminal positions, interspersed either individually among residues in the reference sequence or in one or more contiguous groups within the reference sequence.

As a practical matter, whether any particular polypeptide is at least 80%, 85%,
90%, 95%, 96%, 97%, 98% or 99% identical to, for instance, an amino acid
10 sequences shown in Table 1 (SEQ ID NO:Y) or to the amino acid sequence encoded by cDNA contained in a deposited clone can be determined conventionally using known computer programs. A preferred method for determining the best overall match between a query sequence (a sequence of the present invention) and a subject
sequence, also referred to as a global sequence alignment, can be determined using
15 the FASTDB computer program based on the algorithm of Brutlag et al. (Comp. App. Biosci. 6:237-245(1990)). In a sequence alignment the query and subject sequences are either both nucleotide sequences or both amino acid sequences. The result of said global sequence alignment is in percent identity. Preferred parameters used in a
FASTDB amino acid alignment are: Matrix=PAM 0, k-tuple=2, Mismatch
20 Penalty=1, Joining Penalty=20, Randomization Group Length=0, Cutoff Score=1, Window Size=sequence length, Gap Penalty=5, Gap Size Penalty=0.05, Window Size=500 or the length of the subject amino acid sequence, whichever is shorter.

If the subject sequence is shorter than the query sequence due to N- or C-terminal deletions, not because of internal deletions, a manual correction must be
25 made to the results. This is because the FASTDB program does not account for N- and C-terminal truncations of the subject sequence when calculating global percent identity. For subject sequences truncated at the N- and C-termini, relative to the query sequence, the percent identity is corrected by calculating the number of residues of the query sequence that are N- and C-terminal of the subject sequence, which are
30 not matched/aligned with a corresponding subject residue, as a percent of the total bases of the query sequence. Whether a residue is matched/aligned is determined by

results of the FASTDB sequence alignment. This percentage is then subtracted from the percent identity, calculated by the above FASTDB program using the specified parameters, to arrive at a final percent identity score. This final percent identity score is what is used for the purposes of the present invention. Only residues to the N- and C-termini of the subject sequence, which are not matched/aligned with the query sequence, are considered for the purposes of manually adjusting the percent identity score. That is, only query residue positions outside the farthest N- and C-terminal residues of the subject sequence.

For example, a 90 amino acid residue subject sequence is aligned with a 100 residue query sequence to determine percent identity. The deletion occurs at the N-terminus of the subject sequence and therefore, the FASTDB alignment does not show a matching/alignment of the first 10 residues at the N-terminus. The 10 unpaired residues represent 10% of the sequence (number of residues at the N- and C-termini not matched/total number of residues in the query sequence) so 10% is subtracted from the percent identity score calculated by the FASTDB program. If the remaining 90 residues were perfectly matched the final percent identity would be 90%. In another example, a 90 residue subject sequence is compared with a 100 residue query sequence. This time the deletions are internal deletions so there are no residues at the N- or C-termini of the subject sequence which are not matched/aligned with the query. In this case the percent identity calculated by FASTDB is not manually corrected. Once again, only residue positions outside the N- and C-terminal ends of the subject sequence, as displayed in the FASTDB alignment, which are not matched/aligned with the query sequence are manually corrected for. No other manual corrections are to be made for the purposes of the present invention.

The variants may contain alterations in the coding regions, non-coding regions, or both. Especially preferred are polynucleotide variants containing alterations which produce silent substitutions, additions, or deletions, but do not alter the properties or activities of the encoded polypeptide. Nucleotide variants produced by silent substitutions due to the degeneracy of the genetic code are preferred. Moreover, variants in which 5-10, 1-5, or 1-2 amino acids are substituted, deleted, or added in any combination are also preferred. Polynucleotide variants can be produced

for a variety of reasons, e.g., to optimize codon expression for a particular host (change codons in the human mRNA to those preferred by a bacterial host such as *E. coli*).

Naturally occurring variants are called "allelic variants," and refer to one of several alternate forms of a gene occupying a given locus on a chromosome of an organism. (Genes II, Lewin, B., ed., John Wiley & Sons, New York (1985).) These allelic variants can vary at either the polynucleotide and/or polypeptide level and are included in the present invention. Alternatively, non-naturally occurring variants may be produced by mutagenesis techniques or by direct synthesis.

Using known methods of protein engineering and recombinant DNA technology, variants may be generated to improve or alter the characteristics of the polypeptides of the present invention. For instance, one or more amino acids can be deleted from the N-terminus or C-terminus of the secreted protein without substantial loss of biological function. The authors of Ron et al., *J. Biol. Chem.* 268: 2984-2988 (1993), reported variant KGF proteins having heparin binding activity even after deleting 3, 8, or 27 amino-terminal amino acid residues. Similarly, Interferon gamma exhibited up to ten times higher activity after deleting 8-10 amino acid residues from the carboxy terminus of this protein. (Dobeli et al., *J. Biotechnology* 7:199-216 (1988).)

Moreover, ample evidence demonstrates that variants often retain a biological activity similar to that of the naturally occurring protein. For example, Gayle and coworkers (*J. Biol. Chem.* 268:22105-22111 (1993)) conducted extensive mutational analysis of human cytokine IL-1a. They used random mutagenesis to generate over 3,500 individual IL-1a mutants that averaged 2.5 amino acid changes per variant over the entire length of the molecule. Multiple mutations were examined at every possible amino acid position. The investigators found that "[m]ost of the molecule could be altered with little effect on either [binding or biological activity]." (See, Abstract.) In fact, only 23 unique amino acid sequences, out of more than 3,500 nucleotide sequences examined, produced a protein that significantly differed in activity from wild-type.

Furthermore, even if deleting one or more amino acids from the N-terminus or C-terminus of a polypeptide results in modification or loss of one or more biological functions, other biological activities may still be retained. For example, the ability of a deletion variant to induce and/or to bind antibodies which recognize the secreted form will likely be retained when less than the majority of the residues of the secreted form are removed from the N-terminus or C-terminus. Whether a particular polypeptide lacking N- or C-terminal residues of a protein retains such immunogenic activities can readily be determined by routine methods described herein and otherwise known in the art.

Thus, the invention further includes polypeptide variants which show substantial biological activity. Such variants include deletions, insertions, inversions, repeats, and substitutions selected according to general rules known in the art so as to have little effect on activity. For example, guidance concerning how to make phenotypically silent amino acid substitutions is provided in Bowie et al., Science 247:1306-1310 (1990), wherein the authors indicate that there are two main strategies for studying the tolerance of an amino acid sequence to change.

The first strategy exploits the tolerance of amino acid substitutions by natural selection during the process of evolution. By comparing amino acid sequences in different species, conserved amino acids can be identified. These conserved amino acids are likely important for protein function. In contrast, the amino acid positions where substitutions have been tolerated by natural selection indicates that these positions are not critical for protein function. Thus, positions tolerating amino acid substitution could be modified while still maintaining biological activity of the protein.

The second strategy uses genetic engineering to introduce amino acid changes at specific positions of a cloned gene to identify regions critical for protein function. For example, site directed mutagenesis or alanine-scanning mutagenesis (introduction of single alanine mutations at every residue in the molecule) can be used. (Cunningham and Wells, Science 244:1081-1085 (1989).) The resulting mutant molecules can then be tested for biological activity.

As the authors state, these two strategies have revealed that proteins are surprisingly tolerant of amino acid substitutions. The authors further indicate which amino acid changes are likely to be permissive at certain amino acid positions in the protein. For example, most buried (within the tertiary structure of the protein) amino acid residues require nonpolar side chains, whereas few features of surface side chains are generally conserved. Moreover, tolerated conservative amino acid substitutions involve replacement of the aliphatic or hydrophobic amino acids Ala, Val, Leu and Ile; replacement of the hydroxyl residues Ser and Thr; replacement of the acidic residues Asp and Glu; replacement of the amide residues Asn and Gln, replacement of the basic residues Lys, Arg, and His; replacement of the aromatic residues Phe, Tyr, and Trp, and replacement of the small-sized amino acids Ala, Ser, Thr, Met, and Gly.

Besides conservative amino acid substitution, variants of the present invention include (i) substitutions with one or more of the non-conserved amino acid residues, where the substituted amino acid residues may or may not be one encoded by the genetic code, or (ii) substitution with one or more of amino acid residues having a substituent group, or (iii) fusion of the mature polypeptide with another compound, such as a compound to increase the stability and/or solubility of the polypeptide (for example, polyethylene glycol), or (iv) fusion of the polypeptide with additional amino acids, such as, for example, an IgG Fc fusion region peptide, or leader or secretory sequence, or a sequence facilitating purification. Such variant polypeptides are deemed to be within the scope of those skilled in the art from the teachings herein.

For example, polypeptide variants containing amino acid substitutions of charged amino acids with other charged or neutral amino acids may produce proteins with improved characteristics, such as less aggregation. Aggregation of pharmaceutical formulations both reduces activity and increases clearance due to the aggregate's immunogenic activity. (Pinckard et al., Clin. Exp. Immunol. 2:331-340 (1967); Robbins et al., Diabetes 36: 838-845 (1987); Cleland et al., Crit. Rev. Therapeutic Drug Carrier Systems 10:307-377 (1993).)

A further embodiment of the invention relates to a polypeptide which comprises the amino acid sequence of the present invention having an amino acid sequence which contains at least one amino acid substitution, but not more than 50

amino acid substitutions, even more preferably, not more than 40 amino acid substitutions, still more preferably, not more than 30 amino acid substitutions, and still even more preferably, not more than 20 amino acid substitutions. Of course, in order of ever-increasing preference, it is highly preferable for a peptide or polypeptide to have an amino acid sequence which comprises the amino acid sequence of the present invention, which contains at least one, but not more than 10, 9, 8, 7, 6, 5, 4, 3, 2 or 1 amino acid substitutions. In specific embodiments, the number of additions, substitutions, and/or deletions in the amino acid sequence of the present invention or fragments thereof (e.g., the mature form and/or other fragments described herein), is 1-5, 5-10, 5-25, 5-50, 10-50 or 50-150, conservative amino acid substitutions are preferable.

Polynucleotide and Polypeptide Fragments

The present invention is also directed to polynucleotide fragments of the polynucleotides of the invention.

In the present invention, a "polynucleotide fragment" refers to a short polynucleotide having a nucleic acid sequence which: is a portion of that contained in a deposited clone, or encoding the polypeptide encoded by the cDNA in a deposited clone; is a portion of that shown in SEQ ID NO:X or the complementary strand thereto, or is a portion of a polynucleotide sequence encoding the polypeptide of SEQ ID NO:Y. The nucleotide fragments of the invention are preferably at least about 15 nt, and more preferably at least about 20 nt, still more preferably at least about 30 nt, and even more preferably, at least about 40 nt, at least about 50 nt, at least about 75 nt, or at least about 150 nt in length. A fragment "at least 20 nt in length," for example, is intended to include 20 or more contiguous bases from the cDNA sequence contained in a deposited clone or the nucleotide sequence shown in SEQ ID NO:X. In this context "about" includes the particularly recited value, a value larger or smaller by several (5, 4, 3, 2, or 1) nucleotides, at either terminus or at both termini. These nucleotide fragments have uses that include, but are not limited to, as diagnostic probes and primers as discussed herein. Of course, larger fragments (e.g., 50, 150, 500, 600, 2000 nucleotides) are preferred.

Moreover, representative examples of polynucleotide fragments of the invention, include, for example, fragments comprising, or alternatively consisting of, a sequence from about nucleotide number 1-50, 51-100, 101-150, 151-200, 201-250, 251-300, 301-350, 351-400, 401-450, 451-500, 501-550, 551-600, 651-700, 701-750, 751-800, 800-850, 851-900, 901-950, 951-1000, 1001-1050, 1051-1100, 1101-1150, 1151-1200, 1201-1250, 1251-1300, 1301-1350, 1351-1400, 1401-1450, 1451-1500, 1501-1550, 1551-1600, 1601-1650, 1651-1700, 1701-1750, 1751-1800, 1801-1850, 1851-1900, 1901-1950, 1951-2000, or 2001 to the end of SEQ ID NO:X, or the complementary strand thereto, or the cDNA contained in a deposited clone. In this context "about" includes the particularly recited ranges, and ranges larger or smaller by several (5, 4, 3, 2, or 1) nucleotides, at either terminus or at both termini.

Preferably, these fragments encode a polypeptide which has biological activity. More preferably, these polynucleotides can be used as probes or primers as discussed herein. Polynucleotides which hybridize to these nucleic acid molecules under stringent hybridization conditions or lower stringency conditions are also encompassed by the invention, as are polypeptides encoded by these polynucleotides.

In the present invention, a "polypeptide fragment" refers to an amino acid sequence which is a portion of that contained in SEQ ID NO:Y or encoded by the cDNA contained in a deposited clone. Protein (polypeptide) fragments may be "free-standing," or comprised within a larger polypeptide of which the fragment forms a part or region, most preferably as a single continuous region. Representative examples of polypeptide fragments of the invention, include, for example, fragments comprising, or alternatively consisting of, from about amino acid number 1-20, 21-40, 41-60, 61-80, 81-100, 102-120, 121-140, 141-160, or 161 to the end of the coding region. Moreover, polypeptide fragments can be about 20, 30, 40, 50, 60, 70, 80, 90, 100, 110, 120, 130, 140, or 150 amino acids in length. In this context "about" includes the particularly recited ranges or values, and ranges or values larger or smaller by several (5, 4, 3, 2, or 1) amino acids, at either extreme or at both extremes. Polynucleotides encoding these polypeptides are also encompassed by the invention.

Preferred polypeptide fragments include the secreted protein as well as the mature form. Further preferred polypeptide fragments include the secreted protein or

the mature form having a continuous series of deleted residues from the amino or the carboxy terminus, or both. For example, any number of amino acids, ranging from 1-60, can be deleted from the amino terminus of either the secreted polypeptide or the mature form. Similarly, any number of amino acids, ranging from 1-30, can be
5 deleted from the carboxy terminus of the secreted protein or mature form.

Furthermore, any combination of the above amino and carboxy terminus deletions are preferred. Similarly, polynucleotides encoding these polypeptide fragments are also preferred.

Also preferred are polypeptide and polynucleotide fragments characterized by
10 structural or functional domains, such as fragments that comprise alpha-helix and alpha-helix forming regions, beta-sheet and beta-sheet-forming regions, turn and turn-forming regions, coil and coil-forming regions, hydrophilic regions, hydrophobic regions, alpha amphipathic regions, beta amphipathic regions, flexible regions, surface-forming regions, substrate binding region, and high antigenic index regions.
15 Polypeptide fragments of SEQ ID NO:Y falling within conserved domains are specifically contemplated by the present invention. Moreover, polynucleotides encoding these domains are also contemplated.

Other preferred polypeptide fragments are biologically active fragments. Biologically active fragments are those exhibiting activity similar, but not necessarily
20 identical, to an activity of the polypeptide of the present invention. The biological activity of the fragments may include an improved desired activity, or a decreased undesirable activity. Polynucleotides encoding these polypeptide fragments are also encompassed by the invention.

Preferably, the polynucleotide fragments of the invention encode a
25 polypeptide which demonstrates a functional activity. By a polypeptide demonstrating a "functional activity" is meant, a polypeptide capable of displaying one or more known functional activities associated with a full-length (complete) polypeptide of invention protein. Such functional activities include, but are not limited to, biological activity, antigenicity [ability to bind (or compete with a
30 polypeptide of the invention for binding) to an antibody to the polypeptide of the invention], immunogenicity (ability to generate antibody which binds to a polypeptide

of the invention), ability to form multimers with polypeptides of the invention, and ability to bind to a receptor or ligand for a polypeptide of the invention.

The functional activity of polypeptides of the invention, and fragments, variants derivatives, and analogs thereof, can be assayed by various methods.

5 For example, in one embodiment where one is assaying for the ability to bind or compete with full-length polypeptide of the invention for binding to an antibody of the polypeptide of the invention, various immunoassays known in the art can be used, including but not limited to, competitive and non-competitive assay systems using techniques such as radioimmunoassays, ELISA (enzyme linked immunosorbent
10 assay), "sandwich" immunoassays, immunoradiometric assays, gel diffusion precipitation reactions, immunodiffusion assays, in situ immunoassays (using colloidal gold, enzyme or radioisotope labels, for example), western blots, precipitation reactions, agglutination assays (e.g., gel agglutination assays, hemagglutination assays), complement fixation assays, immunofluorescence assays,
15 protein A assays, and immunoelectrophoresis assays, etc. In one embodiment, antibody binding is detected by detecting a label on the primary antibody. In another embodiment, the primary antibody is detected by detecting binding of a secondary antibody or reagent to the primary antibody. In a further embodiment, the secondary antibody is labeled. Many means are known in the art for detecting binding in an
20 immunoassay and are within the scope of the present invention.

In another embodiment, where a ligand for a polypeptide of the invention identified, or the ability of a polypeptide fragment, variant or derivative of the invention to multimerize is being evaluated, binding can be assayed, e.g., by means well-known in the art, such as, for example, reducing and non-reducing gel
25 chromatography, protein affinity chromatography, and affinity blotting. See generally, Phizicky, E., et al., 1995, Microbiol. Rev. 59:94-123. In another embodiment, physiological correlates of binding of a polypeptide of the invention to its substrates (signal transduction) can be assayed.

In addition, assays described herein (see Examples) and otherwise known in
30 the art may routinely be applied to measure the ability of polypeptides of the invention and fragments, variants derivatives and analogs thereof to elicit related

biological activity related to that of the polypeptide of the invention (either in vitro or in vivo). Other methods will be known to the skilled artisan and are within the scope of the invention.

5

Epitopes and Antibodies

The present invention encompasses polypeptides comprising, or alternatively consisting of, an epitope of the polypeptide having an amino acid sequence of SEQ ID NO:Y, or an epitope of the polypeptide sequence encoded by a polynucleotide sequence contained in ATCC deposit No. Z or encoded by a polynucleotide that
10 hybridizes to the complement of the sequence of SEQ ID NO:X or contained in ATCC deposit No. Z under stringent hybridization conditions or lower stringency hybridization conditions as defined supra. The present invention further encompasses polynucleotide sequences encoding an epitope of a polypeptide sequence of the invention (such as, for example, the sequence disclosed in SEQ ID NO:X),
15 polynucleotide sequences of the complementary strand of a polynucleotide sequence encoding an epitope of the invention, and polynucleotide sequences which hybridize to the complementary strand under stringent hybridization conditions or lower stringency hybridization conditions defined supra.

The term "epitopes," as used herein, refers to portions of a polypeptide having
20 antigenic or immunogenic activity in an animal, preferably a mammal, and most preferably in a human. In a preferred embodiment, the present invention encompasses a polypeptide comprising an epitope, as well as the polynucleotide encoding this polypeptide. An "immunogenic epitope," as used herein, is defined as a portion of a protein that elicits an antibody response in an animal, as determined by
25 any method known in the art, for example, by the methods for generating antibodies described infra. (See, for example, Geysen et al., Proc. Natl. Acad. Sci. USA 81:3998- 4002 (1983)). The term "antigenic epitope," as used herein, is defined as a portion of a protein to which an antibody can immunospecifically bind its antigen as determined by any method well known in the art, for example, by the immunoassays
30 described herein. Immunospecific binding excludes non-specific binding but does not

necessarily exclude cross- reactivity with other antigens. Antigenic epitopes need not necessarily be immunogenic.

Fragments which function as epitopes may be produced by any conventional means. (See, e.g., Houghten, Proc. Natl. Acad. Sci. USA 82:5131-5135 (1985),
5 further described in U.S. Patent No. 4,631,211).

In the present invention, antigenic epitopes preferably contain a sequence of at least 4, at least 5, at least 6, at least 7, more preferably at least 8, at least 9, at least 10, at least 11, at least 12, at least 13, at least 14, at least 15, at least 20, at least 25, at least 30, at least 40, at least 50, and, most preferably, between about 15 to about 30
10 amino acids. Preferred polypeptides comprising immunogenic or antigenic epitopes are at least 10, 15, 20, 25, 30, 35, 40, 45, 50, 55, 60, 65, 70, 75, 80, 85, 90, 95, or 100 amino acid residues in length. Additional non-exclusive preferred antigenic epitopes include the antigenic epitopes disclosed herein, as well as portions thereof. Antigenic epitopes are useful, for example, to raise antibodies, including monoclonal antibodies,
15 that specifically bind the epitope. Preferred antigenic epitopes include the antigenic epitopes disclosed herein, as well as any combination of two, three, four, five or more of these antigenic epitopes. Antigenic epitopes can be used as the target molecules in immunoassays. (See, for instance, Wilson et al., Cell 37:767-778 (1984); Sutcliffe et al., Science 219:660-666 (1983)).

20 Similarly, immunogenic epitopes can be used, for example, to induce antibodies according to methods well known in the art. (See, for instance, Sutcliffe et al., supra; Wilson et al., supra; Chow et al., Proc. Natl. Acad. Sci. USA 82:910-914; and Bittle et al., J. Gen. Virol. 66:2347-2354 (1985). Preferred immunogenic epitopes include the immunogenic epitopes disclosed herein, as well as any
25 combination of two, three, four, five or more of these immunogenic epitopes. The polypeptides comprising one or more immunogenic epitopes may be presented for eliciting an antibody response together with a carrier protein, such as an albumin, to an animal system (such as rabbit or mouse), or, if the polypeptide is of sufficient length (at least about 25 amino acids), the polypeptide may be presented without a
30 carrier. However, immunogenic epitopes comprising as few as 8 to 10 amino acids

have been shown to be sufficient to raise antibodies capable of binding to, at the very least, linear epitopes in a denatured polypeptide (e.g., in Western blotting).

Epitope-bearing polypeptides of the present invention may be used to induce antibodies according to methods well known in the art including, but not limited to, in vivo immunization, in vitro immunization, and phage display methods. See, e.g., Sutcliffe et al., supra; Wilson et al., supra, and Bittle et al., J. Gen. Virol., 66:2347-2354 (1985). If in vivo immunization is used, animals may be immunized with free peptide; however, anti-peptide antibody titer may be boosted by coupling the peptide to a macromolecular carrier, such as keyhole limpet hemacyanin (KLH) or tetanus toxoid. For instance, peptides containing cysteine residues may be coupled to a carrier using a linker such as maleimidobenzoyl- N-hydroxysuccinimide ester (MBS), while other peptides may be coupled to carriers using a more general linking agent such as glutaraldehyde. Animals such as rabbits, rats and mice are immunized with either free or carrier- coupled peptides, for instance, by intraperitoneal and/or intradermal injection of emulsions containing about 100 µg of peptide or carrier protein and Freund's adjuvant or any other adjuvant known for stimulating an immune response. Several booster injections may be needed, for instance, at intervals of about two weeks, to provide a useful titer of anti-peptide antibody which can be detected, for example, by ELISA assay using free peptide adsorbed to a solid surface. The titer of anti-peptide antibodies in serum from an immunized animal may be increased by selection of anti-peptide antibodies, for instance, by adsorption to the peptide on a solid support and elution of the selected antibodies according to methods well known in the art.

As one of skill in the art will appreciate, and as discussed above, the polypeptides of the present invention comprising an immunogenic or antigenic epitope can be fused to other polypeptide sequences. For example, the polypeptides of the present invention may be fused with the constant domain of immunoglobulins (IgA, IgE, IgG, IgM), or portions thereof (CH1, CH2, CH3, or any combination thereof and portions thereof) resulting in chimeric polypeptides. Such fusion proteins may facilitate purification and may increase half-life in vivo. This has been shown for chimeric proteins consisting of the first two domains of the human CD4-

polypeptide and various domains of the constant regions of the heavy or light chains of mammalian immunoglobulins. See, e.g., EP 394,827; Traunecker et al., *Nature*, 331:84-86 (1988). Enhanced delivery of an antigen across the epithelial barrier to the immune system has been demonstrated for antigens (e.g., insulin) conjugated to an FcRn binding partner such as IgG or Fc fragments (see, e.g., PCT Publications WO 96/22024 and WO 99/04813). IgG Fusion proteins that have a disulfide-linked dimeric structure due to the IgG portion disulfide bonds have also been found to be more efficient in binding and neutralizing other molecules than monomeric polypeptides or fragments thereof alone. See, e.g., Fountoulakis et al., *J. Biochem.*, 120:3958-3964 (1995). Nucleic acids encoding the above epitopes can also be recombined with a gene of interest as an epitope tag (e.g., the hemagglutinin ("HA") tag or flag tag) to aid in detection and purification of the expressed polypeptide. For example, a system described by Janknecht et al. allows for the ready purification of non-denatured fusion proteins expressed in human cell lines (Janknecht et al., 1991, *Proc. Natl. Acad. Sci. USA* 88:8972- 897). In this system, the gene of interest is subcloned into a vaccinia recombination plasmid such that the open reading frame of the gene is translationally fused to an amino-terminal tag consisting of six histidine residues. The tag serves as a matrix binding domain for the fusion protein. Extracts from cells infected with the recombinant vaccinia virus are loaded onto Ni²⁺-nitriloacetic acid-agarose column and histidine-tagged proteins can be selectively eluted with imidazole-containing buffers.

Additional fusion proteins of the invention may be generated through the techniques of gene-shuffling, motif-shuffling, exon-shuffling, and/or codon-shuffling (collectively referred to as "DNA shuffling"). DNA shuffling may be employed to modulate the activities of polypeptides of the invention, such methods can be used to generate polypeptides with altered activity, as well as agonists and antagonists of the polypeptides. See, generally, U.S. Patent Nos. 5,605,793; 5,811,238; 5,830,721; 5,834,252; and 5,837,458, and Patten et al., *Curr. Opin. Biotechnol.* 8:724-33 (1997); Harayama, *Trends Biotechnol.* 16(2):76-82 (1998); Hansson, et al., *J. Mol. Biol.* 287:265-76 (1999); and Lorenzo and Blasco, *Biotechniques* 24(2):308- 13 (1998) (each of these patents and publications are hereby incorporated by reference in

its entirety). In one embodiment, alteration of polynucleotides corresponding to SEQ ID NO:X and the polypeptides encoded by these polynucleotides may be achieved by DNA shuffling. DNA shuffling involves the assembly of two or more DNA segments by homologous or site-specific recombination to generate variation in the polynucleotide sequence. In another embodiment, polynucleotides of the invention, or the encoded polypeptides, may be altered by being subjected to random mutagenesis by error-prone PCR, random nucleotide insertion or other methods prior to recombination. In another embodiment, one or more components, motifs, sections, parts, domains, fragments, etc., of a polynucleotide encoding a polypeptide of the invention may be recombined with one or more components, motifs, sections, parts, domains, fragments, etc. of one or more heterologous molecules.

Antibodies

Further polypeptides of the invention relate to antibodies and T-cell antigen receptors (TCR) which immunospecifically bind a polypeptide, polypeptide fragment, or variant of SEQ ID NO:Y, and/or an epitope, of the present invention (as determined by immunoassays well known in the art for assaying specific antibody-antigen binding). Antibodies of the invention include, but are not limited to, polyclonal, monoclonal, multispecific, human, humanized or chimeric antibodies, single chain antibodies, Fab fragments, F(ab') fragments, fragments produced by a Fab expression library, anti-idiotypic (anti-Id) antibodies (including, e.g., anti-Id antibodies to antibodies of the invention), and epitope-binding fragments of any of the above. The term "antibody," as used herein, refers to immunoglobulin molecules and immunologically active portions of immunoglobulin molecules, i.e., molecules that contain an antigen binding site that immunospecifically binds an antigen. The immunoglobulin molecules of the invention can be of any type (e.g., IgG, IgE, IgM, IgD, IgA and IgY), class (e.g., IgG1, IgG2, IgG3, IgG4, IgA1 and IgA2) or subclass of immunoglobulin molecule.

Most preferably the antibodies are human antigen-binding antibody fragments of the present invention and include, but are not limited to, Fab, Fab' and F(ab')₂, Fd, single-chain Fvs (scFv), single-chain antibodies, disulfide-linked Fvs (sdFv) and

fragments comprising either a VL or VH domain. Antigen-binding antibody fragments, including single-chain antibodies, may comprise the variable region(s) alone or in combination with the entirety or a portion of the following: hinge region, CH1, CH2, and CH3 domains. Also included in the invention are antigen-binding
5 fragments also comprising any combination of variable region(s) with a hinge region, CH1, CH2, and CH3 domains. The antibodies of the invention may be from any animal origin including birds and mammals. Preferably, the antibodies are human, murine (e.g., mouse and rat), donkey, ship rabbit, goat, guinea pig, camel, horse, or chicken. As used herein, "human" antibodies include antibodies having the amino
10 acid sequence of a human immunoglobulin and include antibodies isolated from human immunoglobulin libraries or from animals transgenic for one or more human immunoglobulin and that do not express endogenous immunoglobulins, as described infra and, for example in, U.S. Patent No. 5,939,598 by Kucherlapati et al.

The antibodies of the present invention may be monospecific, bispecific,
15 trispecific or of greater multispecificity. Multispecific antibodies may be specific for different epitopes of a polypeptide of the present invention or may be specific for both a polypeptide of the present invention as well as for a heterologous epitope, such as a heterologous polypeptide or solid support material. See, e.g., PCT publications WO 93/17715; WO 92/08802; WO 91/00360; WO 92/05793; Tutt, et al., J. Immunol.
20 147:60-69 (1991); U.S. Patent Nos. 4,474,893; 4,714,681; 4,925,648; 5,573,920; 5,601,819; Kostelny et al., J. Immunol. 148:1547-1553 (1992).

Antibodies of the present invention may be described or specified in terms of the epitope(s) or portion(s) of a polypeptide of the present invention which they recognize or specifically bind. The epitope(s) or polypeptide portion(s) may be
25 specified as described herein, e.g., by N-terminal and C-terminal positions, by size in contiguous amino acid residues, or listed in the Tables and Figures. Antibodies which specifically bind any epitope or polypeptide of the present invention may also be excluded. Therefore, the present invention includes antibodies that specifically bind polypeptides of the present invention, and allows for the exclusion of the same.

30 Antibodies of the present invention may also be described or specified in terms of their cross-reactivity. Antibodies that do not bind any other analog,

ortholog, or homolog of a polypeptide of the present invention are included.

Antibodies that bind polypeptides with at least 95%, at least 90%, at least 85%, at least 80%, at least 75%, at least 70%, at least 65%, at least 60%, at least 55%, and at least 50% identity (as calculated using methods known in the art and described

5 herein) to a polypeptide of the present invention are also included in the present invention. In specific embodiments, antibodies of the present invention cross-react with murine, rat and/or rabbit homologs of human proteins and the corresponding epitopes thereof. Antibodies that do not bind polypeptides with less than 95%, less than 90%, less than 85%, less than 80%, less than 75%, less than 70%, less than 65%,
10 less than 60%, less than 55%, and less than 50% identity (as calculated using methods known in the art and described herein) to a polypeptide of the present invention are also included in the present invention. In a specific embodiment, the above-described cross-reactivity is with respect to any single specific antigenic or immunogenic polypeptide, or combination(s) of 2, 3, 4, 5, or more of the specific
15 antigenic and/or immunogenic polypeptides disclosed herein. Further included in the present invention are antibodies which bind polypeptides encoded by polynucleotides which hybridize to a polynucleotide of the present invention under stringent hybridization conditions (as described herein). Antibodies of the present invention may also be described or specified in terms of their binding affinity to a polypeptide
20 of the invention. Preferred binding affinities include those with a dissociation constant or K_d less than 5×10^{-2} M, 10^{-2} M, 5×10^{-3} M, 10^{-3} M, 5×10^{-4} M, 10^{-4} M, 5×10^{-5} M, 10^{-5} M, 5×10^{-6} M, 10^{-6} M, 5×10^{-7} M, 10^{-7} M, 5×10^{-8} M, 10^{-8} M, 5×10^{-9} M, 10^{-9} M, 5×10^{-10} M, 10^{-10} M, 5×10^{-11} M, 10^{-11} M, 5×10^{-12} M, 10^{-12} M, 5×10^{-13} M, 10^{-13} M, 5×10^{-14} M, 10^{-14} M, 5×10^{-15} M, or 10^{-15} M.

25 The invention also provides antibodies that competitively inhibit binding of an antibody to an epitope of the invention as determined by any method known in the art for determining competitive binding, for example, the immunoassays described herein. In preferred embodiments, the antibody competitively inhibits binding to the epitope by at least 95%, at least 90%, at least 85 %, at least 80%, at least 75%, at least
30 70%, at least 60%, or at least 50%.

Antibodies of the present invention may act as agonists or antagonists of the polypeptides of the present invention. For example, the present invention includes antibodies which disrupt the receptor/ligand interactions with the polypeptides of the invention either partially or fully. Preferably, antibodies of the present invention
5 bind an antigenic epitope disclosed herein, or a portion thereof. The invention features both receptor-specific antibodies and ligand-specific antibodies. The invention also features receptor-specific antibodies which do not prevent ligand binding but prevent receptor activation. Receptor activation (i.e., signaling) may be determined by techniques described herein or otherwise known in the art. For
10 example, receptor activation can be determined by detecting the phosphorylation (e.g., tyrosine or serine/threonine) of the receptor or its substrate by immunoprecipitation followed by western blot analysis (for example, as described supra). In specific embodiments, antibodies are provided that inhibit ligand activity or receptor activity by at least 95%, at least 90%, at least 85%, at least 80%, at least
15 75%, at least 70%, at least 60%, or at least 50% of the activity in absence of the antibody.

The invention also features receptor-specific antibodies which both prevent ligand binding and receptor activation as well as antibodies that recognize the receptor-ligand complex, and, preferably, do not specifically recognize the unbound
20 receptor or the unbound ligand. Likewise, included in the invention are neutralizing antibodies which bind the ligand and prevent binding of the ligand to the receptor, as well as antibodies which bind the ligand, thereby preventing receptor activation, but do not prevent the ligand from binding the receptor. Further included in the invention are antibodies which activate the receptor. These antibodies may act as receptor
25 agonists, i.e., potentiate or activate either all or a subset of the biological activities of the ligand-mediated receptor activation, for example, by inducing dimerization of the receptor. The antibodies may be specified as agonists, antagonists or inverse agonists for biological activities comprising the specific biological activities of the peptides of the invention disclosed herein. The above antibody agonists can be made using
30 methods known in the art. See, e.g., PCT publication WO 96/40281; U.S. Patent No. 5,811,097; Deng et al., Blood 92(6):1981-1988 (1998); Chen et al., Cancer Res.

58(16):3668-3678 (1998); Harrop et al., J. Immunol. 161(4):1786-1794 (1998); Zhu et al., Cancer Res. 58(15):3209-3214 (1998); Yoon et al., J. Immunol. 160(7):3170-3179 (1998); Prat et al., J. Cell. Sci. 111(Pt2):237-247 (1998); Pitard et al., J. Immunol. Methods 205(2):177-190 (1997); Liautard et al., Cytokine 9(4):233-241 (1997); Carlson et al., J. Biol. Chem. 272(17):11295-11301 (1997); Taryman et al., Neuron 14(4):755-762 (1995); Muller et al., Structure 6(9):1153-1167 (1998); Bartunek et al., Cytokine 8(1):14-20 (1996) (which are all incorporated by reference herein in their entireties).

Antibodies of the present invention may be used, for example, but not limited to, to purify, detect, and target the polypeptides of the present invention, including both in vitro and in vivo diagnostic and therapeutic methods. For example, the antibodies have use in immunoassays for qualitatively and quantitatively measuring levels of the polypeptides of the present invention in biological samples. See, e.g., Harlow et al., Antibodies: A Laboratory Manual, (Cold Spring Harbor Laboratory Press, 2nd ed. 1988) (incorporated by reference herein in its entirety).

As discussed in more detail below, the antibodies of the present invention may be used either alone or in combination with other compositions. The antibodies may further be recombinantly fused to a heterologous polypeptide at the N- or C-terminus or chemically conjugated (including covalently and non-covalently conjugations) to polypeptides or other compositions. For example, antibodies of the present invention may be recombinantly fused or conjugated to molecules useful as labels in detection assays and effector molecules such as heterologous polypeptides, drugs, radionuclides, or toxins. See, e.g., PCT publications WO 92/08495; WO 91/14438; WO 89/12624; U.S. Patent No. 5,314,995; and EP 396,387.

The antibodies of the invention include derivatives that are modified, i.e., by the covalent attachment of any type of molecule to the antibody such that covalent attachment does not prevent the antibody from generating an anti-idiotypic response. For example, but not by way of limitation, the antibody derivatives include antibodies that have been modified, e.g., by glycosylation, acetylation, pegylation, phosphorylation, amidation, derivatization by known protecting/blocking groups, proteolytic cleavage, linkage to a cellular ligand or other protein, etc. Any of

numerous chemical modifications may be carried out by known techniques, including, but not limited to specific chemical cleavage, acetylation, formylation, metabolic synthesis of tunicamycin, etc. Additionally, the derivative may contain one or more non-classical amino acids.

5 The antibodies of the present invention may be generated by any suitable method known in the art. Polyclonal antibodies to an antigen-of- interest can be produced by various procedures well known in the art. For example, a polypeptide of the invention can be administered to various host animals including, but not limited to, rabbits, mice, rats, etc. to induce the production of sera containing polyclonal
10 antibodies specific for the antigen. Various adjuvants may be used to increase the immunological response, depending on the host species, and include but are not limited to, Freund's (complete and incomplete), mineral gels such as aluminum hydroxide, surface active substances such as lysolecithin, pluronic polyols, polyanions, peptides, oil emulsions, keyhole limpet hemocyanins, dinitrophenol, and
15 potentially useful human adjuvants such as BCG (bacille Calmette-Guerin) and corynebacterium parvum. Such adjuvants are also well known in the art.

Monoclonal antibodies can be prepared using a wide variety of techniques known in the art including the use of hybridoma, recombinant, and phage display technologies, or a combination thereof. For example, monoclonal antibodies can be
20 produced using hybridoma techniques including those known in the art and taught, for example, in Harlow et al., *Antibodies: A Laboratory Manual*, (Cold Spring Harbor Laboratory Press, 2nd ed. 1988); Hammerling, et al., in: *Monoclonal Antibodies and T-Cell Hybridomas* 563-681 (Elsevier, N.Y., 1981) (said references incorporated by reference in their entireties). The term "monoclonal antibody" as used herein is not
25 limited to antibodies produced through hybridoma technology. The term "monoclonal antibody" refers to an antibody that is derived from a single clone, including any eukaryotic, prokaryotic, or phage clone, and not the method by which it is produced.

Methods for producing and screening for specific antibodies using hybridoma
30 technology are routine and well known in the art and are discussed in detail in the Examples (e.g., Example 16). In a non-limiting example, mice can be immunized

with a polypeptide of the invention or a cell expressing such peptide. Once an immune response is detected, e.g., antibodies specific for the antigen are detected in the mouse serum, the mouse spleen is harvested and splenocytes isolated. The splenocytes are then fused by well known techniques to any suitable myeloma cells, for example cells from cell line SP20 available from the ATCC. Hybridomas are selected and cloned by limited dilution. The hybridoma clones are then assayed by methods known in the art for cells that secrete antibodies capable of binding a polypeptide of the invention. Ascites fluid, which generally contains high levels of antibodies, can be generated by immunizing mice with positive hybridoma clones.

Accordingly, the present invention provides methods of generating monoclonal antibodies as well as antibodies produced by the method comprising culturing a hybridoma cell secreting an antibody of the invention wherein, preferably, the hybridoma is generated by fusing splenocytes isolated from a mouse immunized with an antigen of the invention with myeloma cells and then screening the hybridomas resulting from the fusion for hybridoma clones that secrete an antibody able to bind a polypeptide of the invention.

Antibody fragments which recognize specific epitopes may be generated by known techniques. For example, Fab and F(ab')₂ fragments of the invention may be produced by proteolytic cleavage of immunoglobulin molecules, using enzymes such as papain (to produce Fab fragments) or pepsin (to produce F(ab')₂ fragments). F(ab')₂ fragments contain the variable region, the light chain constant region and the CH1 domain of the heavy chain.

For example, the antibodies of the present invention can also be generated using various phage display methods known in the art. In phage display methods, functional antibody domains are displayed on the surface of phage particles which carry the polynucleotide sequences encoding them. In a particular embodiment, such phage can be utilized to display antigen binding domains expressed from a repertoire or combinatorial antibody library (e.g., human or murine). Phage expressing an antigen binding domain that binds the antigen of interest can be selected or identified with antigen, e.g., using labeled antigen or antigen bound or captured to a solid surface or bead. Phage used in these methods are typically filamentous phage

including fd and M13 binding domains expressed from phage with Fab, Fv or disulfide stabilized Fv antibody domains recombinantly fused to either the phage gene III or gene VIII protein. Examples of phage display methods that can be used to make the antibodies of the present invention include those disclosed in Brinkman et al., J. Immunol. Methods 182:41-50 (1995); Ames et al., J. Immunol. Methods 184:177-186 (1995); Kettleborough et al., Eur. J. Immunol. 24:952-958 (1994); Persic et al., Gene 187 9-18 (1997); Burton et al., Advances in Immunology 57:191-280 (1994); PCT application No. PCT/GB91/01134; PCT publications WO 90/02809; WO 91/10737; WO 92/01047; WO 92/18619; WO 93/11236; WO 95/15982; WO 95/20401; and U.S. Patent Nos. 5,698,426; 5,223,409; 5,403,484; 5,580,717; 5,427,908; 5,750,753; 5,821,047; 5,571,698; 5,427,908; 5,516,637; 5,780,225; 5,658,727; 5,733,743 and 5,969,108; each of which is incorporated herein by reference in its entirety.

As described in the above references, after phage selection, the antibody coding regions from the phage can be isolated and used to generate whole antibodies, including human antibodies, or any other desired antigen binding fragment, and expressed in any desired host, including mammalian cells, insect cells, plant cells, yeast, and bacteria, e.g., as described in detail below. For example, techniques to recombinantly produce Fab, Fab' and F(ab')₂ fragments can also be employed using methods known in the art such as those disclosed in PCT publication WO 92/22324; Mullinax et al., BioTechniques 12(6):864-869 (1992); and Sawai et al., AJRI 34:26-34 (1995); and Better et al., Science 240:1041-1043 (1988) (said references incorporated by reference in their entireties).

Examples of techniques which can be used to produce single-chain Fvs and antibodies include those described in U.S. Patents 4,946,778 and 5,258,498; Huston et al., Methods in Enzymology 203:46-88 (1991); Shu et al., PNAS 90:7995-7999 (1993); and Skerra et al., Science 240:1038-1040 (1988). For some uses, including in vivo use of antibodies in humans and in vitro detection assays, it may be preferable to use chimeric, humanized, or human antibodies. A chimeric antibody is a molecule in which different portions of the antibody are derived from different animal species, such as antibodies having a variable region derived from a murine monoclonal

antibody and a human immunoglobulin constant region. Methods for producing chimeric antibodies are known in the art. See e.g., Morrison, Science 229:1202 (1985); Oi et al., BioTechniques 4:214 (1986); Gillies et al., (1989) J. Immunol. Methods 125:191-202; U.S. Patent Nos. 5,807,715; 4,816,567; and 4,816,397, which
5 are incorporated herein by reference in their entirety. Humanized antibodies are antibody molecules from non-human species antibody that binds the desired antigen having one or more complementarity determining regions (CDRs) from the non-human species and a framework regions from a human immunoglobulin molecule. Often, framework residues in the human framework regions will be substituted with
10 the corresponding residue from the CDR donor antibody to alter, preferably improve, antigen binding. These framework substitutions are identified by methods well known in the art, e.g., by modeling of the interactions of the CDR and framework residues to identify framework residues important for antigen binding and sequence comparison to identify unusual framework residues at particular positions. (See, e.g.,
15 Queen et al., U.S. Patent No. 5,585,089; Riechmann et al., Nature 332:323 (1988), which are incorporated herein by reference in their entirety.) Antibodies can be humanized using a variety of techniques known in the art including, for example, CDR-grafting (EP 239,400; PCT publication WO 91/09967; U.S. Patent Nos. 5,225,539; 5,530,101; and 5,585,089), veneering or resurfacing (EP 592,106; EP
20 519,596; Padlan, Molecular Immunology 28(4/5):489-498 (1991); Studnicka et al., Protein Engineering 7(6):805-814 (1994); Roguska. et al., PNAS 91:969-973 (1994)), and chain shuffling (U.S. Patent No. 5,565,332).

Completely human antibodies are particularly desirable for therapeutic treatment of human patients. Human antibodies can be made by a variety of methods
25 known in the art including phage display methods described above using antibody libraries derived from human immunoglobulin sequences. See also, U.S. Patent Nos. 4,444,887 and 4,716,111; and PCT publications WO 98/46645, WO 98/50433, WO 98/24893, WO 98/16654, WO 96/34096, WO 96/33735, and WO 91/10741; each of which is incorporated herein by reference in its entirety.

30 Human antibodies can also be produced using transgenic mice which are incapable of expressing functional endogenous immunoglobulins, but which can

express human immunoglobulin genes. For example, the human heavy and light chain immunoglobulin gene complexes may be introduced randomly or by homologous recombination into mouse embryonic stem cells. Alternatively, the human variable region, constant region, and diversity region may be introduced into mouse embryonic stem cells in addition to the human heavy and light chain genes. The mouse heavy and light chain immunoglobulin genes may be rendered non-functional separately or simultaneously with the introduction of human immunoglobulin loci by homologous recombination. In particular, homozygous deletion of the JH region prevents endogenous antibody production. The modified embryonic stem cells are expanded and microinjected into blastocysts to produce chimeric mice. The chimeric mice are then bred to produce homozygous offspring which express human antibodies. The transgenic mice are immunized in the normal fashion with a selected antigen, e.g., all or a portion of a polypeptide of the invention. Monoclonal antibodies directed against the antigen can be obtained from the immunized, transgenic mice using conventional hybridoma technology. The human immunoglobulin transgenes harbored by the transgenic mice rearrange during B cell differentiation, and subsequently undergo class switching and somatic mutation. Thus, using such a technique, it is possible to produce therapeutically useful IgG, IgA, IgM and IgE antibodies. For an overview of this technology for producing human antibodies, see Lonberg and Huszar, *Int. Rev. Immunol.* 13:65-93 (1995). For a detailed discussion of this technology for producing human antibodies and human monoclonal antibodies and protocols for producing such antibodies, see, e.g., PCT publications WO 98/24893; WO 92/01047; WO 96/34096; WO 96/33735; European Patent No. 0 598 877; U.S. Patent Nos. 5,413,923; 5,625,126; 5,633,425; 5,569,825; 5,661,016; 5,545,806; 5,814,318; 5,885,793; 5,916,771; and 5,939,598, which are incorporated by reference herein in their entirety. In addition, companies such as Abgenix, Inc. (Freemont, CA) and Genpharm (San Jose, CA) can be engaged to provide human antibodies directed against a selected antigen using technology similar to that described above.

Completely human antibodies which recognize a selected epitope can be generated using a technique referred to as "guided selection." In this approach a

selected non-human monoclonal antibody, e.g., a mouse antibody, is used to guide the selection of a completely human antibody recognizing the same epitope. (Jespers et al., Bio/technology 12:899-903 (1988)).

Further, antibodies to the polypeptides of the invention can, in turn, be utilized to generate anti-idiotypic antibodies that "mimic" polypeptides of the invention using techniques well known to those skilled in the art. (See, e.g., Greenspan & Bona, FASEB J. 7(5):437-444; (1989) and Nissinoff, J. Immunol. 147(8):2429-2438 (1991)). For example, antibodies which bind to and competitively inhibit polypeptide multimerization and/or binding of a polypeptide of the invention to a ligand can be used to generate anti-idiotypes that "mimic" the polypeptide multimerization and/or binding domain and, as a consequence, bind to and neutralize polypeptide and/or its ligand. Such neutralizing anti-idiotypes or Fab fragments of such anti-idiotypes can be used in therapeutic regimens to neutralize polypeptide ligand. For example, such anti-idiotypic antibodies can be used to bind a polypeptide of the invention and/or to bind its ligands/receptors, and thereby block its biological activity.

Polynucleotides Encoding Antibodies

The invention further provides polynucleotides comprising a nucleotide sequence encoding an antibody of the invention and fragments thereof. The invention also encompasses polynucleotides that hybridize under stringent or lower stringency hybridization conditions, e.g., as defined supra, to polynucleotides that encode an antibody, preferably, that specifically binds to a polypeptide of the invention, preferably, an antibody that binds to a polypeptide having the amino acid sequence of SEQ ID NO:Y.

The polynucleotides may be obtained, and the nucleotide sequence of the polynucleotides determined, by any method known in the art. For example, if the nucleotide sequence of the antibody is known, a polynucleotide encoding the antibody may be assembled from chemically synthesized oligonucleotides (e.g., as described in Kutmeier et al., BioTechniques 17:242 (1994)), which, briefly, involves the synthesis of overlapping oligonucleotides containing portions of the sequence

encoding the antibody, annealing and ligating of those oligonucleotides, and then amplification of the ligated oligonucleotides by PCR.

Alternatively, a polynucleotide encoding an antibody may be generated from nucleic acid from a suitable source. If a clone containing a nucleic acid encoding a particular antibody is not available, but the sequence of the antibody molecule is known, a nucleic acid encoding the immunoglobulin may be chemically synthesized or obtained from a suitable source (e.g., an antibody cDNA library, or a cDNA library generated from, or nucleic acid, preferably poly A+ RNA, isolated from, any tissue or cells expressing the antibody, such as hybridoma cells selected to express an antibody of the invention) by PCR amplification using synthetic primers hybridizable to the 3' and 5' ends of the sequence or by cloning using an oligonucleotide probe specific for the particular gene sequence to identify, e.g., a cDNA clone from a cDNA library that encodes the antibody. Amplified nucleic acids generated by PCR may then be cloned into replicable cloning vectors using any method well known in the art.

Once the nucleotide sequence and corresponding amino acid sequence of the antibody is determined, the nucleotide sequence of the antibody may be manipulated using methods well known in the art for the manipulation of nucleotide sequences, e.g., recombinant DNA techniques, site directed mutagenesis, PCR, etc. (see, for example, the techniques described in Sambrook et al., 1990, Molecular Cloning, A Laboratory Manual, 2d Ed., Cold Spring Harbor Laboratory, Cold Spring Harbor, NY and Ausubel et al., eds., 1998, Current Protocols in Molecular Biology, John Wiley & Sons, NY, which are both incorporated by reference herein in their entireties), to generate antibodies having a different amino acid sequence, for example to create amino acid substitutions, deletions, and/or insertions.

In a specific embodiment, the amino acid sequence of the heavy and/or light chain variable domains may be inspected to identify the sequences of the complementarity determining regions (CDRs) by methods that are well known in the art, e.g., by comparison to known amino acid sequences of other heavy and light chain variable regions to determine the regions of sequence hypervariability. Using routine recombinant DNA techniques, one or more of the CDRs may be inserted

within framework regions, e.g., into human framework regions to humanize a non-human antibody, as described supra. The framework regions may be naturally occurring or consensus framework regions, and preferably human framework regions (see, e.g., Chothia et al., *J. Mol. Biol.* 278: 457-479 (1998) for a listing of human framework regions). Preferably, the polynucleotide generated by the combination of the framework regions and CDRs encodes an antibody that specifically binds a polypeptide of the invention. Preferably, as discussed supra, one or more amino acid substitutions may be made within the framework regions, and, preferably, the amino acid substitutions improve binding of the antibody to its antigen. Additionally, such methods may be used to make amino acid substitutions or deletions of one or more variable region cysteine residues participating in an intrachain disulfide bond to generate antibody molecules lacking one or more intrachain disulfide bonds. Other alterations to the polynucleotide are encompassed by the present invention and within the skill of the art.

In addition, techniques developed for the production of "chimeric antibodies" (Morrison et al., *Proc. Natl. Acad. Sci.* 81:851-855 (1984); Neuberger et al., *Nature* 312:604-608 (1984); Takeda et al., *Nature* 314:452-454 (1985)) by splicing genes from a mouse antibody molecule of appropriate antigen specificity together with genes from a human antibody molecule of appropriate biological activity can be used. As described supra, a chimeric antibody is a molecule in which different portions are derived from different animal species, such as those having a variable region derived from a murine mAb and a human immunoglobulin constant region, e.g., humanized antibodies.

Alternatively, techniques described for the production of single chain antibodies (U.S. Patent No. 4,946,778; Bird, *Science* 242:423-42 (1988); Huston et al., *Proc. Natl. Acad. Sci. USA* 85:5879-5883 (1988); and Ward et al., *Nature* 334:544-54 (1989)) can be adapted to produce single chain antibodies. Single chain antibodies are formed by linking the heavy and light chain fragments of the Fv region via an amino acid bridge, resulting in a single chain polypeptide. Techniques for the assembly of functional Fv fragments in *E. coli* may also be used (Skerra et al., *Science* 242:1038-1041 (1988)).

Methods of Producing Antibodies

The antibodies of the invention can be produced by any method known in the art for the synthesis of antibodies, in particular, by chemical synthesis or preferably, by recombinant expression techniques.

Recombinant expression of an antibody of the invention, or fragment, derivative or analog thereof, (e.g., a heavy or light chain of an antibody of the invention or a single chain antibody of the invention), requires construction of an expression vector containing a polynucleotide that encodes the antibody. Once a polynucleotide encoding an antibody molecule or a heavy or light chain of an antibody, or portion thereof (preferably containing the heavy or light chain variable domain), of the invention has been obtained, the vector for the production of the antibody molecule may be produced by recombinant DNA technology using techniques well known in the art. Thus, methods for preparing a protein by expressing a polynucleotide containing an antibody encoding nucleotide sequence are described herein. Methods which are well known to those skilled in the art can be used to construct expression vectors containing antibody coding sequences and appropriate transcriptional and translational control signals. These methods include, for example, in vitro recombinant DNA techniques, synthetic techniques, and in vivo genetic recombination. The invention, thus, provides replicable vectors comprising a nucleotide sequence encoding an antibody molecule of the invention, or a heavy or light chain thereof, or a heavy or light chain variable domain, operably linked to a promoter. Such vectors may include the nucleotide sequence encoding the constant region of the antibody molecule (see, e.g., PCT Publication WO 86/05807; PCT Publication WO 89/01036; and U.S. Patent No. 5,122,464) and the variable domain of the antibody may be cloned into such a vector for expression of the entire heavy or light chain.

The expression vector is transferred to a host cell by conventional techniques and the transfected cells are then cultured by conventional techniques to produce an antibody of the invention. Thus, the invention includes host cells containing a polynucleotide encoding an antibody of the invention, or a heavy or light chain

thereof, or a single chain antibody of the invention, operably linked to a heterologous promoter. In preferred embodiments for the expression of double-chained antibodies, vectors encoding both the heavy and light chains may be co-expressed in the host cell for expression of the entire immunoglobulin molecule, as detailed below.

5 A variety of host-expression vector systems may be utilized to express the antibody molecules of the invention. Such host-expression systems represent vehicles by which the coding sequences of interest may be produced and subsequently purified, but also represent cells which may, when transformed or transfected with the appropriate nucleotide coding sequences, express an antibody molecule of the
10 invention in situ. These include but are not limited to microorganisms such as bacteria (e.g., *E. coli*, *B. subtilis*) transformed with recombinant bacteriophage DNA, plasmid DNA or cosmid DNA expression vectors containing antibody coding sequences; yeast (e.g., *Saccharomyces*, *Pichia*) transformed with recombinant yeast expression vectors containing antibody coding sequences; insect cell systems
15 infected with recombinant virus expression vectors (e.g., baculovirus) containing antibody coding sequences; plant cell systems infected with recombinant virus expression vectors (e.g., cauliflower mosaic virus, CaMV; tobacco mosaic virus, TMV) or transformed with recombinant plasmid expression vectors (e.g., Ti plasmid) containing antibody coding sequences; or mammalian cell systems (e.g., COS, CHO,
20 BHK, 293, 3T3 cells) harboring recombinant expression constructs containing promoters derived from the genome of mammalian cells (e.g., metallothionein promoter) or from mammalian viruses (e.g., the adenovirus late promoter; the vaccinia virus 7.5K promoter). Preferably, bacterial cells such as *Escherichia coli*, and more preferably, eukaryotic cells, especially for the expression of whole
25 recombinant antibody molecule, are used for the expression of a recombinant antibody molecule. For example, mammalian cells such as Chinese hamster ovary cells (CHO), in conjunction with a vector such as the major intermediate early gene promoter element from human cytomegalovirus is an effective expression system for antibodies (Foecking et al., *Gene* 45:101 (1986); Cockett et al., *Bio/Technology* 8:2
30 (1990)).

In bacterial systems, a number of expression vectors may be advantageously selected depending upon the use intended for the antibody molecule being expressed. For example, when a large quantity of such a protein is to be produced, for the generation of pharmaceutical compositions of an antibody molecule, vectors which
5 direct the expression of high levels of fusion protein products that are readily purified may be desirable. Such vectors include, but are not limited, to the *E. coli* expression vector pUR278 (Ruther et al., EMBO J. 2:1791 (1983)), in which the antibody coding sequence may be ligated individually into the vector in frame with the lac Z coding region so that a fusion protein is produced; pIN vectors (Inouye & Inouye, Nucleic
10 Acids Res. 13:3101-3109 (1985); Van Heeke & Schuster, J. Biol. Chem. 24:5503-5509 (1989)); and the like. pGEX vectors may also be used to express foreign polypeptides as fusion proteins with glutathione S-transferase (GST). In general, such fusion proteins are soluble and can easily be purified from lysed cells by adsorption and binding to matrix glutathione-agarose beads followed by elution in the presence
15 of free glutathione. The pGEX vectors are designed to include thrombin or factor Xa protease cleavage sites so that the cloned target gene product can be released from the GST moiety.

In an insect system, *Autographa californica* nuclear polyhedrosis virus (AcNPV) is used as a vector to express foreign genes. The virus grows in
20 *Spodoptera frugiperda* cells. The antibody coding sequence may be cloned individually into non-essential regions (for example the polyhedrin gene) of the virus and placed under control of an AcNPV promoter (for example the polyhedrin promoter).

In mammalian host cells, a number of viral-based expression systems may be
25 utilized. In cases where an adenovirus is used as an expression vector, the antibody coding sequence of interest may be ligated to an adenovirus transcription/translation control complex, e.g., the late promoter and tripartite leader sequence. This chimeric gene may then be inserted in the adenovirus genome by in vitro or in vivo recombination. Insertion in a non-essential region of the viral genome (e.g., region
30 E1 or E3) will result in a recombinant virus that is viable and capable of expressing the antibody molecule in infected hosts. (e.g., see Logan & Shenk, Proc. Natl. Acad.

Sci. USA 81:355-359 (1984)). Specific initiation signals may also be required for efficient translation of inserted antibody coding sequences. These signals include the ATG initiation codon and adjacent sequences. Furthermore, the initiation codon must be in phase with the reading frame of the desired coding sequence to ensure translation of the entire insert. These exogenous translational control signals and initiation codons can be of a variety of origins, both natural and synthetic. The efficiency of expression may be enhanced by the inclusion of appropriate transcription enhancer elements, transcription terminators, etc. (see Bittner et al., Methods in Enzymol. 153:51-544 (1987)).

10 In addition, a host cell strain may be chosen which modulates the expression of the inserted sequences, or modifies and processes the gene product in the specific fashion desired. Such modifications (e.g., glycosylation) and processing (e.g., cleavage) of protein products may be important for the function of the protein. Different host cells have characteristic and specific mechanisms for the post-
15 translational processing and modification of proteins and gene products. Appropriate cell lines or host systems can be chosen to ensure the correct modification and processing of the foreign protein expressed. To this end, eukaryotic host cells which possess the cellular machinery for proper processing of the primary transcript, glycosylation, and phosphorylation of the gene product may be used. Such
20 mammalian host cells include but are not limited to CHO, VERY, BHK, Hela, COS, MDCK, 293, 3T3, WI38, and in particular, breast cancer cell lines such as, for example, BT483, Hs578T, HTB2, BT20 and T47D, and normal mammary gland cell line such as, for example, CRL7030 and Hs578Bst.

For long-term, high-yield production of recombinant proteins, stable
25 expression is preferred. For example, cell lines which stably express the antibody molecule may be engineered. Rather than using expression vectors which contain viral origins of replication, host cells can be transformed with DNA controlled by appropriate expression control elements (e.g., promoter, enhancer, sequences, transcription terminators, polyadenylation sites, etc.), and a selectable marker.
30 Following the introduction of the foreign DNA, engineered cells may be allowed to grow for 1-2 days in an enriched media, and then are switched to a selective media.

The selectable marker in the recombinant plasmid confers resistance to the selection and allows cells to stably integrate the plasmid into their chromosomes and grow to form foci which in turn can be cloned and expanded into cell lines. This method may advantageously be used to engineer cell lines which express the antibody molecule.

- 5 Such engineered cell lines may be particularly useful in screening and evaluation of compounds that interact directly or indirectly with the antibody molecule.

A number of selection systems may be used, including but not limited to the herpes simplex virus thymidine kinase (Wigler et al., *Cell* 11:223 (1977)), hypoxanthine-guanine phosphoribosyltransferase (Szybalska & Szybalski, *Proc. Natl. Acad. Sci. USA* 48:202 (1992)), and adenine phosphoribosyltransferase (Lowy et al., *Cell* 22:817 (1980)) genes can be employed in tk-, hgp^rt- or ap^rt- cells, respectively. Also, antimetabolite resistance can be used as the basis of selection for the following genes: dhfr, which confers resistance to methotrexate (Wigler et al., *Natl. Acad. Sci. USA* 77:357 (1980); O'Hare et al., *Proc. Natl. Acad. Sci. USA* 78:1527 (1981)); gpt, which confers resistance to mycophenolic acid (Mulligan & Berg, *Proc. Natl. Acad. Sci. USA* 78:2072 (1981)); neo, which confers resistance to the aminoglycoside G-418 (*Clinical Pharmacy* 12:488-505; Wu and Wu, *Biotherapy* 3:87-95 (1991); Tolstoshev, *Ann. Rev. Pharmacol. Toxicol.* 32:573-596 (1993); Mulligan, *Science* 260:926-932 (1993); and Morgan and Anderson, *Ann. Rev. Biochem.* 62:191-217 (1993); May, 1993, *TIB TECH* 11(5):155-215); and hyg^r, which confers resistance to hygromycin (Santerre et al., *Gene* 30:147 (1984)). Methods commonly known in the art of recombinant DNA technology may be routinely applied to select the desired recombinant clone, and such methods are described, for example, in Ausubel et al. (eds.), *Current Protocols in Molecular Biology*, John Wiley & Sons, NY (1993); Kriegler, *Gene Transfer and Expression, A Laboratory Manual*, Stockton Press, NY (1990); and in Chapters 12 and 13, Dracopoli et al. (eds), *Current Protocols in Human Genetics*, John Wiley & Sons, NY (1994); Colberre-Garapin et al., *J. Mol. Biol.* 150:1 (1981), which are incorporated by reference herein in their entireties.

- 25 The expression levels of an antibody molecule can be increased by vector amplification (for a review, see Bebbington and Hentschel, *The use of vectors based on gene amplification for the expression of cloned genes in mammalian cells in DNA*
- 30

cloning, Vol.3. (Academic Press, New York, 1987)). When a marker in the vector system expressing antibody is amplifiable, increase in the level of inhibitor present in culture of host cell will increase the number of copies of the marker gene. Since the amplified region is associated with the antibody gene, production of the antibody will also increase (Crouse et al., Mol. Cell. Biol. 3:257 (1983)).

The host cell may be co-transfected with two expression vectors of the invention, the first vector encoding a heavy chain derived polypeptide and the second vector encoding a light chain derived polypeptide. The two vectors may contain identical selectable markers which enable equal expression of heavy and light chain polypeptides. Alternatively, a single vector may be used which encodes, and is capable of expressing, both heavy and light chain polypeptides. In such situations, the light chain should be placed before the heavy chain to avoid an excess of toxic free heavy chain (Proudfoot, Nature 322:52 (1986); Kohler, Proc. Natl. Acad. Sci. USA 77:2197 (1980)). The coding sequences for the heavy and light chains may comprise cDNA or genomic DNA.

Once an antibody molecule of the invention has been produced by an animal, chemically synthesized, or recombinantly expressed, it may be purified by any method known in the art for purification of an immunoglobulin molecule, for example, by chromatography (e.g., ion exchange, affinity, particularly by affinity for the specific antigen after Protein A, and sizing column chromatography), centrifugation, differential solubility, or by any other standard technique for the purification of proteins. In addition, the antibodies of the present invention or fragments thereof can be fused to heterologous polypeptide sequences described herein or otherwise known in the art, to facilitate purification.

The present invention encompasses antibodies recombinantly fused or chemically conjugated (including both covalently and non-covalently conjugations) to a polypeptide (or portion thereof, preferably at least 10, 20, 30, 40, 50, 60, 70, 80, 90 or 100 amino acids of the polypeptide) of the present invention to generate fusion proteins. The fusion does not necessarily need to be direct, but may occur through linker sequences. The antibodies may be specific for antigens other than polypeptides (or portion thereof, preferably at least 10, 20, 30, 40, 50, 60, 70, 80, 90 or 100 amino

acids of the polypeptide) of the present invention. For example, antibodies may be used to target the polypeptides of the present invention to particular cell types, either in vitro or in vivo, by fusing or conjugating the polypeptides of the present invention to antibodies specific for particular cell surface receptors. Antibodies fused or
5 conjugated to the polypeptides of the present invention may also be used in in vitro immunoassays and purification methods using methods known in the art. See e.g., Harbor et al., supra, and PCT publication WO 93/21232; EP 439,095; Naramura et al., Immunol. Lett. 39:91-99 (1994); U.S. Patent 5,474,981; Gillies et al., PNAS 89:1428-1432 (1992); Fell et al., J. Immunol. 146:2446-2452(1991), which are
10 incorporated by reference in their entireties.

The present invention further includes compositions comprising the polypeptides of the present invention fused or conjugated to antibody domains other than the variable regions. For example, the polypeptides of the present invention may be fused or conjugated to an antibody Fc region, or portion thereof. The antibody
15 portion fused to a polypeptide of the present invention may comprise the constant region, hinge region, CH1 domain, CH2 domain, and CH3 domain or any combination of whole domains or portions thereof. The polypeptides may also be fused or conjugated to the above antibody portions to form multimers. For example, Fc portions fused to the polypeptides of the present invention can form dimers
20 through disulfide bonding between the Fc portions. Higher multimeric forms can be made by fusing the polypeptides to portions of IgA and IgM. Methods for fusing or conjugating the polypeptides of the present invention to antibody portions are known in the art. See, e.g., U.S. Patent Nos. 5,336,603; 5,622,929; 5,359,046; 5,349,053; 5,447,851; 5,112,946; EP 307,434; EP 367,166; PCT publications WO 96/04388; WO
25 91/06570; Ashkenazi et al., Proc. Natl. Acad. Sci. USA 88:10535-10539 (1991); Zheng et al., J. Immunol. 154:5590-5600 (1995); and Vil et al., Proc. Natl. Acad. Sci. USA 89:11337-11341(1992) (said references incorporated by reference in their entireties).

As discussed, supra, the polypeptides corresponding to a polypeptide,
30 polypeptide fragment, or a variant of SEQ ID NO:Y may be fused or conjugated to the above antibody portions to increase the in vivo half life of the polypeptides or for

use in immunoassays using methods known in the art. Further, the polypeptides corresponding to SEQ ID NO:Y may be fused or conjugated to the above antibody portions to facilitate purification. One reported example describes chimeric proteins consisting of the first two domains of the human CD4-polypeptide and various domains of the constant regions of the heavy or light chains of mammalian immunoglobulins. (EP 394,827; Traunecker et al., *Nature* 331:84-86 (1988)). The polypeptides of the present invention fused or conjugated to an antibody having disulfide-linked dimeric structures (due to the IgG) may also be more efficient in binding and neutralizing other molecules, than the monomeric secreted protein or protein fragment alone. (Fountoulakis et al., *J. Biochem.* 270:3958-3964 (1995)). In many cases, the Fc part in a fusion protein is beneficial in therapy and diagnosis, and thus can result in, for example, improved pharmacokinetic properties. (EP A 232,262). Alternatively, deleting the Fc part after the fusion protein has been expressed, detected, and purified, would be desired. For example, the Fc portion may hinder therapy and diagnosis if the fusion protein is used as an antigen for immunizations. In drug discovery, for example, human proteins, such as hIL-5, have been fused with Fc portions for the purpose of high-throughput screening assays to identify antagonists of hIL-5. (See, Bennett et al., *J. Molecular Recognition* 8:52-58 (1995); Johanson et al., *J. Biol. Chem.* 270:9459-9471 (1995)).

Moreover, the antibodies or fragments thereof of the present invention can be fused to marker sequences, such as a peptide to facilitate purification. In preferred embodiments, the marker amino acid sequence is a hexa-histidine peptide, such as the tag provided in a pQE vector (QIAGEN, Inc., 9259 Eton Avenue, Chatsworth, CA, 91311), among others, many of which are commercially available. As described in Gentz et al., *Proc. Natl. Acad. Sci. USA* 86:821-824 (1989), for instance, hexa-histidine provides for convenient purification of the fusion protein. Other peptide tags useful for purification include, but are not limited to, the "HA" tag, which corresponds to an epitope derived from the influenza hemagglutinin protein (Wilson et al., *Cell* 37:767 (1984)) and the "flag" tag.

The present invention further encompasses antibodies or fragments thereof conjugated to a diagnostic or therapeutic agent. The antibodies can be used

diagnostically to, for example, monitor the development or progression of a tumor as part of a clinical testing procedure to, e.g., determine the efficacy of a given treatment regimen. Detection can be facilitated by coupling the antibody to a detectable substance. Examples of detectable substances include various enzymes, prosthetic groups, fluorescent materials, luminescent materials, bioluminescent materials, radioactive materials, positron emitting metals using various positron emission tomographies, and nonradioactive paramagnetic metal ions. The detectable substance may be coupled or conjugated either directly to the antibody (or fragment thereof) or indirectly, through an intermediate (such as, for example, a linker known in the art) using techniques known in the art. See, for example, U.S. Patent No. 4,741,900 for metal ions which can be conjugated to antibodies for use as diagnostics according to the present invention. Examples of suitable enzymes include horseradish peroxidase, alkaline phosphatase, beta-galactosidase, or acetylcholinesterase; examples of suitable prosthetic group complexes include streptavidin/biotin and avidin/biotin; examples of suitable fluorescent materials include umbelliferone, fluorescein, fluorescein isothiocyanate, rhodamine, dichlorotriazinylamine fluorescein, dansyl chloride or phycoerythrin; an example of a luminescent material includes luminol; examples of bioluminescent materials include luciferase, luciferin, and aequorin; and examples of suitable radioactive material include ^{125}I , ^{131}I , ^{111}In or ^{99}Tc .

Further, an antibody or fragment thereof may be conjugated to a therapeutic moiety such as a cytotoxin, e.g., a cytostatic or cytocidal agent, a therapeutic agent or a radioactive metal ion, e.g., alpha-emitters such as, for example, ^{213}Bi . A cytotoxin or cytotoxic agent includes any agent that is detrimental to cells. Examples include paclitaxol, cytochalasin B, gramicidin D, ethidium bromide, emetine, mitomycin, etoposide, tenoposide, vincristine, vinblastine, colchicin, doxorubicin, daunorubicin, dihydroxy anthracin dione, mitoxantrone, mithramycin, actinomycin D, 1-dehydrotestosterone, glucocorticoids, procaine, tetracaine, lidocaine, propranolol, and puromycin and analogs or homologs thereof. Therapeutic agents include, but are not limited to, antimetabolites (e.g., methotrexate, 6-mercaptopurine, 6-thioguanine, cytarabine, 5-fluorouracil decarbazine), alkylating agents (e.g., mechlorethamine,

thioepa chlorambucil, melphalan, carmustine (BSNU) and lomustine (CCNU), cyclophosphamide, busulfan, dibromomannitol, streptozotocin, mitomycin C, and cis-dichlorodiamine platinum (II) (DDP) cisplatin), anthracyclines (e.g., daunorubicin (formerly daunomycin) and doxorubicin), antibiotics (e.g., dactinomycin (formerly actinomycin), bleomycin, mithramycin, and anthramycin (AMC)), and anti-mitotic agents (e.g., vincristine and vinblastine).

The conjugates of the invention can be used for modifying a given biological response, the therapeutic agent or drug moiety is not to be construed as limited to classical chemical therapeutic agents. For example, the drug moiety may be a protein or polypeptide possessing a desired biological activity. Such proteins may include, for example, a toxin such as abrin, ricin A, pseudomonas exotoxin, or diphtheria toxin; a protein such as tumor necrosis factor, α -interferon, β -interferon, nerve growth factor, platelet derived growth factor, tissue plasminogen activator, an apoptotic agent, e.g., TNF- α , TNF- β , AIM I (See, International Publication No. WO 97/33899), AIM II (See, International Publication No. WO 97/34911), Fas Ligand (Takahashi *et al.*, *Int. Immunol.*, 6:1567-1574 (1994)), VEGF (See, International Publication No. WO 99/23105), a thrombotic agent or an anti-angiogenic agent, e.g., angiostatin or endostatin; or, biological response modifiers such as, for example, lymphokines, interleukin-1 ("IL-1"), interleukin-2 ("IL-2"), interleukin-6 ("IL-6"), granulocyte macrophage colony stimulating factor ("GM-CSF"), granulocyte colony stimulating factor ("G-CSF"), or other growth factors.

Antibodies may also be attached to solid supports, which are particularly useful for immunoassays or purification of the target antigen. Such solid supports include, but are not limited to, glass, cellulose, polyacrylamide, nylon, polystyrene, polyvinyl chloride or polypropylene.

Techniques for conjugating such therapeutic moiety to antibodies are well known, see, e.g., Arnon *et al.*, "Monoclonal Antibodies For Immunotargeting Of Drugs In Cancer Therapy", in *Monoclonal Antibodies And Cancer Therapy*, Reisfeld *et al.* (eds.), pp. 243-56 (Alan R. Liss, Inc. 1985); Hellstrom *et al.*, "Antibodies For Drug Delivery", in *Controlled Drug Delivery* (2nd Ed.), Robinson *et al.* (eds.), pp. 623-53 (Marcel Dekker, Inc. 1987); Thorpe, "Antibody Carriers Of Cytotoxic Agents

In Cancer Therapy: A Review", in *Monoclonal Antibodies '84: Biological And Clinical Applications*, Pinchera et al. (eds.), pp. 475-506 (1985); "Analysis, Results, And Future Prospective Of The Therapeutic Use Of Radiolabeled Antibody In Cancer Therapy", in *Monoclonal Antibodies For Cancer Detection And Therapy*, Baldwin et al. (eds.), pp. 303-16 (Academic Press 1985), and Thorpe et al., "The Preparation And Cytotoxic Properties Of Antibody-Toxin Conjugates", *Immunol. Rev.* 62:119-58 (1982).

Alternatively, an antibody can be conjugated to a second antibody to form an antibody heteroconjugate as described by Segal in U.S. Patent No. 4,676,980, which is incorporated herein by reference in its entirety.

An antibody, with or without a therapeutic moiety conjugated to it, administered alone or in combination with cytotoxic factor(s) and/or cytokine(s) can be used as a therapeutic.

Immunophenotyping

The antibodies of the invention may be utilized for immunophenotyping of cell lines and biological samples. The translation product of the gene of the present invention may be useful as a cell specific marker, or more specifically as a cellular marker that is differentially expressed at various stages of differentiation and/or maturation of particular cell types. Monoclonal antibodies directed against a specific epitope, or combination of epitopes, will allow for the screening of cellular populations expressing the marker. Various techniques can be utilized using monoclonal antibodies to screen for cellular populations expressing the marker(s), and include magnetic separation using antibody-coated magnetic beads, "panning" with antibody attached to a solid matrix (i.e., plate), and flow cytometry (See, e.g., U.S. Patent 5,985,660; and Morrison *et al.*, *Cell*, 96:737-49 (1999)).

These techniques allow for the screening of particular populations of cells, such as might be found with hematological malignancies (i.e. minimal residual disease (MRD) in acute leukemic patients) and "non-self" cells in transplantations to prevent Graft-versus-Host Disease (GVHD). Alternatively, these techniques allow for

the screening of hematopoietic stem and progenitor cells capable of undergoing proliferation and/or differentiation, as might be found in human umbilical cord blood.

Assays For Antibody Binding

5 The antibodies of the invention may be assayed for immunospecific binding by any method known in the art. The immunoassays which can be used include but are not limited to competitive and non-competitive assay systems using techniques such as western blots, radioimmunoassays, ELISA (enzyme linked immunosorbent assay), "sandwich" immunoassays, immunoprecipitation assays, precipitin reactions,
10 gel diffusion precipitin reactions, immunodiffusion assays, agglutination assays, complement-fixation assays, immunoradiometric assays, fluorescent immunoassays, protein A immunoassays, to name but a few. Such assays are routine and well known in the art (see, e.g., Ausubel et al, eds, 1994, Current Protocols in Molecular Biology, Vol. 1, John Wiley & Sons, Inc., New York, which is incorporated by
15 reference herein in its entirety). Exemplary immunoassays are described briefly below (but are not intended by way of limitation).

Immunoprecipitation protocols generally comprise lysing a population of cells in a lysis buffer such as RIPA buffer (1% NP-40 or Triton X- 100, 1% sodium deoxycholate, 0.1% SDS, 0.15 M NaCl, 0.01 M sodium phosphate at pH 7.2, 1%
20 Trasyllol) supplemented with protein phosphatase and/or protease inhibitors (e.g., EDTA, PMSF, aprotinin, sodium vanadate), adding the antibody of interest to the cell lysate, incubating for a period of time (e.g., 1-4 hours) at 4° C, adding protein A and/or protein G sepharose beads to the cell lysate, incubating for about an hour or more at 4° C, washing the beads in lysis buffer and resuspending the beads in
25 SDS/sample buffer. The ability of the antibody of interest to immunoprecipitate a particular antigen can be assessed by, e.g., western blot analysis. One of skill in the art would be knowledgeable as to the parameters that can be modified to increase the binding of the antibody to an antigen and decrease the background (e.g., pre-clearing the cell lysate with sepharose beads). For further discussion regarding
30 immunoprecipitation protocols see, e.g., Ausubel et al, eds, 1994, Current Protocols in Molecular Biology, Vol. 1, John Wiley & Sons, Inc., New York at 10.16.1.

Western blot analysis generally comprises preparing protein samples, electrophoresis of the protein samples in a polyacrylamide gel (e.g., 8%- 20% SDS-PAGE depending on the molecular weight of the antigen), transferring the protein sample from the polyacrylamide gel to a membrane such as nitrocellulose, PVDF or nylon, blocking the membrane in blocking solution (e.g., PBS with 3% BSA or non-fat milk), washing the membrane in washing buffer (e.g., PBS-Tween 20), blocking the membrane with primary antibody (the antibody of interest) diluted in blocking buffer, washing the membrane in washing buffer, blocking the membrane with a secondary antibody (which recognizes the primary antibody, e.g., an anti-human antibody) conjugated to an enzymatic substrate (e.g., horseradish peroxidase or alkaline phosphatase) or radioactive molecule (e.g., ^{32}P or ^{125}I) diluted in blocking buffer, washing the membrane in wash buffer, and detecting the presence of the antigen. One of skill in the art would be knowledgeable as to the parameters that can be modified to increase the signal detected and to reduce the background noise. For further discussion regarding western blot protocols see, e.g., Ausubel et al, eds, 1994, Current Protocols in Molecular Biology, Vol. 1, John Wiley & Sons, Inc., New York at 10.8.1.

ELISAs comprise preparing antigen, coating the well of a 96 well microtiter plate with the antigen, adding the antibody of interest conjugated to a detectable compound such as an enzymatic substrate (e.g., horseradish peroxidase or alkaline phosphatase) to the well and incubating for a period of time, and detecting the presence of the antigen. In ELISAs the antibody of interest does not have to be conjugated to a detectable compound; instead, a second antibody (which recognizes the antibody of interest) conjugated to a detectable compound may be added to the well. Further, instead of coating the well with the antigen, the antibody may be coated to the well. In this case, a second antibody conjugated to a detectable compound may be added following the addition of the antigen of interest to the coated well. One of skill in the art would be knowledgeable as to the parameters that can be modified to increase the signal detected as well as other variations of ELISAs known in the art. For further discussion regarding ELISAs see, e.g., Ausubel et al,

eds, 1994, Current Protocols in Molecular Biology, Vol. 1, John Wiley & Sons, Inc., New York at 11.2.1.

The binding affinity of an antibody to an antigen and the off-rate of an antibody-antigen interaction can be determined by competitive binding assays. One
5 example of a competitive binding assay is a radioimmunoassay comprising the incubation of labeled antigen (e.g., ^3H or ^{125}I) with the antibody of interest in the presence of increasing amounts of unlabeled antigen, and the detection of the antibody bound to the labeled antigen. The affinity of the antibody of interest for a particular antigen and the binding off-rates can be determined from the data by
10 scatchard plot analysis. Competition with a second antibody can also be determined using radioimmunoassays. In this case, the antigen is incubated with antibody of interest conjugated to a labeled compound (e.g., ^3H or ^{125}I) in the presence of increasing amounts of an unlabeled second antibody.

15 Therapeutic Uses

The present invention is further directed to antibody-based therapies which involve administering antibodies of the invention to an animal, preferably a mammal, and most preferably a human, patient for treating one or more of the disclosed diseases, disorders, or conditions. Therapeutic compounds of the invention include,
20 but are not limited to, antibodies of the invention (including fragments, analogs and derivatives thereof as described herein) and nucleic acids encoding antibodies of the invention (including fragments, analogs and derivatives thereof and anti-idiotypic antibodies as described herein). The antibodies of the invention can be used to treat, inhibit or prevent diseases, disorders or conditions associated with aberrant expression
25 and/or activity of a polypeptide of the invention, including, but not limited to, any one or more of the diseases, disorders, or conditions described herein. The treatment and/or prevention of diseases, disorders, or conditions associated with aberrant expression and/or activity of a polypeptide of the invention includes, but is not limited to, alleviating symptoms associated with those diseases, disorders or
30 conditions. Antibodies of the invention may be provided in pharmaceutically acceptable compositions as known in the art or as described herein.

A summary of the ways in which the antibodies of the present invention may be used therapeutically includes binding polynucleotides or polypeptides of the present invention locally or systemically in the body or by direct cytotoxicity of the antibody, e.g. as mediated by complement (CDC) or by effector cells (ADCC).

- 5 Some of these approaches are described in more detail below. Armed with the teachings provided herein, one of ordinary skill in the art will know how to use the antibodies of the present invention for diagnostic, monitoring or therapeutic purposes without undue experimentation.

- 10 The antibodies of this invention may be advantageously utilized in combination with other monoclonal or chimeric antibodies, or with lymphokines or hematopoietic growth factors (such as, e.g., IL-2, IL-3 and IL-7), for example, which serve to increase the number or activity of effector cells which interact with the antibodies.

- 15 The antibodies of the invention may be administered alone or in combination with other types of treatments (e.g., radiation therapy, chemotherapy, hormonal therapy, immunotherapy and anti-tumor agents). Generally, administration of products of a species origin or species reactivity (in the case of antibodies) that is the same species as that of the patient is preferred. Thus, in a preferred embodiment, human antibodies, fragments derivatives, analogs, or nucleic acids, are administered
20 to a human patient for therapy or prophylaxis.

- It is preferred to use high affinity and/or potent in vivo inhibiting and/or neutralizing antibodies against polypeptides or polynucleotides of the present invention, fragments or regions thereof, for both immunoassays directed to and therapy of disorders related to polynucleotides or polypeptides, including fragments
25 thereof, of the present invention. Such antibodies, fragments, or regions, will preferably have an affinity for polynucleotides or polypeptides of the invention, including fragments thereof. Preferred binding affinities include those with a dissociation constant or K_d less than 5×10^{-2} M, 10^{-2} M, 5×10^{-3} M, 10^{-3} M, 5×10^{-4} M, 10^{-4} M, 5×10^{-5} M, 10^{-5} M, 5×10^{-6} M, 10^{-6} M, 5×10^{-7} M, 10^{-7} M, 5×10^{-8} M,
30 10^{-8} M, 5×10^{-9} M, 10^{-9} M, 5×10^{-10} M, 10^{-10} M, 5×10^{-11} M, 10^{-11} M, 5×10^{-12} M, 10^{-12} M, 5×10^{-13} M, 10^{-13} M, 5×10^{-14} M, 10^{-14} M, 5×10^{-15} M, and 10^{-15} M.

Gene Therapy

In a specific embodiment, nucleic acids comprising sequences encoding antibodies or functional derivatives thereof, are administered to treat, inhibit or prevent a disease or disorder associated with aberrant expression and/or activity of a polypeptide of the invention, by way of gene therapy. Gene therapy refers to therapy performed by the administration to a subject of an expressed or expressible nucleic acid. In this embodiment of the invention, the nucleic acids produce their encoded protein that mediates a therapeutic effect.

Any of the methods for gene therapy available in the art can be used according to the present invention. Exemplary methods are described below.

For general reviews of the methods of gene therapy, see Goldspiel et al., *Clinical Pharmacy* 12:488-505 (1993); Wu and Wu, *Biotherapy* 3:87-95 (1991); Tolstoshev, *Ann. Rev. Pharmacol. Toxicol.* 32:573-596 (1993); Mulligan, *Science* 260:926-932 (1993); and Morgan and Anderson, *Ann. Rev. Biochem.* 62:191-217 (1993); May, *TIBTECH* 11(5):155-215 (1993). Methods commonly known in the art of recombinant DNA technology which can be used are described in Ausubel et al. (eds.), *Current Protocols in Molecular Biology*, John Wiley & Sons, NY (1993); and Kriegler, *Gene Transfer and Expression, A Laboratory Manual*, Stockton Press, NY (1990).

In a preferred aspect, the compound comprises nucleic acid sequences encoding an antibody, said nucleic acid sequences being part of expression vectors that express the antibody or fragments or chimeric proteins or heavy or light chains thereof in a suitable host. In particular, such nucleic acid sequences have promoters operably linked to the antibody coding region, said promoter being inducible or constitutive, and, optionally, tissue-specific. In another particular embodiment, nucleic acid molecules are used in which the antibody coding sequences and any other desired sequences are flanked by regions that promote homologous recombination at a desired site in the genome, thus providing for intrachromosomal expression of the antibody encoding nucleic acids (Koller and Smithies, *Proc. Natl. Acad. Sci. USA* 86:8932-8935 (1989); Zijlstra et al., *Nature* 342:435-438 (1989). In specific

embodiments, the expressed antibody molecule is a single chain antibody; alternatively, the nucleic acid sequences include sequences encoding both the heavy and light chains, or fragments thereof, of the antibody.

Delivery of the nucleic acids into a patient may be either direct, in which case the patient is directly exposed to the nucleic acid or nucleic acid- carrying vectors, or indirect, in which case, cells are first transformed with the nucleic acids in vitro, then transplanted into the patient. These two approaches are known, respectively, as in vivo or ex vivo gene therapy.

In a specific embodiment, the nucleic acid sequences are directly administered in vivo, where it is expressed to produce the encoded product. This can be accomplished by any of numerous methods known in the art, e.g., by constructing them as part of an appropriate nucleic acid expression vector and administering it so that they become intracellular, e.g., by infection using defective or attenuated retrovirals or other viral vectors (see U.S. Patent No. 4,980,286), or by direct injection of naked DNA, or by use of microparticle bombardment (e.g., a gene gun; Biolistic; Dupont), or coating with lipids or cell-surface receptors or transfecting agents, encapsulation in liposomes, microparticles, or microcapsules, or by administering them in linkage to a peptide which is known to enter the nucleus, by administering it in linkage to a ligand subject to receptor-mediated endocytosis (see, e.g., Wu and Wu, J. Biol. Chem. 262:4429-4432 (1987)) (which can be used to target cell types specifically expressing the receptors), etc. In another embodiment, nucleic acid-ligand complexes can be formed in which the ligand comprises a fusogenic viral peptide to disrupt endosomes, allowing the nucleic acid to avoid lysosomal degradation. In yet another embodiment, the nucleic acid can be targeted in vivo for cell specific uptake and expression, by targeting a specific receptor (see, e.g., PCT Publications WO 92/06180; WO 92/22635; WO92/20316; WO93/14188, WO 93/20221). Alternatively, the nucleic acid can be introduced intracellularly and incorporated within host cell DNA for expression, by homologous recombination (Koller and Smithies, Proc. Natl. Acad. Sci. USA 86:8932-8935 (1989); Zijlstra et al., Nature 342:435-438 (1989)).

In a specific embodiment, viral vectors that contains nucleic acid sequences encoding an antibody of the invention are used. For example, a retroviral vector can be used (see Miller et al., Meth. Enzymol. 217:581-599 (1993)). These retroviral vectors contain the components necessary for the correct packaging of the viral genome and integration into the host cell DNA. The nucleic acid sequences encoding the antibody to be used in gene therapy are cloned into one or more vectors, which facilitates delivery of the gene into a patient. More detail about retroviral vectors can be found in Boesen et al., Biotherapy 6:291-302 (1994), which describes the use of a retroviral vector to deliver the *mdr1* gene to hematopoietic stem cells in order to make the stem cells more resistant to chemotherapy. Other references illustrating the use of retroviral vectors in gene therapy are: Clowes et al., J. Clin. Invest. 93:644-651 (1994); Kiem et al., Blood 83:1467-1473 (1994); Salmons and Gunzberg, Human Gene Therapy 4:129-141 (1993); and Grossman and Wilson, Curr. Opin. in Genetics and Devel. 3:110-114 (1993).

Adenoviruses are other viral vectors that can be used in gene therapy. Adenoviruses are especially attractive vehicles for delivering genes to respiratory epithelia. Adenoviruses naturally infect respiratory epithelia where they cause a mild disease. Other targets for adenovirus-based delivery systems are liver, the central nervous system, endothelial cells, and muscle. Adenoviruses have the advantage of being capable of infecting non-dividing cells. Kozarsky and Wilson, Current Opinion in Genetics and Development 3:499-503 (1993) present a review of adenovirus-based gene therapy. Bout et al., Human Gene Therapy 5:3-10 (1994) demonstrated the use of adenovirus vectors to transfer genes to the respiratory epithelia of rhesus monkeys. Other instances of the use of adenoviruses in gene therapy can be found in Rosenfeld et al., Science 252:431-434 (1991); Rosenfeld et al., Cell 68:143-155 (1992); Mastrangeli et al., J. Clin. Invest. 91:225-234 (1993); PCT Publication WO94/12649; and Wang, et al., Gene Therapy 2:775-783 (1995). In a preferred embodiment, adenovirus vectors are used.

Adeno-associated virus (AAV) has also been proposed for use in gene therapy (Walsh et al., Proc. Soc. Exp. Biol. Med. 204:289-300 (1993); U.S. Patent No. 5,436,146).

Another approach to gene therapy involves transferring a gene to cells in tissue culture by such methods as electroporation, lipofection, calcium phosphate mediated transfection, or viral infection. Usually, the method of transfer includes the transfer of a selectable marker to the cells. The cells are then placed under selection to isolate those cells that have taken up and are expressing the transferred gene. Those cells are then delivered to a patient.

In this embodiment, the nucleic acid is introduced into a cell prior to administration in vivo of the resulting recombinant cell. Such introduction can be carried out by any method known in the art, including but not limited to transfection, electroporation, microinjection, infection with a viral or bacteriophage vector containing the nucleic acid sequences, cell fusion, chromosome-mediated gene transfer, microcell-mediated gene transfer, spheroplast fusion, etc. Numerous techniques are known in the art for the introduction of foreign genes into cells (see, e.g., Loeffler and Behr, *Meth. Enzymol.* 217:599-618 (1993); Cohen et al., *Meth. Enzymol.* 217:618-644 (1993); Cline, *Pharmac. Ther.* 29:69-92m (1985) and may be used in accordance with the present invention, provided that the necessary developmental and physiological functions of the recipient cells are not disrupted. The technique should provide for the stable transfer of the nucleic acid to the cell, so that the nucleic acid is expressible by the cell and preferably heritable and expressible by its cell progeny.

The resulting recombinant cells can be delivered to a patient by various methods known in the art. Recombinant blood cells (e.g., hematopoietic stem or progenitor cells) are preferably administered intravenously. The amount of cells envisioned for use depends on the desired effect, patient state, etc., and can be determined by one skilled in the art.

Cells into which a nucleic acid can be introduced for purposes of gene therapy encompass any desired, available cell type, and include but are not limited to epithelial cells, endothelial cells, keratinocytes, fibroblasts, muscle cells, hepatocytes; blood cells such as Tlymphocytes, Blymphocytes, monocytes, macrophages, neutrophils, eosinophils, megakaryocytes, granulocytes; various stem or progenitor

cells, in particular hematopoietic stem or progenitor cells, e.g., as obtained from bone marrow, umbilical cord blood, peripheral blood, fetal liver, etc.

In a preferred embodiment, the cell used for gene therapy is autologous to the patient.

5 In an embodiment in which recombinant cells are used in gene therapy, nucleic acid sequences encoding an antibody are introduced into the cells such that they are expressible by the cells or their progeny, and the recombinant cells are then administered in vivo for therapeutic effect. In a specific embodiment, stem or progenitor cells are used. Any stem and/or progenitor cells which can be isolated and
10 maintained in vitro can potentially be used in accordance with this embodiment of the present invention (see e.g. PCT Publication WO 94/08598; Stemple and Anderson, Cell 71:973-985 (1992); Rheinwald, Meth. Cell Bio. 21A:229 (1980); and Pittelkow and Scott, Mayo Clinic Proc. 61:771 (1986)).

In a specific embodiment, the nucleic acid to be introduced for purposes of
15 gene therapy comprises an inducible promoter operably linked to the coding region, such that expression of the nucleic acid is controllable by controlling the presence or absence of the appropriate inducer of transcription.

Demonstration of Therapeutic or Prophylactic Activity

20 The compounds or pharmaceutical compositions of the invention are preferably tested in vitro, and then in vivo for the desired therapeutic or prophylactic activity, prior to use in humans. For example, in vitro assays to demonstrate the therapeutic or prophylactic utility of a compound or pharmaceutical composition include, the effect of a compound on a cell line or a patient tissue sample. The effect
25 of the compound or composition on the cell line and/or tissue sample can be determined utilizing techniques known to those of skill in the art including, but not limited to, rosette formation assays and cell lysis assays. In accordance with the invention, in vitro assays which can be used to determine whether administration of a specific compound is indicated, include in vitro cell culture assays in which a patient
30 tissue sample is grown in culture, and exposed to or otherwise administered a compound, and the effect of such compound upon the tissue sample is observed.

Therapeutic/Prophylactic Administration and Composition

The invention provides methods of treatment, inhibition and prophylaxis by administration to a subject of an effective amount of a compound or pharmaceutical composition of the invention, preferably an antibody of the invention. In a preferred aspect, the compound is substantially purified (e.g., substantially free from substances that limit its effect or produce undesired side-effects). The subject is preferably an animal, including but not limited to animals such as cows, pigs, horses, chickens, cats, dogs, etc., and is preferably a mammal, and most preferably human.

Formulations and methods of administration that can be employed when the compound comprises a nucleic acid or an immunoglobulin are described above; additional appropriate formulations and routes of administration can be selected from among those described herein below.

Various delivery systems are known and can be used to administer a compound of the invention, e.g., encapsulation in liposomes, microparticles, microcapsules, recombinant cells capable of expressing the compound, receptor-mediated endocytosis (see, e.g., Wu and Wu, J. Biol. Chem. 262:4429-4432 (1987)), construction of a nucleic acid as part of a retroviral or other vector, etc. Methods of introduction include but are not limited to intradermal, intramuscular, intraperitoneal, intravenous, subcutaneous, intranasal, epidural, and oral routes. The compounds or compositions may be administered by any convenient route, for example by infusion or bolus injection, by absorption through epithelial or mucocutaneous linings (e.g., oral mucosa, rectal and intestinal mucosa, etc.) and may be administered together with other biologically active agents. Administration can be systemic or local. In addition, it may be desirable to introduce the pharmaceutical compounds or compositions of the invention into the central nervous system by any suitable route, including intraventricular and intrathecal injection; intraventricular injection may be facilitated by an intraventricular catheter, for example, attached to a reservoir, such as an Ommaya reservoir. Pulmonary administration can also be employed, e.g., by use of an inhaler or nebulizer, and formulation with an aerosolizing agent.

In a specific embodiment, it may be desirable to administer the pharmaceutical compounds or compositions of the invention locally to the area in need of treatment; this may be achieved by, for example, and not by way of limitation, local infusion during surgery, topical application, e.g., in conjunction with a wound dressing after surgery, by injection, by means of a catheter, by means of a suppository, or by means of an implant, said implant being of a porous, non-porous, or gelatinous material, including membranes, such as sialastic membranes, or fibers. Preferably, when administering a protein, including an antibody, of the invention, care must be taken to use materials to which the protein does not absorb.

10 In another embodiment, the compound or composition can be delivered in a vesicle, in particular a liposome (see Langer, *Science* 249:1527-1533 (1990); Treat et al., in *Liposomes in the Therapy of Infectious Disease and Cancer*, Lopez-Berestein and Fidler (eds.), Liss, New York, pp. 353- 365 (1989); Lopez-Berestein, *ibid.*, pp. 317-327; see generally *ibid.*)

15 In yet another embodiment, the compound or composition can be delivered in a controlled release system. In one embodiment, a pump may be used (see Langer, *supra*; Sefton, *CRC Crit. Ref. Biomed. Eng.* 14:201 (1987); Buchwald et al., *Surgery* 88:507 (1980); Saudek et al., *N. Engl. J. Med.* 321:574 (1989)). In another embodiment, polymeric materials can be used (see *Medical Applications of*
20 *Controlled Release*, Langer and Wise (eds.), CRC Pres., Boca Raton, Florida (1974); *Controlled Drug Bioavailability, Drug Product Design and Performance*, Smolen and Ball (eds.), Wiley, New York (1984); Ranger and Peppas, J., *Macromol. Sci. Rev. Macromol. Chem.* 23:61 (1983); see also Levy et al., *Science* 228:190 (1985); During et al., *Ann. Neurol.* 25:351 (1989); Howard et al., *J.Neurosurg.* 71:105 (1989)). In yet
25 another embodiment, a controlled release system can be placed in proximity of the therapeutic target, i.e., the brain, thus requiring only a fraction of the systemic dose (see, e.g., Goodson, in *Medical Applications of Controlled Release*, *supra*, vol. 2, pp. 115-138 (1984)).

Other controlled release systems are discussed in the review by Langer
30 (*Science* 249:1527-1533 (1990)).

In a specific embodiment where the compound of the invention is a nucleic acid encoding a protein, the nucleic acid can be administered in vivo to promote expression of its encoded protein, by constructing it as part of an appropriate nucleic acid expression vector and administering it so that it becomes intracellular, e.g., by use of a retroviral vector (see U.S. Patent No. 4,980,286), or by direct injection, or by use of microparticle bombardment (e.g., a gene gun; Biolistic, Dupont), or coating with lipids or cell-surface receptors or transfecting agents, or by administering it in linkage to a homeobox-like peptide which is known to enter the nucleus (see e.g., Joliet et al., Proc. Natl. Acad. Sci. USA 88:1864-1868 (1991)), etc. Alternatively, a nucleic acid can be introduced intracellularly and incorporated within host cell DNA for expression, by homologous recombination.

The present invention also provides pharmaceutical compositions. Such compositions comprise a therapeutically effective amount of a compound, and a pharmaceutically acceptable carrier. In a specific embodiment, the term "pharmaceutically acceptable" means approved by a regulatory agency of the Federal or a state government or listed in the U.S. Pharmacopeia or other generally recognized pharmacopeia for use in animals, and more particularly in humans. The term "carrier" refers to a diluent, adjuvant, excipient, or vehicle with which the therapeutic is administered. Such pharmaceutical carriers can be sterile liquids, such as water and oils, including those of petroleum, animal, vegetable or synthetic origin, such as peanut oil, soybean oil, mineral oil, sesame oil and the like. Water is a preferred carrier when the pharmaceutical composition is administered intravenously. Saline solutions and aqueous dextrose and glycerol solutions can also be employed as liquid carriers, particularly for injectable solutions. Suitable pharmaceutical excipients include starch, glucose, lactose, sucrose, gelatin, malt, rice, flour, chalk, silica gel, sodium stearate, glycerol monostearate, talc, sodium chloride, dried skim milk, glycerol, propylene, glycol, water, ethanol and the like. The composition, if desired, can also contain minor amounts of wetting or emulsifying agents, or pH buffering agents. These compositions can take the form of solutions, suspensions, emulsion, tablets, pills, capsules, powders, sustained-release formulations and the like. The composition can be formulated as a suppository, with traditional binders and carriers

such as triglycerides. Oral formulation can include standard carriers such as pharmaceutical grades of mannitol, lactose, starch, magnesium stearate, sodium saccharine, cellulose, magnesium carbonate, etc. Examples of suitable pharmaceutical carriers are described in "Remington's Pharmaceutical Sciences" by E.W. Martin. Such compositions will contain a therapeutically effective amount of the compound, preferably in purified form, together with a suitable amount of carrier so as to provide the form for proper administration to the patient. The formulation should suit the mode of administration.

In a preferred embodiment, the composition is formulated in accordance with routine procedures as a pharmaceutical composition adapted for intravenous administration to human beings. Typically, compositions for intravenous administration are solutions in sterile isotonic aqueous buffer. Where necessary, the composition may also include a solubilizing agent and a local anesthetic such as lignocaine to ease pain at the site of the injection. Generally, the ingredients are supplied either separately or mixed together in unit dosage form, for example, as a dry lyophilized powder or water free concentrate in a hermetically sealed container such as an ampoule or sachette indicating the quantity of active agent. Where the composition is to be administered by infusion, it can be dispensed with an infusion bottle containing sterile pharmaceutical grade water or saline. Where the composition is administered by injection, an ampoule of sterile water for injection or saline can be provided so that the ingredients may be mixed prior to administration.

The compounds of the invention can be formulated as neutral or salt forms. Pharmaceutically acceptable salts include those formed with anions such as those derived from hydrochloric, phosphoric, acetic, oxalic, tartaric acids, etc., and those formed with cations such as those derived from sodium, potassium, ammonium, calcium, ferric hydroxides, isopropylamine, triethylamine, 2-ethylamino ethanol, histidine, procaine, etc.

The amount of the compound of the invention which will be effective in the treatment, inhibition and prevention of a disease or disorder associated with aberrant expression and/or activity of a polypeptide of the invention can be determined by standard clinical techniques. In addition, in vitro assays may optionally be employed

to help identify optimal dosage ranges. The precise dose to be employed in the formulation will also depend on the route of administration, and the seriousness of the disease or disorder, and should be decided according to the judgment of the practitioner and each patient's circumstances. Effective doses may be extrapolated
5 from dose-response curves derived from in vitro or animal model test systems.

For antibodies, the dosage administered to a patient is typically 0.1 mg/kg to 100 mg/kg of the patient's body weight. Preferably, the dosage administered to a patient is between 0.1 mg/kg and 20 mg/kg of the patient's body weight, more preferably 1 mg/kg to 10 mg/kg of the patient's body weight. Generally, human
10 antibodies have a longer half-life within the human body than antibodies from other species due to the immune response to the foreign polypeptides. Thus, lower dosages of human antibodies and less frequent administration is often possible. Further, the dosage and frequency of administration of antibodies of the invention may be reduced by enhancing uptake and tissue penetration (e.g., into the brain) of the
15 antibodies by modifications such as, for example, lipidation.

The invention also provides a pharmaceutical pack or kit comprising one or more containers filled with one or more of the ingredients of the pharmaceutical compositions of the invention. Optionally associated with such container(s) can be a notice in the form prescribed by a governmental agency regulating the manufacture,
20 use or sale of pharmaceuticals or biological products, which notice reflects approval by the agency of manufacture, use or sale for human administration. Diagnosis and Imaging

Labeled antibodies, and derivatives and analogs thereof, which specifically bind to a polypeptide of interest can be used for diagnostic purposes to detect,
25 diagnose, or monitor diseases and/or disorders associated with the aberrant expression and/or activity of a polypeptide of the invention. The invention provides for the detection of aberrant expression of a polypeptide of interest, comprising (a) assaying the expression of the polypeptide of interest in cells or body fluid of an individual using one or more antibodies specific to the polypeptide interest and (b) comparing
30 the level of gene expression with a standard gene expression level, whereby an

increase or decrease in the assayed polypeptide gene expression level compared to the standard expression level is indicative of aberrant expression.

The invention provides a diagnostic assay for diagnosing a disorder, comprising (a) assaying the expression of the polypeptide of interest in cells or body fluid of an individual using one or more antibodies specific to the polypeptide interest and (b) comparing the level of gene expression with a standard gene expression level, whereby an increase or decrease in the assayed polypeptide gene expression level compared to the standard expression level is indicative of a particular disorder. With respect to cancer, the presence of a relatively high amount of transcript in biopsied tissue from an individual may indicate a predisposition for the development of the disease, or may provide a means for detecting the disease prior to the appearance of actual clinical symptoms. A more definitive diagnosis of this type may allow health professionals to employ preventative measures or aggressive treatment earlier thereby preventing the development or further progression of the cancer.

Antibodies of the invention can be used to assay protein levels in a biological sample using classical immunohistological methods known to those of skill in the art (e.g., see Jalkanen, et al., J. Cell. Biol. 101:976-985 (1985); Jalkanen, et al., J. Cell Biol. 105:3087-3096 (1987)). Other antibody-based methods useful for detecting protein gene expression include immunoassays, such as the enzyme linked immunosorbent assay (ELISA) and the radioimmunoassay (RIA). Suitable antibody assay labels are known in the art and include enzyme labels, such as, glucose oxidase; radioisotopes, such as iodine (^{125}I , ^{121}I), carbon (^{14}C), sulfur (^{35}S), tritium (^3H), indium (^{112}In), and technetium (^{99}Tc); luminescent labels, such as luminol; and fluorescent labels, such as fluorescein and rhodamine, and biotin.

One aspect of the invention is the detection and diagnosis of a disease or disorder associated with aberrant expression of a polypeptide of interest in an animal, preferably a mammal and most preferably a human. In one embodiment, diagnosis comprises: a) administering (for example, parenterally, subcutaneously, or intraperitoneally) to a subject an effective amount of a labeled molecule which specifically binds to the polypeptide of interest; b) waiting for a time interval following the administering for permitting the labeled molecule to preferentially

concentrate at sites in the subject where the polypeptide is expressed (and for unbound labeled molecule to be cleared to background level); c) determining background level; and d) detecting the labeled molecule in the subject, such that detection of labeled molecule above the background level indicates that the subject has a particular disease or disorder associated with aberrant expression of the polypeptide of interest. Background level can be determined by various methods including, comparing the amount of labeled molecule detected to a standard value previously determined for a particular system.

It will be understood in the art that the size of the subject and the imaging system used will determine the quantity of imaging moiety needed to produce diagnostic images. In the case of a radioisotope moiety, for a human subject, the quantity of radioactivity injected will normally range from about 5 to 20 millicuries of ^{99m}Tc. The labeled antibody or antibody fragment will then preferentially accumulate at the location of cells which contain the specific protein. In vivo tumor imaging is described in S.W. Burchiel et al., "Immunopharmacokinetics of Radiolabeled Antibodies and Their Fragments." (Chapter 13 in Tumor Imaging: The Radiochemical Detection of Cancer, S.W. Burchiel and B. A. Rhodes, eds., Masson Publishing Inc. (1982).

Depending on several variables, including the type of label used and the mode of administration, the time interval following the administration for permitting the labeled molecule to preferentially concentrate at sites in the subject and for unbound labeled molecule to be cleared to background level is 6 to 48 hours or 6 to 24 hours or 6 to 12 hours. In another embodiment the time interval following administration is 5 to 20 days or 5 to 10 days.

In an embodiment, monitoring of the disease or disorder is carried out by repeating the method for diagnosing the disease or disease, for example, one month after initial diagnosis, six months after initial diagnosis, one year after initial diagnosis, etc.

Presence of the labeled molecule can be detected in the patient using methods known in the art for in vivo scanning. These methods depend upon the type of label used. Skilled artisans will be able to determine the appropriate method for detecting a

particular label. Methods and devices that may be used in the diagnostic methods of the invention include, but are not limited to, computed tomography (CT), whole body scan such as position emission tomography (PET), magnetic resonance imaging (MRI), and sonography.

5 In a specific embodiment, the molecule is labeled with a radioisotope and is detected in the patient using a radiation responsive surgical instrument (Thurston et al., U.S. Patent No. 5,441,050). In another embodiment, the molecule is labeled with a fluorescent compound and is detected in the patient using a fluorescence responsive scanning instrument. In another embodiment, the molecule is labeled with a positron
10 emitting metal and is detected in the patient using positron emission-tomography. In yet another embodiment, the molecule is labeled with a paramagnetic label and is detected in a patient using magnetic resonance imaging (MRI). Kits

 The present invention provides kits that can be used in the above methods. In one embodiment, a kit comprises an antibody of the invention, preferably a purified
15 antibody, in one or more containers. In a specific embodiment, the kits of the present invention contain a substantially isolated polypeptide comprising an epitope which is specifically immunoreactive with an antibody included in the kit. Preferably, the kits of the present invention further comprise a control antibody which does not react with the polypeptide of interest. In another specific embodiment, the kits of the present
20 invention contain a means for detecting the binding of an antibody to a polypeptide of interest (e.g., the antibody may be conjugated to a detectable substrate such as a fluorescent compound, an enzymatic substrate, a radioactive compound or a luminescent compound, or a second antibody which recognizes the first antibody may be conjugated to a detectable substrate).

25 In another specific embodiment of the present invention, the kit is a diagnostic kit for use in screening serum containing antibodies specific against proliferative and/or cancerous polynucleotides and polypeptides. Such a kit may include a control antibody that does not react with the polypeptide of interest. Such a kit may include a substantially isolated polypeptide antigen comprising an epitope which is specifically
30 immunoreactive with at least one anti-polypeptide antigen antibody. Further, such a kit includes means for detecting the binding of said antibody to the antigen (e.g., the

antibody may be conjugated to a fluorescent compound such as fluorescein or rhodamine which can be detected by flow cytometry). In specific embodiments, the kit may include a recombinantly produced or chemically synthesized polypeptide antigen. The polypeptide antigen of the kit may also be attached to a solid support.

5 In a more specific embodiment the detecting means of the above-described kit includes a solid support to which said polypeptide antigen is attached. Such a kit may also include a non-attached reporter-labeled anti-human antibody. In this embodiment, binding of the antibody to the polypeptide antigen can be detected by binding of the said reporter-labeled antibody.

10 In an additional embodiment, the invention includes a diagnostic kit for use in screening serum containing antigens of the polypeptide of the invention. The diagnostic kit includes a substantially isolated antibody specifically immunoreactive with polypeptide or polynucleotide antigens, and means for detecting the binding of the polynucleotide or polypeptide antigen to the antibody. In one embodiment, the
15 antibody is attached to a solid support. In a specific embodiment, the antibody may be a monoclonal antibody. The detecting means of the kit may include a second, labeled monoclonal antibody. Alternatively, or in addition, the detecting means may include a labeled, competing antigen.

In one diagnostic configuration, test serum is reacted with a solid phase
20 reagent having a surface-bound antigen obtained by the methods of the present invention. After binding with specific antigen antibody to the reagent and removing unbound serum components by washing, the reagent is reacted with reporter-labeled anti-human antibody to bind reporter to the reagent in proportion to the amount of bound anti-antigen antibody on the solid support. The reagent is again washed to
25 remove unbound labeled antibody, and the amount of reporter associated with the reagent is determined. Typically, the reporter is an enzyme which is detected by incubating the solid phase in the presence of a suitable fluorometric, luminescent or colorimetric substrate (Sigma, St. Louis, MO).

The solid surface reagent in the above assay is prepared by known techniques
30 for attaching protein material to solid support material, such as polymeric beads, dip sticks, 96-well plate or filter material. These attachment methods generally include

non-specific adsorption of the protein to the support or covalent attachment of the protein, typically through a free amine group, to a chemically reactive group on the solid support, such as an activated carboxyl, hydroxyl, or aldehyde group.

Alternatively, streptavidin coated plates can be used in conjunction with biotinylated antigen(s).

Thus, the invention provides an assay system or kit for carrying out this diagnostic method. The kit generally includes a support with surface-bound recombinant antigens, and a reporter-labeled anti-human antibody for detecting surface-bound anti-antigen antibody.

Fusion Proteins

Any polypeptide of the present invention can be used to generate fusion proteins. For example, the polypeptide of the present invention, when fused to a second protein, can be used as an antigenic tag. Antibodies raised against the polypeptide of the present invention can be used to indirectly detect the second protein by binding to the polypeptide. Moreover, because secreted proteins target cellular locations based on trafficking signals, the polypeptides of the present invention can be used as targeting molecules once fused to other proteins.

Examples of domains that can be fused to polypeptides of the present invention include not only heterologous signal sequences, but also other heterologous functional regions. The fusion does not necessarily need to be direct, but may occur through linker sequences.

Moreover, fusion proteins may also be engineered to improve characteristics of the polypeptide of the present invention. For instance, a region of additional amino acids, particularly charged amino acids, may be added to the N-terminus of the polypeptide to improve stability and persistence during purification from the host cell or subsequent handling and storage. Also, peptide moieties may be added to the polypeptide to facilitate purification. Such regions may be removed prior to final preparation of the polypeptide. The addition of peptide moieties to facilitate handling of polypeptides are familiar and routine techniques in the art.

Moreover, polypeptides of the present invention, including fragments, and specifically epitopes, can be combined with parts of the constant domain of immunoglobulins (IgA, IgE, IgG, IgM) or portions thereof (CH1, CH2, CH3, and any combination thereof, including both entire domains and portions thereof), resulting in
5 chimeric polypeptides. These fusion proteins facilitate purification and show an increased half-life in vivo. One reported example describes chimeric proteins consisting of the first two domains of the human CD4-polypeptide and various domains of the constant regions of the heavy or light chains of mammalian immunoglobulins. (EP A 394,827; Traunecker et al., Nature 331:84-86 (1988).)
10 Fusion proteins having disulfide-linked dimeric structures (due to the IgG) can also be more efficient in binding and neutralizing other molecules, than the monomeric secreted protein or protein fragment alone. (Fountoulakis et al., J. Biochem. 270:3958-3964 (1995).)

Similarly, EP-A-O 464 533 (Canadian counterpart 2045869) discloses fusion
15 proteins comprising various portions of constant region of immunoglobulin molecules together with another human protein or part thereof. In many cases, the Fc part in a fusion protein is beneficial in therapy and diagnosis, and thus can result in, for example, improved pharmacokinetic properties. (EP-A 0232 262.) Alternatively, deleting the Fc part after the fusion protein has been expressed, detected, and purified,
20 would be desired. For example, the Fc portion may hinder therapy and diagnosis if the fusion protein is used as an antigen for immunizations. In drug discovery, for example, human proteins, such as hIL-5, have been fused with Fc portions for the purpose of high-throughput screening assays to identify antagonists of hIL-5. (See, D. Bennett et al., J. Molecular Recognition 8:52-58 (1995); K. Johanson et al., J. Biol.
25 Chem. 270:9459-9471 (1995).)

Moreover, the polypeptides of the present invention can be fused to marker sequences, such as a peptide which facilitates purification of the fused polypeptide. In preferred embodiments, the marker amino acid sequence is a hexa-histidine peptide, such as the tag provided in a pQE vector (QIAGEN, Inc., 9259 Eton Avenue,
30 Chatsworth, CA, 91311), among others, many of which are commercially available. As described in Gentz et al., Proc. Natl. Acad. Sci. USA 86:821-824 (1989), for

instance, hexa-histidine provides for convenient purification of the fusion protein. Another peptide tag useful for purification, the "HA" tag, corresponds to an epitope derived from the influenza hemagglutinin protein. (Wilson et al., Cell 37:767 (1984).)

- 5 Thus, any of these above fusions can be engineered using the polynucleotides or the polypeptides of the present invention.

Vectors, Host Cells, and Protein Production

- 10 The present invention also relates to vectors containing the polynucleotide of the present invention, host cells, and the production of polypeptides by recombinant techniques. The vector may be, for example, a phage, plasmid, viral, or retroviral vector. Retroviral vectors may be replication competent or replication defective. In the latter case, viral propagation generally will occur only in complementing host cells.

- 15 The polynucleotides may be joined to a vector containing a selectable marker for propagation in a host. Generally, a plasmid vector is introduced in a precipitate, such as a calcium phosphate precipitate, or in a complex with a charged lipid. If the vector is a virus, it may be packaged in vitro using an appropriate packaging cell line and then transduced into host cells.

- 20 The polynucleotide insert should be operatively linked to an appropriate promoter, such as the phage lambda PL promoter, the E. coli lac, trp, phoA and tac promoters, the SV40 early and late promoters and promoters of retroviral LTRs, to name a few. Other suitable promoters will be known to the skilled artisan. The expression constructs will further contain sites for transcription initiation, termination, and, in the transcribed region, a ribosome binding site for translation. The coding
25 portion of the transcripts expressed by the constructs will preferably include a translation initiating codon at the beginning and a termination codon (UAA, UGA or UAG) appropriately positioned at the end of the polypeptide to be translated.

- As indicated, the expression vectors will preferably include at least one
30 selectable marker. Such markers include dihydrofolate reductase, G418 or neomycin resistance for eukaryotic cell culture and tetracycline, kanamycin or ampicillin

resistance genes for culturing in *E. coli* and other bacteria. Representative examples of appropriate hosts include, but are not limited to, bacterial cells, such as *E. coli*, *Streptomyces* and *Salmonella typhimurium* cells; fungal cells, such as yeast cells; insect cells such as *Drosophila* S2 and *Spodoptera* Sf9 cells; animal cells such as CHO, COS, 293, and Bowes melanoma cells; and plant cells. Appropriate culture mediums and conditions for the above-described host cells are known in the art.

Among vectors preferred for use in bacteria include pQE70, pQE60 and pQE-9, available from QIAGEN, Inc.; pBluescript vectors, Phagescript vectors, pNH8A, pNH16a, pNH18A, pNH46A, available from Stratagene Cloning Systems, Inc.; and ptrc99a, pKK223-3, pKK233-3, pDR540, pRIT5 available from Pharmacia Biotech, Inc. Among preferred eukaryotic vectors are pWLNEO, pSV2CAT, pOG44, pXT1 and pSG available from Stratagene; and pSVK3, pBPV, pMSG and pSVL available from Pharmacia. Other suitable vectors will be readily apparent to the skilled artisan.

Introduction of the construct into the host cell can be effected by calcium phosphate transfection, DEAE-dextran mediated transfection, cationic lipid-mediated transfection, electroporation, transduction, infection, or other methods. Such methods are described in many standard laboratory manuals, such as Davis et al., *Basic Methods In Molecular Biology* (1986). It is specifically contemplated that the polypeptides of the present invention may in fact be expressed by a host cell lacking a recombinant vector.

A polypeptide of this invention can be recovered and purified from recombinant cell cultures by well-known methods including ammonium sulfate or ethanol precipitation, acid extraction, anion or cation exchange chromatography, phosphocellulose chromatography, hydrophobic interaction chromatography, affinity chromatography, hydroxylapatite chromatography and lectin chromatography. Most preferably, high performance liquid chromatography ("HPLC") is employed for purification.

Polypeptides of the present invention, and preferably the secreted form, can also be recovered from: products purified from natural sources, including bodily fluids, tissues and cells, whether directly isolated or cultured; products of chemical synthetic procedures; and products produced by recombinant techniques from a

prokaryotic or eukaryotic host, including, for example, bacterial, yeast, higher plant, insect, and mammalian cells. Depending upon the host employed in a recombinant production procedure, the polypeptides of the present invention may be glycosylated or may be non-glycosylated. In addition, polypeptides of the invention may also
5 include an initial modified methionine residue, in some cases as a result of host-mediated processes. Thus, it is well known in the art that the N-terminal methionine encoded by the translation initiation codon generally is removed with high efficiency from any protein after translation in all eukaryotic cells. While the N-terminal methionine on most proteins also is efficiently removed in most prokaryotes, for some
10 proteins, this prokaryotic removal process is inefficient, depending on the nature of the amino acid to which the N-terminal methionine is covalently linked.

In addition to encompassing host cells containing the vector constructs discussed herein, the invention also encompasses primary, secondary, and immortalized host cells of vertebrate origin, particularly mammalian origin, that have
15 been engineered to delete or replace endogenous genetic material (e.g., coding sequence), and/or to include genetic material (e.g., heterologous polynucleotide sequences) that is operably associated with the polynucleotides of the invention, and which activates, alters, and/or amplifies endogenous polynucleotides. For example, techniques known in the art may be used to operably associate heterologous control
20 regions (e.g., promoter and/or enhancer) and endogenous polynucleotide sequences via homologous recombination, resulting in the formation of a new transcription unit (see, e.g., U.S. Patent No. 5,641,670, issued June 24, 1997; U.S. Patent No. 5,733,761, issued March 31, 1998; International Publication No. WO 96/29411, published September 26, 1996; International Publication No. WO 94/12650,
25 published August 4, 1994; Koller et al., Proc. Natl. Acad. Sci. USA 86:8932-8935 (1989); and Zijlstra et al., Nature 342:435-438 (1989), the disclosures of each of which are incorporated by reference in their entireties).

In addition, polypeptides of the invention can be chemically synthesized using techniques known in the art (e.g., see Creighton, 1983, *Proteins: Structures and*
30 *Molecular Principles*, W.H. Freeman & Co., N.Y., and Hunkapiller et al., *Nature*, 310:105-111 (1984)). For example, a polypeptide corresponding to a fragment of a

polypeptide sequence of the invention can be synthesized by use of a peptide synthesizer. Furthermore, if desired, nonclassical amino acids or chemical amino acid analogs can be introduced as a substitution or addition into the polypeptide sequence. Non-classical amino acids include, but are not limited to, to the D-isomers of the
5 common amino acids, 2,4-diaminobutyric acid, α -amino isobutyric acid, 4-aminobutyric acid, Abu, 2-amino butyric acid, γ -Abu, ϵ -Ahx, 6-amino hexanoic acid, Aib, 2-amino isobutyric acid, 3-amino propionic acid, ornithine, norleucine, norvaline, hydroxyproline, sarcosine, citrulline, homocitrulline, cysteic acid, t-butylglycine, t-butylalanine, phenylglycine, cyclohexylalanine, β -alanine, fluoro-
10 amino acids, designer amino acids such as β -methyl amino acids, Ca-methyl amino acids, Na-methyl amino acids, and amino acid analogs in general. Furthermore, the amino acid can be D (dextrorotary) or L (levorotary).

The invention encompasses polypeptides which are differentially modified during or after translation, *e.g.*, by glycosylation, acetylation, phosphorylation, amidation, derivatization by known protecting/blocking groups, proteolytic cleavage,
15 linkage to an antibody molecule or other cellular ligand, etc. Any of numerous chemical modifications may be carried out by known techniques, including but not limited, to specific chemical cleavage by cyanogen bromide, trypsin, chymotrypsin, papain, V8 protease, NaBH_4 ; acetylation, formylation, oxidation, reduction; metabolic
20 synthesis in the presence of tunicamycin; etc.

Additional post-translational modifications encompassed by the invention include, for example, *e.g.*, N-linked or O-linked carbohydrate chains, processing of N-terminal or C-terminal ends), attachment of chemical moieties to the amino acid backbone, chemical modifications of N-linked or O-linked carbohydrate chains, and
25 addition or deletion of an N-terminal methionine residue as a result of procaryotic host cell expression. The polypeptides may also be modified with a detectable label, such as an enzymatic, fluorescent, isotopic or affinity label to allow for detection and isolation of the protein.

Also provided by the invention are chemically modified derivatives of the
30 polypeptides of the invention which may provide additional advantages such as increased solubility, stability and circulating time of the polypeptide, or decreased

immunogenicity (see U.S. Patent NO: 4,179,337). The chemical moieties for derivitization may be selected from water soluble polymers such as polyethylene glycol, ethylene glycol/propylene glycol copolymers, carboxymethylcellulose, dextran, polyvinyl alcohol and the like. The polypeptides may be modified at random positions within the molecule, or at predetermined positions within the molecule and may include one, two, three or more attached chemical moieties.

The polymer may be of any molecular weight, and may be branched or unbranched. For polyethylene glycol, the preferred molecular weight is between about 1 kDa and about 100 kDa (the term "about" indicating that in preparations of polyethylene glycol, some molecules will weigh more, some less, than the stated molecular weight) for ease in handling and manufacturing. Other sizes may be used, depending on the desired therapeutic profile (e.g., the duration of sustained release desired, the effects, if any on biological activity, the ease in handling, the degree or lack of antigenicity and other known effects of the polyethylene glycol to a therapeutic protein or analog).

The polyethylene glycol molecules (or other chemical moieties) should be attached to the protein with consideration of effects on functional or antigenic domains of the protein. There are a number of attachment methods available to those skilled in the art, e.g., EP 0 401 384, herein incorporated by reference (coupling PEG to G-CSF), see also Malik et al., Exp. Hematol. 20:1028-1035 (1992) (reporting pegylation of GM-CSF using tresyl chloride). For example, polyethylene glycol may be covalently bound through amino acid residues via a reactive group, such as, a free amino or carboxyl group. Reactive groups are those to which an activated polyethylene glycol molecule may be bound. The amino acid residues having a free amino group may include lysine residues and the N-terminal amino acid residues; those having a free carboxyl group may include aspartic acid residues glutamic acid residues and the C-terminal amino acid residue. Sulfhydryl groups may also be used as a reactive group for attaching the polyethylene glycol molecules. Preferred for therapeutic purposes is attachment at an amino group, such as attachment at the N-terminus or lysine group.

One may specifically desire proteins chemically modified at the N-terminus. Using polyethylene glycol as an illustration of the present composition, one may select from a variety of polyethylene glycol molecules (by molecular weight, branching, etc.), the proportion of polyethylene glycol molecules to protein (polypeptide) molecules in the reaction mix, the type of pegylation reaction to be performed, and the method of obtaining the selected N-terminally pegylated protein. The method of obtaining the N-terminally pegylated preparation (i.e., separating this moiety from other monopegylated moieties if necessary) may be by purification of the N-terminally pegylated material from a population of pegylated protein molecules.

Selective proteins chemically modified at the N-terminus modification may be accomplished by reductive alkylation which exploits differential reactivity of different types of primary amino groups (lysine versus the N-terminal) available for derivatization in a particular protein. Under the appropriate reaction conditions, substantially selective derivatization of the protein at the N-terminus with a carbonyl group containing polymer is achieved.

The polypeptides of the invention may be in monomers or multimers (i.e., dimers, trimers, tetramers and higher multimers). Accordingly, the present invention relates to monomers and multimers of the polypeptides of the invention, their preparation, and compositions (preferably, *Therapeutics*) containing them. In specific embodiments, the polypeptides of the invention are monomers, dimers, trimers or tetramers. In additional embodiments, the multimers of the invention are at least dimers, at least trimers, or at least tetramers.

Multimers encompassed by the invention may be homomers or heteromers. As used herein, the term homomer, refers to a multimer containing only polypeptides corresponding to the amino acid sequence of SEQ ID NO:Y or encoded by the cDNA contained in a deposited clone (including fragments, variants, splice variants, and fusion proteins, corresponding to these polypeptides as described herein). These homomers may contain polypeptides having identical or different amino acid sequences. In a specific embodiment, a homomer of the invention is a multimer containing only polypeptides having an identical amino acid sequence. In another specific embodiment, a homomer of the invention is a multimer containing

polypeptides having different amino acid sequences. In specific embodiments, the multimer of the invention is a homodimer (*e.g.*, containing polypeptides having identical or different amino acid sequences) or a homotrimer (*e.g.*, containing polypeptides having identical and/or different amino acid sequences). In additional
5 embodiments, the homomeric multimer of the invention is at least a homodimer, at least a homotrimer, or at least a homotetramer.

As used herein, the term heteromer refers to a multimer containing one or more heterologous polypeptides (*i.e.*, polypeptides of different proteins) in addition to the polypeptides of the invention. In a specific embodiment, the multimer of the
10 invention is a heterodimer, a heterotrimer, or a heterotetramer. In additional embodiments, the heteromeric multimer of the invention is at least a heterodimer, at least a heterotrimer, or at least a heterotetramer.

Multimers of the invention may be the result of hydrophobic, hydrophilic, ionic and/or covalent associations and/or may be indirectly linked, by for example, liposome formation. Thus, in one embodiment, multimers of the invention, such as,
15 for example, homodimers or homotrimers, are formed when polypeptides of the invention contact one another in solution. In another embodiment, heteromultimers of the invention, such as, for example, heterotrimers or heterotetramers, are formed when polypeptides of the invention contact antibodies to the polypeptides of the
20 invention (including antibodies to the heterologous polypeptide sequence in a fusion protein of the invention) in solution. In other embodiments, multimers of the invention are formed by covalent associations with and/or between the polypeptides of the invention. Such covalent associations may involve one or more amino acid residues contained in the polypeptide sequence (*e.g.*, that recited in the sequence
25 listing, or contained in the polypeptide encoded by a deposited clone). In one instance, the covalent associations are cross-linking between cysteine residues located within the polypeptide sequences which interact in the native (*i.e.*, naturally occurring) polypeptide. In another instance, the covalent associations are the consequence of chemical or recombinant manipulation. Alternatively, such covalent
30 associations may involve one or more amino acid residues contained in the heterologous polypeptide sequence in a fusion protein of the invention.

In one example, covalent associations are between the heterologous sequence contained in a fusion protein of the invention (see, e.g., US Patent Number 5,478,925). In a specific example, the covalent associations are between the heterologous sequence contained in an Fc fusion protein of the invention (as
5 described herein). In another specific example, covalent associations of fusion proteins of the invention are between heterologous polypeptide sequence from another protein that is capable of forming covalently associated multimers, such as for example, osteoprotegerin (see, e.g., International Publication NO: WO 98/49305, the contents of which are herein incorporated by reference in its entirety). In another
10 embodiment, two or more polypeptides of the invention are joined through peptide linkers. Examples include those peptide linkers described in U.S. Pat. No. 5,073,627 (hereby incorporated by reference). Proteins comprising multiple polypeptides of the invention separated by peptide linkers may be produced using conventional recombinant DNA technology.

15 Another method for preparing multimer polypeptides of the invention involves use of polypeptides of the invention fused to a leucine zipper or isoleucine zipper polypeptide sequence. Leucine zipper and isoleucine zipper domains are polypeptides that promote multimerization of the proteins in which they are found. Leucine zippers were originally identified in several DNA-binding proteins (Landschulz et al.,
20 Science 240:1759, (1988)), and have since been found in a variety of different proteins. Among the known leucine zippers are naturally occurring peptides and derivatives thereof that dimerize or trimerize. Examples of leucine zipper domains suitable for producing soluble multimeric proteins of the invention are those described in PCT application WO 94/10308, hereby incorporated by reference. Recombinant
25 fusion proteins comprising a polypeptide of the invention fused to a polypeptide sequence that dimerizes or trimerizes in solution are expressed in suitable host cells, and the resulting soluble multimeric fusion protein is recovered from the culture supernatant using techniques known in the art.

30 Trimeric polypeptides of the invention may offer the advantage of enhanced biological activity. Preferred leucine zipper moieties and isoleucine moieties are those that preferentially form trimers. One example is a leucine zipper derived from

lung surfactant protein D (SPD), as described in Hoppe et al. (FEBS Letters 344:191, (1994)) and in U.S. patent application Ser. No. 08/446,922, hereby incorporated by reference. Other peptides derived from naturally occurring trimeric proteins may be employed in preparing trimeric polypeptides of the invention.

5 In another example, proteins of the invention are associated by interactions between Flag® polypeptide sequence contained in fusion proteins of the invention containing Flag® polypeptide sequence. In a further embodiment, associations
10 proteins of the invention are associated by interactions between heterologous polypeptide sequence contained in Flag® fusion proteins of the invention and anti-Flag® antibody.

 The multimers of the invention may be generated using chemical techniques known in the art. For example, polypeptides desired to be contained in the multimers of the invention may be chemically cross-linked using linker molecules and linker
15 molecule length optimization techniques known in the art (see, e.g., US Patent Number 5,478,925, which is herein incorporated by reference in its entirety).
Additionally, multimers of the invention may be generated using techniques known in the art to form one or more inter-molecule cross-links between the cysteine residues located within the sequence of the polypeptides desired to be contained in the
20 multimer (see, e.g., US Patent Number 5,478,925, which is herein incorporated by reference in its entirety). Further, polypeptides of the invention may be routinely modified by the addition of cysteine or biotin to the C terminus or N-terminus of the polypeptide and techniques known in the art may be applied to generate multimers containing one or more of these modified polypeptides (see, e.g., US Patent Number
25 5,478,925, which is herein incorporated by reference in its entirety). Additionally, techniques known in the art may be applied to generate liposomes containing the polypeptide components desired to be contained in the multimer of the invention (see, e.g., US Patent Number 5,478,925, which is herein incorporated by reference in its entirety).

 Alternatively, multimers of the invention may be generated using genetic
30 engineering techniques known in the art. In one embodiment, polypeptides contained in multimers of the invention are produced recombinantly using fusion protein

technology described herein or otherwise known in the art (see, e.g., US Patent Number 5,478,925, which is herein incorporated by reference in its entirety). In a specific embodiment, polynucleotides coding for a homodimer of the invention are generated by ligating a polynucleotide sequence encoding a polypeptide of the invention to a sequence encoding a linker polypeptide and then further to a synthetic polynucleotide encoding the translated product of the polypeptide in the reverse orientation from the original C-terminus to the N-terminus (lacking the leader sequence) (see, e.g., US Patent Number 5,478,925, which is herein incorporated by reference in its entirety). In another embodiment, recombinant techniques described herein or otherwise known in the art are applied to generate recombinant polypeptides of the invention which contain a transmembrane domain (or hydrophobic or signal peptide) and which can be incorporated by membrane reconstitution techniques into liposomes (see, e.g., US Patent Number 5,478,925, which is herein incorporated by reference in its entirety).

Uses of the Polynucleotides

Each of the polynucleotides identified herein can be used in numerous ways as reagents. The following description should be considered exemplary and utilizes known techniques.

The polynucleotides of the present invention are useful for chromosome identification. There exists an ongoing need to identify new chromosome markers, since few chromosome marking reagents, based on actual sequence data (repeat polymorphisms), are presently available. Each polynucleotide of the present invention can be used as a chromosome marker.

Briefly, sequences can be mapped to chromosomes by preparing PCR primers (preferably 15-25 bp) from the sequences shown in SEQ ID NO:X. Primers can be selected using computer analysis so that primers do not span more than one predicted exon in the genomic DNA. These primers are then used for PCR screening of somatic cell hybrids containing individual human chromosomes. Only those hybrids containing the human gene corresponding to the SEQ ID NO:X will yield an amplified fragment.

Similarly, somatic hybrids provide a rapid method of PCR mapping the polynucleotides to particular chromosomes. Three or more clones can be assigned per day using a single thermal cycler. Moreover, sublocalization of the polynucleotides can be achieved with panels of specific chromosome fragments. Other gene mapping strategies that can be used include in situ hybridization, prescreening with labeled flow-sorted chromosomes, and preselection by hybridization to construct chromosome specific-cDNA libraries.

Precise chromosomal location of the polynucleotides can also be achieved using fluorescence in situ hybridization (FISH) of a metaphase chromosomal spread. This technique uses polynucleotides as short as 500 or 600 bases; however, polynucleotides 2,000-4,000 bp are preferred. For a review of this technique, see Verma et al., "Human Chromosomes: a Manual of Basic Techniques," Pergamon Press, New York (1988).

For chromosome mapping, the polynucleotides can be used individually (to mark a single chromosome or a single site on that chromosome) or in panels (for marking multiple sites and/or multiple chromosomes). Preferred polynucleotides correspond to the noncoding regions of the cDNAs because the coding sequences are more likely conserved within gene families, thus increasing the chance of cross hybridization during chromosomal mapping.

Once a polynucleotide has been mapped to a precise chromosomal location, the physical position of the polynucleotide can be used in linkage analysis. Linkage analysis establishes coinheritance between a chromosomal location and presentation of a particular disease. (Disease mapping data are found, for example, in V. McKusick, Mendelian Inheritance in Man (available on line through Johns Hopkins University Welch Medical Library) .) Assuming 1 megabase mapping resolution and one gene per 20 kb, a cDNA precisely localized to a chromosomal region associated with the disease could be one of 50-500 potential causative genes.

Thus, once coinheritance is established, differences in the polynucleotide and the corresponding gene between affected and unaffected individuals can be examined. First, visible structural alterations in the chromosomes, such as deletions or translocations, are examined in chromosome spreads or by PCR. If no structural

alterations exist, the presence of point mutations are ascertained. Mutations observed in some or all affected individuals, but not in normal individuals, indicates that the mutation may cause the disease. However, complete sequencing of the polypeptide and the corresponding gene from several normal individuals is required to distinguish
5 the mutation from a polymorphism. If a new polymorphism is identified, this polymorphic polypeptide can be used for further linkage analysis.

Furthermore, increased or decreased expression of the gene in affected individuals as compared to unaffected individuals can be assessed using polynucleotides of the present invention. Any of these alterations (altered expression,
10 chromosomal rearrangement, or mutation) can be used as a diagnostic or prognostic marker.

Thus, the invention also provides a diagnostic method useful during diagnosis of a disorder, involving measuring the expression level of polynucleotides of the present invention in cells or body fluid from an individual and comparing the
15 measured gene expression level with a standard level of polynucleotide expression level, whereby an increase or decrease in the gene expression level compared to the standard is indicative of a disorder.

In still another embodiment, the invention includes a kit for analyzing samples for the presence of proliferative and/or cancerous polynucleotides derived from a test
20 subject. In a general embodiment, the kit includes at least one polynucleotide probe containing a nucleotide sequence that will specifically hybridize with a polynucleotide of the present invention and a suitable container. In a specific embodiment, the kit includes two polynucleotide probes defining an internal region of the polynucleotide of the present invention, where each probe has one strand
25 containing a 31' mer-end internal to the region. In a further embodiment, the probes may be useful as primers for polymerase chain reaction amplification.

Where a diagnosis of a disorder, has already been made according to conventional methods, the present invention is useful as a prognostic indicator, whereby patients exhibiting enhanced or depressed polynucleotide of the present
30 invention expression will experience a worse clinical outcome relative to patients expressing the gene at a level nearer the standard level.

By "measuring the expression level of polynucleotide of the present invention" is intended qualitatively or quantitatively measuring or estimating the level of the polypeptide of the present invention or the level of the mRNA encoding the polypeptide in a first biological sample either directly (e.g., by determining or
5 estimating absolute protein level or mRNA level) or relatively (e.g., by comparing to the polypeptide level or mRNA level in a second biological sample). Preferably, the polypeptide level or mRNA level in the first biological sample is measured or estimated and compared to a standard polypeptide level or mRNA level, the standard
10 being taken from a second biological sample obtained from an individual not having the disorder or being determined by averaging levels from a population of individuals not having a disorder. As will be appreciated in the art, once a standard polypeptide level or mRNA level is known, it can be used repeatedly as a standard for comparison.

By "biological sample" is intended any biological sample obtained from an
15 individual, body fluid, cell line, tissue culture, or other source which contains the polypeptide of the present invention or mRNA. As indicated, biological samples include body fluids (such as semen, lymph, sera, plasma, urine, synovial fluid and spinal fluid) which contain the polypeptide of the present invention, and other tissue sources found to express the polypeptide of the present invention. Methods for
20 obtaining tissue biopsies and body fluids from mammals are well known in the art. Where the biological sample is to include mRNA, a tissue biopsy is the preferred source.

The method(s) provided above may preferably be applied in a diagnostic method and/or kits in which polynucleotides and/or polypeptides are attached to a
25 solid support. In one exemplary method, the support may be a "gene chip" or a "biological chip" as described in US Patents 5,837,832, 5,874,219, and 5,856,174. Further, such a gene chip with polynucleotides of the present invention attached may be used to identify polymorphisms between the polynucleotide sequences, with polynucleotides isolated from a test subject. The knowledge of such polymorphisms
30 (i.e. their location, as well as, their existence) would be beneficial in identifying disease loci for many disorders, including cancerous diseases and conditions. Such a

method is described in US Patents 5,858,659 and 5,856,104. The US Patents referenced supra are hereby incorporated by reference in their entirety herein.

The present invention encompasses polynucleotides of the present invention that are chemically synthesized, or reproduced as peptide nucleic acids (PNA), or according to other methods known in the art. The use of PNAs would serve as the preferred form if the polynucleotides are incorporated onto a solid support, or gene chip. For the purposes of the present invention, a peptide nucleic acid (PNA) is a polyamide type of DNA analog and the monomeric units for adenine, guanine, thymine and cytosine are available commercially (Perceptive Biosystems). Certain components of DNA, such as phosphorus, phosphorus oxides, or deoxyribose derivatives, are not present in PNAs. As disclosed by P. E. Nielsen, M. Egholm, R. H. Berg and O. Buchardt, *Science* 254, 1497 (1991); and M. Egholm, O. Buchardt, L. Christensen, C. Behrens, S. M. Freier, D. A. Driver, R. H. Berg, S. K. Kim, B. Norden, and P. E. Nielsen, *Nature* 365, 666 (1993), PNAs bind specifically and tightly to complementary DNA strands and are not degraded by nucleases. In fact, PNA binds more strongly to DNA than DNA itself does. This is probably because there is no electrostatic repulsion between the two strands, and also the polyamide backbone is more flexible. Because of this, PNA/DNA duplexes bind under a wider range of stringency conditions than DNA/DNA duplexes, making it easier to perform multiplex hybridization. Smaller probes can be used than with DNA due to the strong binding. In addition, it is more likely that single base mismatches can be determined with PNA/DNA hybridization because a single mismatch in a PNA/DNA 15-mer lowers the melting point ($T_{sub.m}$) by 8°-20° C, vs. 4°-16° C for the DNA/DNA 15-mer duplex. Also, the absence of charge groups in PNA means that hybridization can be done at low ionic strengths and reduce possible interference by salt during the analysis.

The present invention is useful for detecting cancer in mammals. In particular the invention is useful during diagnosis of pathological cell proliferative neoplasias which include, but are not limited to: acute myelogenous leukemias including acute monocytic leukemia, acute myeloblastic leukemia, acute promyelocytic leukemia, acute myelomonocytic leukemia, acute erythroleukemia, acute megakaryocytic

leukemia, and acute undifferentiated leukemia, etc.; and chronic myelogenous leukemias including chronic myelomonocytic leukemia, chronic granulocytic leukemia, etc. Preferred mammals include monkeys, apes, cats, dogs, cows, pigs, horses, rabbits and humans. Particularly preferred are humans.

5 Pathological cell proliferative disorders are often associated with inappropriate activation of proto-oncogenes. (Germann, E. P. et al., "The Etiology of Acute Leukemia: Molecular Genetics and Viral Oncology," in *Neoplastic Diseases of the Blood*, Vol 1., Wiernik, P. H. et al. eds., 161-182 (1985)). Neoplasias are now
10 believed to result from the qualitative alteration of a normal cellular gene product, or from the quantitative modification of gene expression by insertion into the
15 chromosome of a viral sequence, by chromosomal translocation of a gene to a more actively transcribed region, or by some other mechanism. (Germann et al., supra) It is likely that mutated or altered expression of specific genes is involved in the
15 pathogenesis of some leukemias, among other tissues and cell types. (Germann et al., supra) Indeed, the human counterparts of the oncogenes involved in some animal
neoplasias have been amplified or translocated in some cases of human leukemia and carcinoma. (Germann et al., supra).

 For example, c-myc expression is highly amplified in the non-lymphocytic leukemia cell line HL-60. When HL-60 cells are chemically induced to stop
20 proliferation, the level of c-myc is found to be downregulated. (International Publication Number WO 91/15580) However, it has been shown that exposure of
HL-60 cells to a DNA construct that is complementary to the 5' end of c-myc or c-myb blocks translation of the corresponding mRNAs which downregulates expression
of the c-myc or c-myb proteins and causes arrest of cell proliferation and
25 differentiation of the treated cells. (International Publication Number WO 91/15580; Wickstrom et al., *Proc. Natl. Acad. Sci.* 85:1028 (1988); Anfossi et al., *Proc. Natl. Acad. Sci.* 86:3379 (1989)). However, the skilled artisan would appreciate the
present invention's usefulness would not be limited to treatment of proliferative
disorders of hematopoietic cells and tissues, in light of the numerous cells and cell
30 types of varying origins which are known to exhibit proliferative phenotypes.

In addition to the foregoing, a polynucleotide can be used to control gene expression through triple helix formation or antisense DNA or RNA. Antisense techniques are discussed, for example, in Okano, J. *Neurochem.* 56: 560 (1991); "Oligodeoxynucleotides as Antisense Inhibitors of Gene Expression, CRC Press, Boca Raton, FL (1988). Triple helix formation is discussed in, for instance Lee et al., *Nucleic Acids Research* 6: 3073 (1979); Cooney et al., *Science* 241: 456 (1988); and Dervan et al., *Science* 251: 1360 (1991). Both methods rely on binding of the polynucleotide to a complementary DNA or RNA. For these techniques, preferred polynucleotides are usually oligonucleotides 20 to 40 bases in length and complementary to either the region of the gene involved in transcription (triple helix - see Lee et al., *Nucl. Acids Res.* 6:3073 (1979); Cooney et al., *Science* 241:456 (1988); and Dervan et al., *Science* 251:1360 (1991)) or to the mRNA itself (antisense - Okano, J. *Neurochem.* 56:560 (1991); Oligodeoxy-nucleotides as Antisense Inhibitors of Gene Expression, CRC Press, Boca Raton, FL (1988).) Triple helix formation optimally results in a shut-off of RNA transcription from DNA, while antisense RNA hybridization blocks translation of an mRNA molecule into polypeptide. Both techniques are effective in model systems, and the information disclosed herein can be used to design antisense or triple helix polynucleotides in an effort to treat disease.

Polynucleotides of the present invention are also useful in gene therapy. One goal of gene therapy is to insert a normal gene into an organism having a defective gene, in an effort to correct the genetic defect. The polynucleotides disclosed in the present invention offer a means of targeting such genetic defects in a highly accurate manner. Another goal is to insert a new gene that was not present in the host genome, thereby producing a new trait in the host cell.

The polynucleotides are also useful for identifying individuals from minute biological samples. The United States military, for example, is considering the use of restriction fragment length polymorphism (RFLP) for identification of its personnel. In this technique, an individual's genomic DNA is digested with one or more restriction enzymes, and probed on a Southern blot to yield unique bands for identifying personnel. This method does not suffer from the current limitations of

"Dog Tags" which can be lost, switched, or stolen, making positive identification difficult. The polynucleotides of the present invention can be used as additional DNA markers for RFLP.

5 The polynucleotides of the present invention can also be used as an alternative to RFLP, by determining the actual base-by-base DNA sequence of selected portions of an individual's genome. These sequences can be used to prepare PCR primers for amplifying and isolating such selected DNA, which can then be sequenced. Using this technique, individuals can be identified because each individual will have a unique set of DNA sequences. Once a unique ID database is established for an individual, positive identification of that individual, living or dead, can be made from
10 extremely small tissue samples.

Forensic biology also benefits from using DNA-based identification techniques as disclosed herein. DNA sequences taken from very small biological samples such as tissues, e.g., hair or skin, or body fluids, e.g., blood, saliva, semen,
15 synovial fluid, amniotic fluid, breast milk, lymph, pulmonary sputum or surfactant, urine, fecal matter, etc., can be amplified using PCR. In one prior art technique, gene sequences amplified from polymorphic loci, such as DQa class II HLA gene, are used in forensic biology to identify individuals. (Erlich, H., PCR Technology, Freeman and Co. (1992).) Once these specific polymorphic loci are
20 amplified, they are digested with one or more restriction enzymes, yielding an identifying set of bands on a Southern blot probed with DNA corresponding to the DQa class II HLA gene. Similarly, polynucleotides of the present invention can be used as polymorphic markers for forensic purposes.

There is also a need for reagents capable of identifying the source of a
25 particular tissue. Such need arises, for example, in forensics when presented with tissue of unknown origin. Appropriate reagents can comprise, for example, DNA probes or primers specific to particular tissue prepared from the sequences of the present invention. Panels of such reagents can identify tissue by species and/or by organ type. In a similar fashion, these reagents can be used to screen tissue cultures
30 for contamination.

In the very least, the polynucleotides of the present invention can be used as molecular weight markers on Southern gels, as diagnostic probes for the presence of a specific mRNA in a particular cell type, as a probe to "subtract-out" known sequences in the process of discovering novel polynucleotides, for selecting and making
5 oligomers for attachment to a "gene chip" or other support, to raise anti-DNA antibodies using DNA immunization techniques, and as an antigen to elicit an immune response.

Uses of the Polypeptides

10 Each of the polypeptides identified herein can be used in numerous ways. The following description should be considered exemplary and utilizes known techniques.

A polypeptide of the present invention can be used to assay protein levels in a biological sample using antibody-based techniques. For example, protein expression in tissues can be studied with classical immunohistological methods. (Jalkanen, M.,
15 et al., J. Cell. Biol. 101:976-985 (1985); Jalkanen, M., et al., J. Cell. Biol. 105:3087-3096 (1987).) Other antibody-based methods useful for detecting protein gene expression include immunoassays, such as the enzyme linked immunosorbent assay (ELISA) and the radioimmunoassay (RIA). Suitable antibody assay labels are known in the art and include enzyme labels, such as, glucose oxidase, and radioisotopes, such
20 as iodine (125I, 121I), carbon (14C), sulfur (35S), tritium (3H), indium (112In), and technetium (99mTc), and fluorescent labels, such as fluorescein and rhodamine, and biotin.

In addition to assaying secreted protein levels in a biological sample, proteins can also be detected in vivo by imaging. Antibody labels or markers for in vivo
25 imaging of protein include those detectable by X-radiography, NMR or ESR. For X-radiography, suitable labels include radioisotopes such as barium or cesium, which emit detectable radiation but are not overtly harmful to the subject. Suitable markers for NMR and ESR include those with a detectable characteristic spin, such as deuterium, which may be incorporated into the antibody by labeling of nutrients for
30 the relevant hybridoma.

A protein-specific antibody or antibody fragment which has been labeled with an appropriate detectable imaging moiety, such as a radioisotope (for example, ^{131}I , ^{112}In , $^{99\text{m}}\text{Tc}$), a radio-opaque substance, or a material detectable by nuclear magnetic resonance, is introduced (for example, parenterally, subcutaneously, or intraperitoneally) into the mammal. It will be understood in the art that the size of the subject and the imaging system used will determine the quantity of imaging moiety needed to produce diagnostic images. In the case of a radioisotope moiety, for a human subject, the quantity of radioactivity injected will normally range from about 5 to 20 millicuries of $^{99\text{m}}\text{Tc}$. The labeled antibody or antibody fragment will then preferentially accumulate at the location of cells which contain the specific protein. In vivo tumor imaging is described in S.W. Burchiel et al., "Immunopharmacokinetics of Radiolabeled Antibodies and Their Fragments." (Chapter 13 in Tumor Imaging: The Radiochemical Detection of Cancer, S.W. Burchiel and B. A. Rhodes, eds., Masson Publishing Inc. (1982).)

Thus, the invention provides a diagnostic method of a disorder, which involves (a) assaying the expression of a polypeptide of the present invention in cells or body fluid of an individual; (b) comparing the level of gene expression with a standard gene expression level, whereby an increase or decrease in the assayed polypeptide gene expression level compared to the standard expression level is indicative of a disorder. With respect to cancer, the presence of a relatively high amount of transcript in biopsied tissue from an individual may indicate a predisposition for the development of the disease, or may provide a means for detecting the disease prior to the appearance of actual clinical symptoms. A more definitive diagnosis of this type may allow health professionals to employ preventative measures or aggressive treatment earlier thereby preventing the development or further progression of the cancer.

Moreover, polypeptides of the present invention can be used to treat disease. For example, patients can be administered a polypeptide of the present invention in an effort to replace absent or decreased levels of the polypeptide (e.g., insulin), to supplement absent or decreased levels of a different polypeptide (e.g., hemoglobin S for hemoglobin B, SOD, catalase, DNA repair proteins), to inhibit the activity of a

polypeptide (e.g., an oncogene or tumor suppressor), to activate the activity of a polypeptide (e.g., by binding to a receptor), to reduce the activity of a membrane bound receptor by competing with it for free ligand (e.g., soluble TNF receptors used in reducing inflammation), or to bring about a desired response (e.g., blood vessel growth inhibition, enhancement of the immune response to proliferative cells or tissues).

Similarly, antibodies directed to a polypeptide of the present invention can also be used to treat disease. For example, administration of an antibody directed to a polypeptide of the present invention can bind and reduce overproduction of the polypeptide. Similarly, administration of an antibody can activate the polypeptide, such as by binding to a polypeptide bound to a membrane (receptor).

At the very least, the polypeptides of the present invention can be used as molecular weight markers on SDS-PAGE gels or on molecular sieve gel filtration columns using methods well known to those of skill in the art. Polypeptides can also be used to raise antibodies, which in turn are used to measure protein expression from a recombinant cell, as a way of assessing transformation of the host cell. Moreover, the polypeptides of the present invention can be used to test the following biological activities.

Gene Therapy Methods

Another aspect of the present invention is to gene therapy methods for treating disorders, diseases and conditions. The gene therapy methods relate to the introduction of nucleic acid (DNA, RNA and antisense DNA or RNA) sequences into an animal to achieve expression of a polypeptide of the present invention. This method requires a polynucleotide which codes for a polypeptide of the invention that operatively linked to a promoter and any other genetic elements necessary for the expression of the polypeptide by the target tissue. Such gene therapy and delivery techniques are known in the art, see, for example, WO90/11092, which is herein incorporated by reference.

Thus, for example, cells from a patient may be engineered with a polynucleotide (DNA or RNA) comprising a promoter operably linked to a

polynucleotide of the invention *ex vivo*, with the engineered cells then being provided to a patient to be treated with the polypeptide. Such methods are well-known in the art. For example, see Belldégrun et al., J. Natl. Cancer Inst., 85:207-216 (1993); Ferrantini et al., Cancer Research, 53:107-1112 (1993); Ferrantini et al., J. Immunology 153: 4604-4615 (1994); Kaido, T., et al., Int. J. Cancer 60: 221-229 (1995); Ogura et al., Cancer Research 50: 5102-5106 (1990); Santodonato, et al., Human Gene Therapy 7:1-10 (1996); Santodonato, et al., Gene Therapy 4:1246-1255 (1997); and Zhang, et al., Cancer Gene Therapy 3: 31-38 (1996)), which are herein incorporated by reference. In one embodiment, the cells which are engineered are arterial cells. The arterial cells may be reintroduced into the patient through direct injection to the artery, the tissues surrounding the artery, or through catheter injection.

As discussed in more detail below, the polynucleotide constructs can be delivered by any method that delivers injectable materials to the cells of an animal, such as, injection into the interstitial space of tissues (heart, muscle, skin, lung, liver, and the like). The polynucleotide constructs may be delivered in a pharmaceutically acceptable liquid or aqueous carrier.

In one embodiment, the polynucleotide of the invention is delivered as a naked polynucleotide. The term "naked" polynucleotide, DNA or RNA refers to sequences that are free from any delivery vehicle that acts to assist, promote or facilitate entry into the cell, including viral sequences, viral particles, liposome formulations, lipofectin or precipitating agents and the like. However, the polynucleotides of the invention can also be delivered in liposome formulations and lipofectin formulations and the like can be prepared by methods well known to those skilled in the art. Such methods are described, for example, in U.S. Patent Nos. 5,593,972, 5,589,466, and 5,580,859, which are herein incorporated by reference.

The polynucleotide vector constructs of the invention used in the gene therapy method are preferably constructs that will not integrate into the host genome nor will they contain sequences that allow for replication. Appropriate vectors include pWLNEO, pSV2CAT, pOG44, pXT1 and pSG available from Stratagene; pSVK3, pBPV, pMSG and pSVL available from Pharmacia; and pEF1/V5,

pcDNA3.1, and pRc/CMV2 available from Invitrogen. Other suitable vectors will be readily apparent to the skilled artisan.

Any strong promoter known to those skilled in the art can be used for driving the expression of polynucleotide sequence of the invention. Suitable promoters
5 include adenoviral promoters, such as the adenoviral major late promoter; or heterologous promoters, such as the cytomegalovirus (CMV) promoter; the respiratory syncytial virus (RSV) promoter; inducible promoters, such as the MMT promoter, the metallothionein promoter; heat shock promoters; the albumin promoter; the ApoAI promoter; human globin promoters; viral thymidine kinase promoters,
10 such as the Herpes Simplex thymidine kinase promoter; retroviral LTRs; the b-actin promoter; and human growth hormone promoters. The promoter also may be the native promoter for the polynucleotides of the invention.

Unlike other gene therapy techniques, one major advantage of introducing naked nucleic acid sequences into target cells is the transitory nature of the
15 polynucleotide synthesis in the cells. Studies have shown that non-replicating DNA sequences can be introduced into cells to provide production of the desired polypeptide for periods of up to six months.

The polynucleotide construct of the invention can be delivered to the interstitial space of tissues within the an animal, including of muscle, skin, brain, lung, liver,
20 spleen, bone marrow, thymus, heart, lymph, blood, bone, cartilage, pancreas, kidney, gall bladder, stomach, intestine, testis, ovary, uterus, rectum, nervous system, eye, gland, and connective tissue. Interstitial space of the tissues comprises the intercellular, fluid, mucopolysaccharide matrix among the reticular fibers of organ tissues, elastic fibers in the walls of vessels or chambers, collagen fibers of fibrous tissues, or that
25 same matrix within connective tissue ensheathing muscle cells or in the lacunae of bone. It is similarly the space occupied by the plasma of the circulation and the lymph fluid of the lymphatic channels. Delivery to the interstitial space of muscle tissue is preferred for the reasons discussed below. They may be conveniently delivered by injection into the tissues comprising these cells. They are preferably delivered to and
30 expressed in persistent, non-dividing cells which are differentiated, although delivery and expression may be achieved in non-differentiated or less completely differentiated

cells, such as, for example, stem cells of blood or skin fibroblasts. *In vivo* muscle cells are particularly competent in their ability to take up and express polynucleotides.

For the naked *nucleic acid* sequence injection, an effective dosage amount of DNA or RNA will be in the range of from about 0.05 mg/kg body weight to about 50 mg/kg body weight. Preferably the dosage will be from about 0.005 mg/kg to about 20 mg/kg and more preferably from about 0.05 mg/kg to about 5 mg/kg. Of course, as the artisan of ordinary skill will appreciate, this dosage will vary according to the tissue site of injection. The appropriate and effective dosage of nucleic acid sequence can readily be determined by those of ordinary skill in the art and may depend on the condition being treated and the route of administration.

The preferred route of administration is by the parenteral route of injection into the interstitial space of tissues. However, other parenteral routes may also be used, such as, inhalation of an aerosol formulation particularly for delivery to lungs or bronchial tissues, throat or mucous membranes of the nose. In addition, naked DNA constructs can be delivered to arteries during angioplasty by the catheter used in the procedure.

The naked polynucleotides are delivered by any method known in the art, including, but not limited to, direct needle injection at the delivery site, intravenous injection, topical administration, catheter infusion, and so-called "gene guns". These delivery methods are known in the art.

The constructs may also be delivered with delivery vehicles such as viral sequences, viral particles, liposome formulations, lipofectin, precipitating agents, etc. Such methods of delivery are known in the art.

In certain embodiments, the polynucleotide constructs of the invention are complexed in a liposome preparation. Liposomal preparations for use in the instant invention include cationic (positively charged), anionic (negatively charged) and neutral preparations. However, cationic liposomes are particularly preferred because a tight charge complex can be formed between the cationic liposome and the polyanionic nucleic acid. Cationic liposomes have been shown to mediate intracellular delivery of plasmid DNA (Felgner et al., Proc. Natl. Acad. Sci. USA, 84:7413-7416 (1987), which is herein incorporated by reference); mRNA (Malone et

al., Proc. Natl. Acad. Sci. USA , 86:6077-6081 (1989), which is herein incorporated by reference); and purified transcription factors (Debs et al., J. Biol. Chem., 265:10189-10192 (1990), which is herein incorporated by reference), in functional form.

5 Cationic liposomes are readily available. For example, N[1-2,3-dioleoyloxy)propyl]-N,N,N-triethylammonium (DOTMA) liposomes are particularly useful and are available under the trademark Lipofectin, from GIBCO BRL, Grand Island, N.Y. (See, also, Felgner et al., Proc. Natl Acad. Sci. USA , 84:7413-7416 (1987), which is herein incorporated by reference). Other commercially
10 available liposomes include transfectace (DDAB/DOPE) and DOTAP/DOPE (Boehringer).

Other cationic liposomes can be prepared from readily available materials using techniques well known in the art. See, e.g. PCT Publication NO: WO 90/11092 (which is herein incorporated by reference) for a description of the synthesis of
15 DOTAP (1,2-bis(oleoyloxy)-3-(trimethylammonio)propane) liposomes. Preparation of DOTMA liposomes is explained in the literature, see, e.g., Felgner et al., Proc. Natl. Acad. Sci. USA, 84:7413-7417, which is herein incorporated by reference. Similar methods can be used to prepare liposomes from other cationic lipid materials.

Similarly, anionic and neutral liposomes are readily available, such as from
20 Avanti Polar Lipids (Birmingham, Ala.), or can be easily prepared using readily available materials. Such materials include phosphatidyl, choline, cholesterol, phosphatidyl ethanolamine, dioleoylphosphatidyl choline (DOPC), dioleoylphosphatidyl glycerol (DOPG), dioleoylphosphatidyl ethanolamine (DOPE), among others. These materials can also be mixed with the DOTMA and DOTAP
25 starting materials in appropriate ratios. Methods for making liposomes using these materials are well known in the art.

For example, commercially dioleoylphosphatidyl choline (DOPC), dioleoylphosphatidyl glycerol (DOPG), and dioleoylphosphatidyl ethanolamine (DOPE) can be used in various combinations to make conventional liposomes, with or
30 without the addition of cholesterol. Thus, for example, DOPG/DOPC vesicles can be prepared by drying 50 mg each of DOPG and DOPC under a stream of nitrogen gas

into a sonication vial. The sample is placed under a vacuum pump overnight and is hydrated the following day with deionized water. The sample is then sonicated for 2 hours in a capped vial, using a Heat Systems model 350 sonicator equipped with an inverted cup (bath type) probe at the maximum setting while the bath is circulated at 15EC. Alternatively, negatively charged vesicles can be prepared without sonication to produce multilamellar vesicles or by extrusion through nucleopore membranes to produce unilamellar vesicles of discrete size. Other methods are known and available to those of skill in the art.

The liposomes can comprise multilamellar vesicles (MLVs), small unilamellar vesicles (SUVs), or large unilamellar vesicles (LUVs), with SUVs being preferred. The various liposome-nucleic acid complexes are prepared using methods well known in the art. See, e.g., Straubinger et al., *Methods of Immunology*, 101:512-527 (1983), which is herein incorporated by reference. For example, MLVs containing nucleic acid can be prepared by depositing a thin film of phospholipid on the walls of a glass tube and subsequently hydrating with a solution of the material to be encapsulated. SUVs are prepared by extended sonication of MLVs to produce a homogeneous population of unilamellar liposomes. The material to be entrapped is added to a suspension of preformed MLVs and then sonicated. When using liposomes containing cationic lipids, the dried lipid film is resuspended in an appropriate solution such as sterile water or an isotonic buffer solution such as 10 mM Tris/NaCl, sonicated, and then the preformed liposomes are mixed directly with the DNA. The liposome and DNA form a very stable complex due to binding of the positively charged liposomes to the cationic DNA. SUVs find use with small nucleic acid fragments. LUVs are prepared by a number of methods, well known in the art. Commonly used methods include Ca^{2+} -EDTA chelation (Papahadjopoulos et al., *Biochim. Biophys. Acta*, 394:483 (1975); Wilson et al., *Cell*, 17:77 (1979)); ether injection (Deamer et al., *Biochim. Biophys. Acta*, 443:629 (1976); Ostro et al., *Biochem. Biophys. Res. Commun.*, 76:836 (1977); Fraley et al., *Proc. Natl. Acad. Sci. USA*, 76:3348 (1979)); detergent dialysis (Enoch et al., *Proc. Natl. Acad. Sci. USA*, 76:145 (1979)); and reverse-phase evaporation (REV) (Fraley et al., *J. Biol. Chem.*, 255:10431 (1980);

Szoka et al., Proc. Natl. Acad. Sci. USA , 75:145 (1978); Schaefer-Ridder et al., Science, 215:166 (1982)), which are herein incorporated by reference.

Generally, the ratio of DNA to liposomes will be from about 10:1 to about 1:10. Preferably, the ration will be from about 5:1 to about 1:5. More preferably, the
5 ration will be about 3:1 to about 1:3. Still more preferably, the ratio will be about 1:1.

U.S. Patent NO: 5,676,954 (which is herein incorporated by reference) reports on the injection of genetic material, complexed with cationic liposomes carriers, into mice. U.S. Patent Nos. 4,897,355, 4,946,787, 5,049,386, 5,459,127, 5,589,466, 5,693,622, 5,580,859, 5,703,055, and international publication NO: WO 94/9469
10 (which are herein incorporated by reference) provide cationic lipids for use in transfecting DNA into cells and mammals. U.S. Patent Nos. 5,589,466, 5,693,622, 5,580,859, 5,703,055, and international publication NO: WO 94/9469 (which are herein incorporated by reference) provide methods for delivering DNA-cationic lipid complexes to mammals.

15 In certain embodiments, cells are engineered, *ex vivo* or *in vivo*, using a retroviral particle containing RNA which comprises a sequence encoding polypeptides of the invention. Retroviruses from which the retroviral plasmid vectors may be derived include, but are not limited to, Moloney Murine Leukemia Virus, spleen necrosis virus, Rous sarcoma Virus, Harvey Sarcoma Virus, avian leukosis
20 virus, gibbon ape leukemia virus, human immunodeficiency virus, Myeloproliferative Sarcoma Virus, and mammary tumor virus.

The retroviral plasmid vector is employed to transduce packaging cell lines to form producer cell lines. Examples of packaging cells which may be transfected include, but are not limited to, the PE501, PA317, R-2, R-AM, PA12, T19-14X, VT-
25 19-17-H2, RCRE, RCRIP, GP+E-86, GP+envAm12, and DAN cell lines as described in Miller, Human Gene Therapy , 1:5-14 (1990), which is incorporated herein by reference in its entirety. The vector may transduce the packaging cells through any means known in the art. Such means include, but are not limited to, electroporation, the use of liposomes, and CaPO₄ precipitation. In one alternative, the retroviral
30 plasmid vector may be encapsulated into a liposome, or coupled to a lipid, and then administered to a host.

The producer cell line generates infectious retroviral vector particles which include polynucleotide encoding polypeptides of the invention. Such retroviral vector particles then may be employed, to transduce eukaryotic cells, either *in vitro* or *in vivo*. The transduced eukaryotic cells will express polypeptides of the invention.

5 In certain other embodiments, cells are engineered, *ex vivo* or *in vivo*, with polynucleotides of the invention contained in an adenovirus vector. Adenovirus can be manipulated such that it encodes and expresses polypeptides of the invention, and at the same time is inactivated in terms of its ability to replicate in a normal lytic viral life cycle. Adenovirus expression is achieved without integration of the viral DNA
10 into the host cell chromosome, thereby alleviating concerns about insertional mutagenesis. Furthermore, adenoviruses have been used as live enteric vaccines for many years with an excellent safety profile (Schwartz et al., Am. Rev. Respir. Dis., 109:233-238 (1974)). Finally, adenovirus mediated gene transfer has been demonstrated in a number of instances including transfer of alpha-1-antitrypsin and
15 CFTR to the lungs of cotton rats (Rosenfeld et al., Science, 252:431-434 (1991); Rosenfeld et al., Cell, 68:143-155 (1992)). Furthermore, extensive studies to attempt to establish adenovirus as a causative agent in human cancer were uniformly negative (Green et al. Proc. Natl. Acad. Sci. USA, 76:6606 (1979)).

Suitable adenoviral vectors useful in the present invention are described, for
20 example, in Kozarsky and Wilson, Curr. Opin. Genet. Devel., 3:499-503 (1993); Rosenfeld et al., Cell, 68:143-155 (1992); Engelhardt et al., Human Genet. Ther., 4:759-769 (1993); Yang et al., Nature Genet., 7:362-369 (1994); Wilson et al., Nature, 365:691-692 (1993); and U.S. Patent NO: 5,652,224, which are herein incorporated by reference. For example, the adenovirus vector Ad2 is useful and can
25 be grown in human 293 cells. These cells contain the E1 region of adenovirus and constitutively express Ela and Elb, which complement the defective adenoviruses by providing the products of the genes deleted from the vector. In addition to Ad2, other varieties of adenovirus (e.g., Ad3, Ad5, and Ad7) are also useful in the present invention.

30 Preferably, the adenoviruses used in the present invention are replication deficient. Replication deficient adenoviruses require the aid of a helper virus and/or

packaging cell line to form infectious particles. The resulting virus is capable of infecting cells and can express a polynucleotide of interest which is operably linked to a promoter, but cannot replicate in most cells. Replication deficient adenoviruses may be deleted in one or more of all or a portion of the following genes: E1a, E1b,
5 E3, E4, E2a, or L1 through L5.

In certain other embodiments, the cells are engineered, *ex vivo* or *in vivo*, using an adeno-associated virus (AAV). AAVs are naturally occurring defective viruses that require helper viruses to produce infectious particles (Muzyczka, Curr. Topics in Microbiol. Immunol., 158:97 (1992)). It is also one of the few viruses that
10 may integrate its DNA into non-dividing cells. Vectors containing as little as 300 base pairs of AAV can be packaged and can integrate, but space for exogenous DNA is limited to about 4.5 kb. Methods for producing and using such AAVs are known in the art. See, for example, U.S. Patent Nos. 5,139,941, 5,173,414, 5,354,678, 5,436,146, 5,474,935, 5,478,745, and 5,589,377.

For example, an appropriate AAV vector for use in the present invention will include all the sequences necessary for DNA replication, encapsidation, and host-cell integration. The polynucleotide construct containing polynucleotides of the invention is inserted into the AAV vector using standard cloning methods, such as those found in Sambrook et al., Molecular Cloning: A Laboratory Manual, Cold Spring Harbor
20 Press (1989). The recombinant AAV vector is then transfected into packaging cells which are infected with a helper virus, using any standard technique, including lipofection, electroporation, calcium phosphate precipitation, etc. Appropriate helper viruses include adenoviruses, cytomegaloviruses, vaccinia viruses, or herpes viruses. Once the packaging cells are transfected and infected, they will produce infectious
25 AAV viral particles which contain the polynucleotide construct of the invention. These viral particles are then used to transduce eukaryotic cells, either *ex vivo* or *in vivo*. The transduced cells will contain the polynucleotide construct integrated into its genome, and will express the desired gene product.

Another method of gene therapy involves operably associating heterologous
30 control regions and endogenous polynucleotide sequences (e.g. encoding the polypeptide sequence of interest) via homologous recombination (see, e.g., U.S.

Patent NO: 5,641,670, issued June 24, 1997; International Publication NO: WO 96/29411, published September 26, 1996; International Publication NO: WO 94/12650, published August 4, 1994; Koller et al., Proc. Natl. Acad. Sci. USA, 86:8932-8935 (1989); and Zijlstra et al., Nature, 342:435-438 (1989). This method
5 involves the activation of a gene which is present in the target cells, but which is not normally expressed in the cells, or is expressed at a lower level than desired.

Polynucleotide constructs are made, using standard techniques known in the art, which contain the promoter with targeting sequences flanking the promoter. Suitable promoters are described herein. The targeting sequence is sufficiently
10 complementary to an endogenous sequence to permit homologous recombination of the promoter-targeting sequence with the endogenous sequence. The targeting sequence will be sufficiently near the 5' end of the desired endogenous polynucleotide sequence so the promoter will be operably linked to the endogenous sequence upon homologous recombination.

15 The promoter and the targeting sequences can be amplified using PCR. Preferably, the amplified promoter contains distinct restriction enzyme sites on the 5' and 3' ends. Preferably, the 3' end of the first targeting sequence contains the same restriction enzyme site as the 5' end of the amplified promoter and the 5' end of the second targeting sequence contains the same restriction site as the 3' end of the
20 amplified promoter. The amplified promoter and targeting sequences are digested and ligated together.

The promoter-targeting sequence construct is delivered to the cells, either as naked polynucleotide, or in conjunction with transfection-facilitating agents, such as liposomes, viral sequences, viral particles, whole viruses, lipofection, precipitating
25 agents, etc., described in more detail above. The P promoter-targeting sequence can be delivered by any method, included direct needle injection, intravenous injection, topical administration, catheter infusion, particle accelerators, etc. The methods are described in more detail below.

The promoter-targeting sequence construct is taken up by cells. Homologous
30 recombination between the construct and the endogenous sequence takes place, such

that an endogenous sequence is placed under the control of the promoter. The promoter then drives the expression of the endogenous sequence.

The polynucleotides encoding polypeptides of the present invention may be administered along with other polynucleotides encoding other angiogenic proteins. Angiogenic proteins include, but are not limited to, acidic and basic fibroblast growth factors, VEGF-1, VEGF-2 (VEGF-C), VEGF-3 (VEGF-B), epidermal growth factor alpha and beta, platelet-derived endothelial cell growth factor, platelet-derived growth factor, tumor necrosis factor alpha, hepatocyte growth factor, insulin like growth factor, colony stimulating factor, macrophage colony stimulating factor, granulocyte/macrophage colony stimulating factor, and nitric oxide synthase.

Preferably, the polynucleotide encoding a polypeptide of the invention contains a secretory signal sequence that facilitates secretion of the protein. Typically, the signal sequence is positioned in the coding region of the polynucleotide to be expressed towards or at the 5' end of the coding region. The signal sequence may be homologous or heterologous to the polynucleotide of interest and may be homologous or heterologous to the cells to be transfected. Additionally, the signal sequence may be chemically synthesized using methods known in the art.

Any mode of administration of any of the above-described polynucleotides constructs can be used so long as the mode results in the expression of one or more molecules in an amount sufficient to provide a therapeutic effect. This includes direct needle injection, systemic injection, catheter infusion, biolistic injectors, particle accelerators (i.e., "gene guns"), gelfoam sponge depots, other commercially available depot materials, osmotic pumps (e.g., Alza minipumps), oral or suppository solid (tablet or pill) pharmaceutical formulations, and decanting or topical applications during surgery. For example, direct injection of naked calcium phosphate-precipitated plasmid into rat liver and rat spleen or a protein-coated plasmid into the portal vein has resulted in gene expression of the foreign gene in the rat livers. (Kaneda et al., Science, 243:375 (1989)).

A preferred method of local administration is by direct injection. Preferably, a recombinant molecule of the present invention complexed with a delivery vehicle is administered by direct injection into or locally within the area of arteries.

Administration of a composition locally within the area of arteries refers to injecting the composition centimeters and preferably, millimeters within arteries.

Another method of local administration is to contact a polynucleotide construct of the present invention in or around a surgical wound. For example, a patient can undergo surgery and the polynucleotide construct can be coated on the surface of tissue inside the wound or the construct can be injected into areas of tissue inside the wound.

Therapeutic compositions useful in systemic administration, include recombinant molecules of the present invention complexed to a targeted delivery vehicle of the present invention. Suitable delivery vehicles for use with systemic administration comprise liposomes comprising ligands for targeting the vehicle to a particular site.

Preferred methods of systemic administration, include intravenous injection, aerosol, oral and percutaneous (topical) delivery. Intravenous injections can be performed using methods standard in the art. Aerosol delivery can also be performed using methods standard in the art (see, for example, Stribling et al., Proc. Natl. Acad. Sci. USA, 189:11277-11281 (1992), which is incorporated herein by reference). Oral delivery can be performed by complexing a polynucleotide construct of the present invention to a carrier capable of withstanding degradation by digestive enzymes in the gut of an animal. Examples of such carriers, include plastic capsules or tablets, such as those known in the art. Topical delivery can be performed by mixing a polynucleotide construct of the present invention with a lipophilic reagent (e.g., DMSO) that is capable of passing into the skin.

Determining an effective amount of substance to be delivered can depend upon a number of factors including, for example, the chemical structure and biological activity of the substance, the age and weight of the animal, the precise condition requiring treatment and its severity, and the route of administration. The frequency of treatments depends upon a number of factors, such as the amount of polynucleotide constructs administered per dose, as well as the health and history of the subject. The precise amount, number of doses, and timing of doses will be determined by the attending physician or veterinarian. Therapeutic compositions of

the present invention can be administered to any animal, preferably to mammals and birds. Preferred mammals include humans, dogs, cats, mice, rats, rabbits sheep, cattle, horses and pigs, with humans being particularly

5 **Biological Activities**

The polynucleotides or polypeptides, or agonists or antagonists of the present invention can be used in assays to test for one or more biological activities. If these polynucleotides and polypeptides do exhibit activity in a particular assay, it is likely that these molecules may be involved in the diseases associated with the biological
10 activity. Thus, the polynucleotides or polypeptides, or agonists or antagonists could be used to treat the associated disease.

Immune Activity

The polynucleotides or polypeptides, or agonists or antagonists of the present
15 invention may be useful in treating deficiencies or disorders of the immune system, by activating or inhibiting the proliferation, differentiation, or mobilization (chemotaxis) of immune cells. Immune cells develop through a process called hematopoiesis, producing myeloid (platelets, red blood cells, neutrophils, and macrophages) and lymphoid (B and T lymphocytes) cells from pluripotent stem cells.
20 The etiology of these immune deficiencies or disorders may be genetic, somatic, such as cancer or some autoimmune disorders, acquired (e.g., by chemotherapy or toxins), or infectious. Moreover, a polynucleotides or polypeptides, or agonists or antagonists of the present invention can be used as a marker or detector of a particular immune system disease or disorder.

25 A polynucleotides or polypeptides, or agonists or antagonists of the present invention may be useful in treating or detecting deficiencies or disorders of hematopoietic cells. A polynucleotides or polypeptides, or agonists or antagonists of the present invention could be used to increase differentiation and proliferation of hematopoietic cells, including the pluripotent stem cells, in an effort to treat those
30 disorders associated with a decrease in certain (or many) types hematopoietic cells. Examples of immunologic deficiency syndromes include, but are not limited to:

blood protein disorders (e.g. agammaglobulinemia, dysgammaglobulinemia), ataxia
telangiectasia, common variable immunodeficiency, Digeorge Syndrome, HIV
infection, HTLV-BLV infection, leukocyte adhesion deficiency syndrome,
lymphopenia, phagocyte bactericidal dysfunction, severe combined
5 immunodeficiency (SCIDs), Wiskott-Aldrich Disorder, anemia, thrombocytopenia,
or hemoglobinuria.

Moreover, a polynucleotides or polypeptides, or agonists or antagonists of the
present invention could also be used to modulate hemostatic (the stopping of
bleeding) or thrombolytic activity (clot formation). For example, by increasing
10 hemostatic or thrombolytic activity, a polynucleotides or polypeptides, or agonists or
antagonists of the present invention could be used to treat blood coagulation disorders
(e.g., afibrinogenemia, factor deficiencies), blood platelet disorders (e.g.
thrombocytopenia), or wounds resulting from trauma, surgery, or other causes.
Alternatively, a polynucleotides or polypeptides, or agonists or antagonists of the
15 present invention that can decrease hemostatic or thrombolytic activity could be used
to inhibit or dissolve clotting. These molecules could be important in the treatment of
heart attacks (infarction), strokes, or scarring.

A polynucleotides or polypeptides, or agonists or antagonists of the present
invention may also be useful in treating or detecting autoimmune disorders. Many
20 autoimmune disorders result from inappropriate recognition of self as foreign material
by immune cells. This inappropriate recognition results in an immune response
leading to the destruction of the host tissue. Therefore, the administration of a
polynucleotides or polypeptides, or agonists or antagonists of the present invention
that inhibits an immune response, particularly the proliferation, differentiation, or
25 chemotaxis of T-cells, may be an effective therapy in preventing autoimmune
disorders.

Examples of autoimmune disorders that can be treated or detected by the
present invention include, but are not limited to: Addison's Disease, hemolytic
anemia, antiphospholipid syndrome, rheumatoid arthritis, dermatitis, allergic
30 encephalomyelitis, glomerulonephritis, Goodpasture's Syndrome, Graves' Disease,
Multiple Sclerosis, Myasthenia Gravis, Neuritis, Ophthalmia, Bullous Pemphigoid,

Pemphigus, Polyendocrinopathies, Purpura, Reiter's Disease, Stiff-Man Syndrome, Autoimmune Thyroiditis, Systemic Lupus Erythematosus, Autoimmune Pulmonary Inflammation, Guillain-Barre Syndrome, insulin dependent diabetes mellitus, and autoimmune inflammatory eye disease.

5 Similarly, allergic reactions and conditions, such as asthma (particularly allergic asthma) or other respiratory problems, may also be treated by a polynucleotides or polypeptides, or agonists or antagonists of the present invention. Moreover, these molecules can be used to treat anaphylaxis, hypersensitivity to an antigenic molecule, or blood group incompatibility.

10 A polynucleotides or polypeptides, or agonists or antagonists of the present invention may also be used to treat and/or prevent organ rejection or graft-versus-host disease (GVHD). Organ rejection occurs by host immune cell destruction of the transplanted tissue through an immune response. Similarly, an immune response is also involved in GVHD, but, in this case, the foreign transplanted immune cells
15 destroy the host tissues. The administration of a polynucleotides or polypeptides, or agonists or antagonists of the present invention that inhibits an immune response, particularly the proliferation, differentiation, or chemotaxis of T-cells, may be an effective therapy in preventing organ rejection or GVHD.

 Similarly, a polynucleotides or polypeptides, or agonists or antagonists of the
20 present invention may also be used to modulate inflammation. For example, the polypeptide or polynucleotide or agonists or antagonist may inhibit the proliferation and differentiation of cells involved in an inflammatory response. These molecules can be used to treat inflammatory conditions, both chronic and acute conditions, including inflammation associated with infection (e.g., septic shock, sepsis, or
25 systemic inflammatory response syndrome (SIRS)), ischemia-reperfusion injury, endotoxin lethality, arthritis, complement-mediated hyperacute rejection, nephritis, cytokine or chemokine induced lung injury, inflammatory bowel disease, Crohn's disease, or resulting from over production of cytokines (e.g., TNF or IL-1.)

30 **Hyperproliferative Disorders**

A polynucleotides or polypeptides, or agonists or antagonists of the invention can be used to treat or detect hyperproliferative disorders, including neoplasms. A polynucleotides or polypeptides, or agonists or antagonists of the present invention may inhibit the proliferation of the disorder through direct or indirect interactions.

- 5 Alternatively, a polynucleotides or polypeptides, or agonists or antagonists of the present invention may proliferate other cells which can inhibit the hyperproliferative disorder.

- For example, by increasing an immune response, particularly increasing antigenic qualities of the hyperproliferative disorder or by proliferating,
10 differentiating, or mobilizing T-cells, hyperproliferative disorders can be treated. This immune response may be increased by either enhancing an existing immune response, or by initiating a new immune response. Alternatively, decreasing an immune response may also be a method of treating hyperproliferative disorders, such as a chemotherapeutic agent.

- 15 Examples of hyperproliferative disorders that can be treated or detected by a polynucleotides or polypeptides, or agonists or antagonists of the present invention include, but are not limited to neoplasms located in the: abdomen, bone, breast, digestive system, liver, pancreas, peritoneum, endocrine glands (adrenal, parathyroid, pituitary, testicles, ovary, thymus, thyroid), eye, head and neck, nervous (central and
20 peripheral), lymphatic system, pelvic, skin, soft tissue, spleen, thoracic, and urogenital.

- Similarly, other hyperproliferative disorders can also be treated or detected by a polynucleotides or polypeptides, or agonists or antagonists of the present invention. Examples of such hyperproliferative disorders include, but are not limited to:
25 hypergammaglobulinemia, lymphoproliferative disorders, paraproteinemias, purpura, sarcoidosis, Sezary Syndrome, Waldenstron's Macroglobulinemia, Gaucher's Disease, histiocytosis, and any other hyperproliferative disease, besides neoplasia, located in an organ system listed above.

- One preferred embodiment utilizes polynucleotides of the present invention to
30 inhibit aberrant cellular division, by gene therapy using the present invention, and/or protein fusions or fragments thereof.

Thus, the present invention provides a method for treating cell proliferative disorders by inserting into an abnormally proliferating cell a polynucleotide of the present invention, wherein said polynucleotide represses said expression.

Another embodiment of the present invention provides a method of treating
5 cell-proliferative disorders in individuals comprising administration of one or more active gene copies of the present invention to an abnormally proliferating cell or cells. In a preferred embodiment, polynucleotides of the present invention is a DNA construct comprising a recombinant expression vector effective in expressing a DNA sequence encoding said polynucleotides. In another preferred embodiment of the
10 present invention, the DNA construct encoding the poynucleotides of the present invention is inserted into cells to be treated utilizing a retrovirus, or more preferrably an adenoviral vector (See G J. Nabel, et. al., PNAS 1999 96: 324-326, which is hereby incorporated by reference). In a most preferred embodiment, the viral vector is defective and will not transform non-proliferating cells, only proliferating cells.
15 Moreover, in a preferred embodiment, the polynucleotides of the present invention inserted into proliferating cells either alone, or in combination with or fused to other polynucleotides, can then be modulated via an external stimulus (i.e. magnetic, specific small molecule, chemical, or drug administration, etc.), which acts upon the promoter upstream of said polynucleotides to induce expression of the encoded
20 protein product. As such the beneficial therapeutic affect of the present invention may be expressly modulated (i.e. to increase, decrease, or inhibit expression of the present invention) based upon said external stimulus.

Polynucleotides of the present invention may be useful in repressing expression of oncogenic genes or antigens. By "repressing expression of the
25 oncogenic genes " is intended the suppression of the transcription of the gene, the degradation of the gene transcript (pre-message RNA), the inhibition of splicing, the destruction of the messenger RNA, the prevention of the post-translational modifications of the protein, the destruction of the protein, or the inhibition of the normal function of the protein.

30 For local administration to abnormally proliferating cells, polynucleotides of the present invention may be administered by any method known to those of skill in

the art including, but not limited to transfection, electroporation, microinjection of cells, or in vehicles such as liposomes, lipofectin, or as naked polynucleotides, or any other method described throughout the specification. The polynucleotide of the present invention may be delivered by known gene delivery systems such as, but not limited to, retroviral vectors (Gilboa, J. Virology 44:845 (1982); Hocke, Nature 320:275 (1986); Wilson, et al., Proc. Natl. Acad. Sci. U.S.A. 85:3014), vaccinia virus system (Chakrabarty et al., Mol. Cell Biol. 5:3403 (1985) or other efficient DNA delivery systems (Yates et al., Nature 313:812 (1985)) known to those skilled in the art. These references are exemplary only and are hereby incorporated by reference.

10 In order to specifically deliver or transfect cells which are abnormally proliferating and spare non-dividing cells, it is preferable to utilize a retrovirus, or adenoviral (as described in the art and elsewhere herein) delivery system known to those of skill in the art. Since host DNA replication is required for retroviral DNA to integrate and the retrovirus will be unable to self replicate due to the lack of the retrovirus genes

15 needed for its life cycle. Utilizing such a retroviral delivery system for polynucleotides of the present invention will target said gene and constructs to abnormally proliferating cells and will spare the non-dividing normal cells.

The polynucleotides of the present invention may be delivered directly to cell proliferative disorder/disease sites in internal organs, body cavities and the like by use of imaging devices used to guide an injecting needle directly to the disease site. The polynucleotides of the present invention may also be administered to disease sites at the time of surgical intervention.

20

By "cell proliferative disease" is meant any human or animal disease or disorder, affecting any one or any combination of organs, cavities, or body parts, which is characterized by single or multiple local abnormal proliferations of cells, groups of cells, or tissues, whether benign or malignant.

25

Any amount of the polynucleotides of the present invention may be administered as long as it has a biologically inhibiting effect on the proliferation of the treated cells. Moreover, it is possible to administer more than one of the polynucleotide of the present invention simultaneously to the same site. By "biologically inhibiting" is meant partial or total growth inhibition as well as

30

decreases in the rate of proliferation or growth of the cells. The biologically inhibitory dose may be determined by assessing the effects of the polynucleotides of the present invention on target malignant or abnormally proliferating cell growth in tissue culture, tumor growth in animals and cell cultures, or any other method known to one of ordinary skill in the art.

The present invention is further directed to antibody-based therapies which involve administering of anti-polypeptides and anti-polynucleotide antibodies to a mammalian, preferably human, patient for treating one or more of the described disorders. Methods for producing anti-polypeptides and anti-polynucleotide antibodies polyclonal and monoclonal antibodies are described in detail elsewhere herein. Such antibodies may be provided in pharmaceutically acceptable compositions as known in the art or as described herein.

A summary of the ways in which the antibodies of the present invention may be used therapeutically includes binding polynucleotides or polypeptides of the present invention locally or systemically in the body or by direct cytotoxicity of the antibody, e.g. as mediated by complement (CDC) or by effector cells (ADCC). Some of these approaches are described in more detail below. Armed with the teachings provided herein, one of ordinary skill in the art will know how to use the antibodies of the present invention for diagnostic, monitoring or therapeutic purposes without undue experimentation.

In particular, the antibodies, fragments and derivatives of the present invention are useful for treating a subject having or developing cell proliferative and/or differentiation disorders as described herein. Such treatment comprises administering a single or multiple doses of the antibody, or a fragment, derivative, or a conjugate thereof.

The antibodies of this invention may be advantageously utilized in combination with other monoclonal or chimeric antibodies, or with lymphokines or hematopoietic growth factors, for example, which serve to increase the number or activity of effector cells which interact with the antibodies.

It is preferred to use high affinity and/or potent in vivo inhibiting and/or neutralizing antibodies against polypeptides or polynucleotides of the present

invention, fragments or regions thereof, for both immunoassays directed to and therapy of disorders related to polynucleotides or polypeptides, including fragments thereof, of the present invention. Such antibodies, fragments, or regions, will preferably have an affinity for polynucleotides or polypeptides, including fragments thereof. Preferred binding affinities include those with a dissociation constant or K_d less than $5 \times 10^{-6}M$, $10^{-6}M$, $5 \times 10^{-7}M$, $10^{-7}M$, $5 \times 10^{-8}M$, $10^{-8}M$, $5 \times 10^{-9}M$, $10^{-9}M$, $5 \times 10^{-10}M$, $10^{-10}M$, $5 \times 10^{-11}M$, $10^{-11}M$, $5 \times 10^{-12}M$, $10^{-12}M$, $5 \times 10^{-13}M$, $10^{-13}M$, $5 \times 10^{-14}M$, $10^{-14}M$, $5 \times 10^{-15}M$, and $10^{-15}M$.

Moreover, polypeptides of the present invention are useful in inhibiting the angiogenesis of proliferative cells or tissues, either alone, as a protein fusion, or in combination with other polypeptides directly or indirectly, as described elsewhere herein. In a most preferred embodiment, said anti-angiogenesis effect may be achieved indirectly, for example, through the inhibition of hematopoietic, tumor-specific cells, such as tumor-associated macrophages (See Joseph IB, et al. J Natl Cancer Inst, 90(21):1648-53 (1998), which is hereby incorporated by reference). Antibodies directed to polypeptides or polynucleotides of the present invention may also result in inhibition of angiogenesis directly, or indirectly (See Witte L, et al., Cancer Metastasis Rev. 17(2):155-61 (1998), which is hereby incorporated by reference)).

Polypeptides, including protein fusions, of the present invention, or fragments thereof may be useful in inhibiting proliferative cells or tissues through the induction of apoptosis. Said polypeptides may act either directly, or indirectly to induce apoptosis of proliferative cells and tissues, for example in the activation of a death-domain receptor, such as tumor necrosis factor (TNF) receptor-1, CD95 (Fas/APO-1), TNF-receptor-related apoptosis-mediated protein (TRAMP) and TNF-related apoptosis-inducing ligand (TRAIL) receptor-1 and -2 (See Schulze-Osthoff K, et.al., Eur J Biochem 254(3):439-59 (1998), which is hereby incorporated by reference). Moreover, in another preferred embodiment of the present invention, said polypeptides may induce apoptosis through other mechanisms, such as in the activation of other proteins which will activate apoptosis, or through stimulating the expression of said proteins, either alone or in combination with small molecule drugs

or adjuvants, such as apoptonin, galectins, thioredoxins, antiinflammatory proteins (See for example, Mutat Res 400(1-2):447-55 (1998), Med Hypotheses.50(5):423-33 (1998), Chem Biol Interact. Apr 24;111-112:23-34 (1998), J Mol Med.76(6):402-12 (1998), Int J Tissue React;20(1):3-15 (1998), which are all hereby incorporated by
5 reference).

Polypeptides, including protein fusions to, or fragments thereof, of the present invention are useful in inhibiting the metastasis of proliferative cells or tissues. Inhibition may occur as a direct result of administering polypeptides, or antibodies directed to said polypeptides as described elsewhere herein, or indirectly, such as
10 activating the expression of proteins known to inhibit metastasis, for example alpha 4 integrins, (See, e.g., Curr Top Microbiol Immunol 1998;231:125-41, which is hereby incorporated by reference). Such therapeutic affects of the present invention may be achieved either alone, or in combination with small molecule drugs or adjuvants.

In another embodiment, the invention provides a method of delivering
15 compositions containing the polypeptides of the invention (e.g., compositions containing polypeptides or polypeptide antibodies associated with heterologous polypeptides, heterologous nucleic acids, toxins, or prodrugs) to targeted cells expressing the polypeptide of the present invention. Polypeptides or polypeptide antibodies of the invention may be associated with with heterologous polypeptides,
20 heterologous nucleic acids, toxins, or prodrugs via hydrophobic, hydrophilic, ionic and/or covalent interactions.

Polypeptides, protein fusions to, or fragments thereof, of the present invention are useful in enhancing the immunogenicity and/or antigenicity of proliferating cells or tissues, either directly, such as would occur if the polypeptides of the present
25 invention 'vaccinated' the immune response to respond to proliferative antigens and immunogens, or indirectly, such as in activating the expression of proteins known to enhance the immune response (e.g. chemokines), to said antigens and immunogens.

Cardiovascular Disorders

Polynucleotides or polypeptides, or agonists or antagonists of the invention may be used to treat cardiovascular disorders, including peripheral artery disease, such as limb ischemia.

Cardiovascular disorders include cardiovascular abnormalities, such as arterio-
5 arterial fistula, arteriovenous fistula, cerebral arteriovenous malformations, congenital heart defects, pulmonary atresia, and Scimitar Syndrome. Congenital heart defects include aortic coarctation, cor triatriatum, coronary vessel anomalies, crisscross heart, dextrocardia, patent ductus arteriosus, Ebstein's anomaly, Eisenmenger complex,
10 hypoplastic left heart syndrome, levocardia, tetralogy of fallot, transposition of great vessels, double outlet right ventricle, tricuspid atresia, persistent truncus arteriosus, and heart septal defects, such as aortopulmonary septal defect, endocardial cushion defects, Lutembacher's Syndrome, trilogy of Fallot, ventricular heart septal defects.

Cardiovascular disorders also include heart disease, such as arrhythmias, carcinoid heart disease, high cardiac output, low cardiac output, cardiac tamponade,
15 endocarditis (including bacterial), heart aneurysm, cardiac arrest, congestive heart failure, congestive cardiomyopathy, paroxysmal dyspnea, cardiac edema, heart hypertrophy, congestive cardiomyopathy, left ventricular hypertrophy, right ventricular hypertrophy, post-infarction heart rupture, ventricular septal rupture, heart valve diseases, myocardial diseases, myocardial ischemia, pericardial effusion,
20 pericarditis (including constrictive and tuberculous), pneumopericardium, postpericardiotomy syndrome, pulmonary heart disease, rheumatic heart disease, ventricular dysfunction, hyperemia, cardiovascular pregnancy complications, Scimitar Syndrome, cardiovascular syphilis, and cardiovascular tuberculosis.

Arrhythmias include sinus arrhythmia, atrial fibrillation, atrial flutter,
25 bradycardia, extrasystole, Adams-Stokes Syndrome, bundle-branch block, sinoatrial block, long QT syndrome, parasystole, Lown-Ganong-Levine Syndrome, Mahaim-type pre-excitation syndrome, Wolff-Parkinson-White syndrome, sick sinus syndrome, tachycardias, and ventricular fibrillation. Tachycardias include paroxysmal tachycardia, supraventricular tachycardia, accelerated idioventricular
30 rhythm, atrioventricular nodal reentry tachycardia, ectopic atrial tachycardia, ectopic

junctional tachycardia, sinoatrial nodal reentry tachycardia, sinus tachycardia, Torsades de Pointes, and ventricular tachycardia.

Heart valve disease include aortic valve insufficiency, aortic valve stenosis, hear murmurs, aortic valve prolapse, mitral valve prolapse, tricuspid valve prolapse, 5 mitral valve insufficiency, mitral valve stenosis, pulmonary atresia, pulmonary valve insufficiency, pulmonary valve stenosis, tricuspid atresia, tricuspid valve insufficiency, and tricuspid valve stenosis.

Myocardial diseases include alcoholic cardiomyopathy, congestive cardiomyopathy, hypertrophic cardiomyopathy, aortic subvalvular stenosis, 10 pulmonary subvalvular stenosis, restrictive cardiomyopathy, Chagas cardiomyopathy, endocardial fibroelastosis, endomyocardial fibrosis, Kearns Syndrome, myocardial reperfusion injury, and myocarditis.

Myocardial ischemias include coronary disease, such as angina pectoris, coronary aneurysm, coronary arteriosclerosis, coronary thrombosis, coronary 15 vasospasm, myocardial infarction and myocardial stunning.

Cardiovascular diseases also include vascular diseases such as aneurysms, angiodyplasia, angiomatosis, bacillary angiomatosis, Hippel-Lindau Disease, Klippel-Trenaunay-Weber Syndrome, Sturge-Weber Syndrome, angioneurotic edema, aortic diseases, Takayasu's Arteritis, aortitis, Leriche's Syndrome, arterial occlusive 20 diseases, arteritis, enarteritis, polyarteritis nodosa, cerebrovascular disorders, diabetic angiopathies, diabetic retinopathy, embolisms, thrombosis, erythromelalgia, hemorrhoids, hepatic veno-occlusive disease, hypertension, hypotension, ischemia, peripheral vascular diseases, phlebitis, pulmonary veno-occlusive disease, Raynaud's disease, CREST syndrome, retinal vein occlusion, Scimitar syndrome, superior vena 25 cava syndrome, telangiectasia, atacia telangiectasia, hereditary hemorrhagic telangiectasia, varicocele, varicose veins, varicose ulcer, vasculitis, and venous insufficiency.

Aneurysms include dissecting aneurysms, false aneurysms, infected aneurysms, ruptured aneurysms, aortic aneurysms, cerebral aneurysms, coronary 30 aneurysms, heart aneurysms, and iliac aneurysms.

Arterial occlusive diseases include arteriosclerosis, intermittent claudication, carotid stenosis, fibromuscular dysplasias, mesenteric vascular occlusion, Moyamoya disease, renal artery obstruction, retinal artery occlusion, and thromboangiitis obliterans.

- 5 Cerebrovascular disorders include carotid artery diseases, cerebral amyloid angiopathy, cerebral aneurysm, cerebral anoxia, cerebral arteriosclerosis, cerebral arteriovenous malformation, cerebral artery diseases, cerebral embolism and thrombosis, carotid artery thrombosis, sinus thrombosis, Wallenberg's syndrome, cerebral hemorrhage, epidural hematoma, subdural hematoma, subarachnoid
10 hemorrhage, cerebral infarction, cerebral ischemia (including transient), subclavian steal syndrome, periventricular leukomalacia, vascular headache, cluster headache, migraine, and vertebrobasilar insufficiency.

- Embolisms include air embolisms, amniotic fluid embolisms, cholesterol embolisms, blue toe syndrome, fat embolisms, pulmonary embolisms, and
15 thromboembolisms. Thrombosis include coronary thrombosis, hepatic vein thrombosis, retinal vein occlusion, carotid artery thrombosis, sinus thrombosis, Wallenberg's syndrome, and thrombophlebitis.

- Ischemia includes cerebral ischemia, ischemic colitis, compartment syndromes, anterior compartment syndrome, myocardial ischemia, reperfusion
20 injuries, and peripheral limb ischemia. Vasculitis includes aortitis, arteritis, Behcet's Syndrome, Churg-Strauss Syndrome, mucocutaneous lymph node syndrome, thromboangiitis obliterans, hypersensitivity vasculitis, Schoenlein-Henoch purpura, allergic cutaneous vasculitis, and Wegener's granulomatosis.

- Polynucleotides or polypeptides, or agonists or antagonists of the invention,
25 are especially effective for the treatment of critical limb ischemia and coronary disease.

- Polypeptides may be administered using any method known in the art, including, but not limited to, direct needle injection at the delivery site, intravenous injection, topical administration, catheter infusion, biolistic injectors, particle
30 accelerators, gelfoam sponge depots, other commercially available depot materials, osmotic pumps, oral or suppository solid pharmaceutical formulations, decanting or

topical applications during surgery, aerosol delivery. Such methods are known in the art. Polypeptides of the invention may be administered as part of a *Therapeutic*, described in more detail below. Methods of delivering polynucleotides of the invention are described in more detail herein.

5

Anti-Angiogenesis Activity

The naturally occurring balance between endogenous stimulators and inhibitors of angiogenesis is one in which inhibitory influences predominate. Rastinejad *et al.*, *Cell* 56:345-355 (1989). In those rare instances in which

10 neovascularization occurs under normal physiological conditions, such as wound healing, organ regeneration, embryonic development, and female reproductive processes, angiogenesis is stringently regulated and spatially and temporally delimited. Under conditions of pathological angiogenesis such as that characterizing solid tumor growth, these regulatory controls fail. Unregulated angiogenesis becomes

15 pathologic and sustains progression of many neoplastic and non-neoplastic diseases. A number of serious diseases are dominated by abnormal neovascularization including solid tumor growth and metastases, arthritis, some types of eye disorders, and psoriasis. See, e.g., reviews by Moses *et al.*, *Biotech.* 9:630-634 (1991); Folkman *et al.*, *N. Engl. J. Med.*, 333:1757-1763 (1995); Auerbach *et al.*, *J. Microvasc. Res.*

20 29:401-411 (1985); Folkman, *Advances in Cancer Research*, eds. Klein and Weinhouse, Academic Press, New York, pp. 175-203 (1985); Patz, *Am. J. Ophthalmol.* 94:715-743 (1982); and Folkman *et al.*, *Science* 221:719-725 (1983). In a number of pathological conditions, the process of angiogenesis contributes to the disease state. For example, significant data have accumulated which suggest that the

25 growth of solid tumors is dependent on angiogenesis. Folkman and Klagsbrun, *Science* 235:442-447 (1987).

The present invention provides for treatment of diseases or disorders associated with neovascularization by administration of the polynucleotides and/or polypeptides of the invention, as well as agonists or antagonists of the present

30 invention. Malignant and metastatic conditions which can be treated with the polynucleotides and polypeptides, or agonists or antagonists of the invention include,

but are not limited to, malignancies, solid tumors, and cancers described herein and otherwise known in the art (for a review of such disorders, see Fishman *et al.*, Medicine, 2d Ed., J. B. Lippincott Co., Philadelphia (1985)). Thus, the present invention provides a method of treating an angiogenesis-related disease and/or disorder, comprising administering to an individual in need thereof a therapeutically effective amount of a polynucleotide, polypeptide, antagonist and/or agonist of the invention. For example, polynucleotides, polypeptides, antagonists and/or agonists may be utilized in a variety of additional methods in order to therapeutically treat a cancer or tumor. Cancers which may be treated with polynucleotides, polypeptides, antagonists and/or agonists include, but are not limited to solid tumors, including prostate, lung, breast, ovarian, stomach, pancreas, larynx, esophagus, testes, liver, parotid, biliary tract, colon, rectum, cervix, uterus, endometrium, kidney, bladder, thyroid cancer; primary tumors and metastases; melanomas; glioblastoma; Kaposi's sarcoma; leiomyosarcoma; non-small cell lung cancer; colorectal cancer; advanced malignancies; and blood born tumors such as leukemias. For example, polynucleotides, polypeptides, antagonists and/or agonists may be delivered topically, in order to treat cancers such as skin cancer, head and neck tumors, breast tumors, and Kaposi's sarcoma.

Within yet other aspects, polynucleotides, polypeptides, antagonists and/or agonists may be utilized to treat superficial forms of bladder cancer by, for example, intravesical administration. Polynucleotides, polypeptides, antagonists and/or agonists may be delivered directly into the tumor, or near the tumor site, via injection or a catheter. Of course, as the artisan of ordinary skill will appreciate, the appropriate mode of administration will vary according to the cancer to be treated. Other modes of delivery are discussed herein.

Polynucleotides, polypeptides, antagonists and/or agonists may be useful in treating other disorders, besides cancers, which involve angiogenesis. These disorders include, but are not limited to: benign tumors, for example hemangiomas, acoustic neuromas, neurofibromas, trachomas, and pyogenic granulomas; arteriosclerotic plaques; ocular angiogenic diseases, for example, diabetic retinopathy, retinopathy of prematurity, macular degeneration, corneal graft rejection, neovascular

glaucoma, retrolental fibroplasia, rubeosis, retinoblastoma, uveitis and Pterygia (abnormal blood vessel growth) of the eye; rheumatoid arthritis; psoriasis; delayed wound healing; endometriosis; vasculogenesis; granulations; hypertrophic scars (keloids); nonunion fractures; scleroderma; trachoma; vascular adhesions; myocardial angiogenesis; coronary collaterals; cerebral collaterals; arteriovenous malformations; ischemic limb angiogenesis; Osler-Webber Syndrome; plaque neovascularization; telangiectasia; hemophilic joints; angiofibroma; fibromuscular dysplasia; wound granulation; Crohn's disease; and atherosclerosis.

For example, within one aspect of the present invention methods are provided for treating hypertrophic scars and keloids, comprising the step of administering a polynucleotide, polypeptide, antagonist and/or agonist of the invention to a hypertrophic scar or keloid.

Within one embodiment of the present invention polynucleotides, polypeptides, antagonists and/or agonists are directly injected into a hypertrophic scar or keloid, in order to prevent the progression of these lesions. This therapy is of particular value in the prophylactic treatment of conditions which are known to result in the development of hypertrophic scars and keloids (e.g., burns), and is preferably initiated after the proliferative phase has had time to progress (approximately 14 days after the initial injury), but before hypertrophic scar or keloid development. As noted above, the present invention also provides methods for treating neovascular diseases of the eye, including for example, corneal neovascularization, neovascular glaucoma, proliferative diabetic retinopathy, retrolental fibroplasia and macular degeneration.

Moreover, Ocular disorders associated with neovascularization which can be treated with the polynucleotides and polypeptides of the present invention (including agonists and/or antagonists) include, but are not limited to: neovascular glaucoma, diabetic retinopathy, retinoblastoma, retrolental fibroplasia, uveitis, retinopathy of prematurity macular degeneration, corneal graft neovascularization, as well as other eye inflammatory diseases, ocular tumors and diseases associated with choroidal or iris neovascularization. See, e.g., reviews by Waltman *et al.*, *Am. J. Ophthalmol.* 85:704-710 (1978) and Gartner *et al.*, *Surv. Ophthalmol.* 22:291-312 (1978).

Thus, within one aspect of the present invention methods are provided for

treating neovascular diseases of the eye such as corneal neovascularization (including corneal graft neovascularization), comprising the step of administering to a patient a therapeutically effective amount of a compound (as described above) to the cornea, such that the formation of blood vessels is inhibited. Briefly, the cornea is a tissue
5 which normally lacks blood vessels. In certain pathological conditions however, capillaries may extend into the cornea from the pericorneal vascular plexus of the limbus. When the cornea becomes vascularized, it also becomes clouded, resulting in a decline in the patient's visual acuity. Visual loss may become complete if the cornea completely opacitates. A wide variety of disorders can result in corneal
10 neovascularization, including for example, corneal infections (e.g., trachoma, herpes simplex keratitis, leishmaniasis and onchocerciasis), immunological processes (e.g., graft rejection and Stevens-Johnson's syndrome), alkali burns, trauma, inflammation (of any cause), toxic and nutritional deficiency states, and as a complication of wearing contact lenses.

15 Within particularly preferred embodiments of the invention, may be prepared for topical administration in saline (combined with any of the preservatives and antimicrobial agents commonly used in ocular preparations), and administered in eyedrop form. The solution or suspension may be prepared in its pure form and administered several times daily. Alternatively, anti-angiogenic compositions,
20 prepared as described above, may also be administered directly to the cornea. Within preferred embodiments, the anti-angiogenic composition is prepared with a muco-adhesive polymer which binds to cornea. Within further embodiments, the anti-angiogenic factors or anti-angiogenic compositions may be utilized as an adjunct to conventional steroid therapy. Topical therapy may also be useful prophylactically in
25 corneal lesions which are known to have a high probability of inducing an angiogenic response (such as chemical burns). In these instances the treatment, likely in combination with steroids, may be instituted immediately to help prevent subsequent complications.

30 Within other embodiments, the compounds described above may be injected directly into the corneal stroma by an ophthalmologist under microscopic guidance. The preferred site of injection may vary with the morphology of the individual lesion,

but the goal of the administration would be to place the composition at the advancing front of the vasculature (i.e., interspersed between the blood vessels and the normal cornea). In most cases this would involve perilimbic corneal injection to "protect" the cornea from the advancing blood vessels. This method may also be utilized shortly
5 after a corneal insult in order to prophylactically prevent corneal neovascularization. In this situation the material could be injected in the perilimbic cornea interspersed between the corneal lesion and its undesired potential limbic blood supply. Such methods may also be utilized in a similar fashion to prevent capillary invasion of transplanted corneas. In a sustained-release form injections might only be required 2-
10 3 times per year. A steroid could also be added to the injection solution to reduce inflammation resulting from the injection itself.

Within another aspect of the present invention, methods are provided for treating neovascular glaucoma, comprising the step of administering to a patient a therapeutically effective amount of a polynucleotide, polypeptide, antagonist and/or
15 agonist to the eye, such that the formation of blood vessels is inhibited. In one embodiment, the compound may be administered topically to the eye in order to treat early forms of neovascular glaucoma. Within other embodiments, the compound may be implanted by injection into the region of the anterior chamber angle. Within other embodiments, the compound may also be placed in any location such that the
20 compound is continuously released into the aqueous humor. Within another aspect of the present invention, methods are provided for treating proliferative diabetic retinopathy, comprising the step of administering to a patient a therapeutically effective amount of a polynucleotide, polypeptide, antagonist and/or agonist to the eyes, such that the formation of blood vessels is inhibited.

25 Within particularly preferred embodiments of the invention, proliferative diabetic retinopathy may be treated by injection into the aqueous humor or the vitreous, in order to increase the local concentration of the polynucleotide, polypeptide, antagonist and/or agonist in the retina. Preferably, this treatment should be initiated prior to the acquisition of severe disease requiring photocoagulation.

30 Within another aspect of the present invention, methods are provided for treating retrolental fibroplasia, comprising the step of administering to a patient a

therapeutically effective amount of a polynucleotide, polypeptide, antagonist and/or agonist to the eye, such that the formation of blood vessels is inhibited. The compound may be administered topically, via intravitreal injection and/or via intraocular implants.

5 Additionally, disorders which can be treated with the polynucleotides, polypeptides, agonists and/or antagonists include, but are not limited to, hemangioma, arthritis, psoriasis, angiofibroma, atherosclerotic plaques, delayed wound healing, granulations, hemophilic joints, hypertrophic scars, nonunion fractures, Osler-Weber syndrome, pyogenic granuloma, scleroderma, trachoma, and vascular adhesions.

10 Moreover, disorders and/or states, which can be treated with the polynucleotides, polypeptides, agonists and/or antagonists include, but are not limited to, solid tumors, blood born tumors such as leukemias, tumor metastasis, Kaposi's sarcoma, benign tumors, for example hemangiomas, acoustic neuromas, neurofibromas, trachomas, and pyogenic granulomas, rheumatoid arthritis, psoriasis,
15 ocular angiogenic diseases, for example, diabetic retinopathy, retinopathy of prematurity, macular degeneration, corneal graft rejection, neovascular glaucoma, retrolental fibroplasia, rubeosis, retinoblastoma, and uveitis, delayed wound healing, endometriosis, vasculogenesis, granulations, hypertrophic scars (keloids), nonunion fractures, scleroderma, trachoma, vascular adhesions, myocardial angiogenesis,
20 coronary collaterals, cerebral collaterals, arteriovenous malformations, ischemic limb angiogenesis, Osler-Webber Syndrome, plaque neovascularization, telangiectasia, hemophilic joints, angiofibroma fibromuscular dysplasia, wound granulation, Crohn's disease, atherosclerosis, birth control agent by preventing vascularization required for embryo implantation controlling menstruation, diseases that have
25 angiogenesis as a pathologic consequence such as cat scratch disease (Rochelominalia quintosa), ulcers (*Helicobacter pylori*), Bartonellosis and bacillary angiomatosis.

 In one aspect of the birth control method, an amount of the compound sufficient to block embryo implantation is administered before or after intercourse and
30 fertilization have occurred, thus providing an effective method of birth control, possibly a "morning after" method. Polynucleotides, polypeptides, agonists and/or

agonists may also be used in controlling menstruation or administered as either a peritoneal lavage fluid or for peritoneal implantation in the treatment of endometriosis.

Polynucleotides, polypeptides, agonists and/or agonists of the present invention may be incorporated into surgical sutures in order to prevent stitch granulomas.

Polynucleotides, polypeptides, agonists and/or agonists may be utilized in a wide variety of surgical procedures. For example, within one aspect of the present invention a compositions (in the form of, for example, a spray or film) may be utilized to coat or spray an area prior to removal of a tumor, in order to isolate normal surrounding tissues from malignant tissue, and/or to prevent the spread of disease to surrounding tissues. Within other aspects of the present invention, compositions (e.g., in the form of a spray) may be delivered via endoscopic procedures in order to coat tumors, or inhibit angiogenesis in a desired locale. Within yet other aspects of the present invention, surgical meshes which have been coated with anti- angiogenic compositions of the present invention may be utilized in any procedure wherein a surgical mesh might be utilized. For example, within one embodiment of the invention a surgical mesh laden with an anti-angiogenic composition may be utilized during abdominal cancer resection surgery (e.g., subsequent to colon resection) in order to provide support to the structure, and to release an amount of the anti-angiogenic factor.

Within further aspects of the present invention, methods are provided for treating tumor excision sites, comprising administering a polynucleotide, polypeptide, agonist and/or agonist to the resection margins of a tumor subsequent to excision, such that the local recurrence of cancer and the formation of new blood vessels at the site is inhibited. Within one embodiment of the invention, the anti-angiogenic compound is administered directly to the tumor excision site (e.g., applied by swabbing, brushing or otherwise coating the resection margins of the tumor with the anti-angiogenic compound). Alternatively, the anti-angiogenic compounds may be incorporated into known surgical pastes prior to administration. Within particularly

preferred embodiments of the invention, the anti-angiogenic compounds are applied after hepatic resections for malignancy, and after neurosurgical operations.

Within one aspect of the present invention, polynucleotides, polypeptides, agonists and/or agonists may be administered to the resection margin of a wide
5 variety of tumors, including for example, breast, colon, brain and hepatic tumors. For example, within one embodiment of the invention, anti-angiogenic compounds may be administered to the site of a neurological tumor subsequent to excision, such that the formation of new blood vessels at the site are inhibited.

The polynucleotides, polypeptides, agonists and/or agonists of the present
10 invention may also be administered along with other anti-angiogenic factors. Representative examples of other anti-angiogenic factors include: Anti-Invasive Factor, retinoic acid and derivatives thereof, paclitaxel, Suramin, Tissue Inhibitor of Metalloproteinase-1, Tissue Inhibitor of Metalloproteinase-2, Plasminogen Activator Inhibitor-1, Plasminogen Activator Inhibitor-2, and various forms of the lighter "d
15 group" transition metals.

Lighter "d group" transition metals include, for example, vanadium, molybdenum, tungsten, titanium, niobium, and tantalum species. Such transition metal species may form transition metal complexes. Suitable complexes of the above-mentioned transition metal species include oxo transition metal complexes.

Representative examples of vanadium complexes include oxo vanadium
20 complexes such as vanadate and vanadyl complexes. Suitable vanadate complexes include metavanadate and orthovanadate complexes such as, for example, ammonium metavanadate, sodium metavanadate, and sodium orthovanadate. Suitable vanadyl complexes include, for example, vanadyl acetylacetonate and vanadyl sulfate
25 including vanadyl sulfate hydrates such as vanadyl sulfate mono- and trihydrates.

Representative examples of tungsten and molybdenum complexes also include oxo complexes. Suitable oxo tungsten complexes include tungstate and tungsten oxide complexes. Suitable tungstate complexes include ammonium tungstate, calcium tungstate, sodium tungstate dihydrate, and tungstic acid. Suitable tungsten
30 oxides include tungsten (IV) oxide and tungsten (VI) oxide. Suitable oxo molybdenum complexes include molybdate, molybdenum oxide, and molybdenyl

complexes. Suitable molybdate complexes include ammonium molybdate and its hydrates, sodium molybdate and its hydrates, and potassium molybdate and its hydrates. Suitable molybdenum oxides include molybdenum (VI) oxide, molybdenum (VI) oxide, and molybdic acid. Suitable molybdenyl complexes include, for example, molybdenyl acetylacetonate. Other suitable tungsten and molybdenum complexes include hydroxo derivatives derived from, for example, glycerol, tartaric acid, and sugars.

A wide variety of other anti-angiogenic factors may also be utilized within the context of the present invention. Representative examples include platelet factor 4; protamine sulphate; sulphated chitin derivatives (prepared from queen crab shells), (Murata et al., Cancer Res. 51:22-26, 1991); Sulphated Polysaccharide Peptidoglycan Complex (SP- PG) (the function of this compound may be enhanced by the presence of steroids such as estrogen, and tamoxifen citrate); Staurosporine; modulators of matrix metabolism, including for example, proline analogs, cishydroxyproline, d,L-3,4-dehydroproline, Thiaproline, alpha,alpha-dipyridyl, aminopropionitrile fumarate; 4-propyl-5-(4-pyridinyl)-2(3H)-oxazolone; Methotrexate; Mitoxantrone; Heparin; Interferons; 2 Macroglobulin-serum; ChIMP-3 (Pavloff et al., J. Bio. Chem. 267:17321-17326, 1992); Chymostatin (Tomkinson et al., Biochem J. 286:475-480, 1992); Cyclodextrin Tetradasulfate; Eponemycin; Camptothecin; Fumagillin (Ingber et al., Nature 348:555-557, 1990); Gold Sodium Thiomalate ("GST"; Matsubara and Ziff, J. Clin. Invest. 79:1440-1446, 1987); anticollagenase-serum; alpha2-antiplasmin (Holmes et al., J. Biol. Chem. 262(4):1659-1664, 1987); Bisantrene (National Cancer Institute); Lobenzarit disodium (N-(2)-carboxyphenyl-4-chloroanthronilic acid disodium or "CCA"; Takeuchi et al., Agents Actions 36:312-316, 1992); Thalidomide; Angostatic steroid; AGM-1470; carboxynaminolmidazole; and metalloproteinase inhibitors such as BB94.

Diseases at the Cellular Level

Diseases associated with increased cell survival or the inhibition of apoptosis that could be treated or detected by the polynucleotides or polypeptides and/or antagonists or agonists of the invention, include cancers (such as follicular

lymphomas, carcinomas with p53 mutations, and hormone-dependent tumors, including, but not limited to colon cancer, cardiac tumors, pancreatic cancer, melanoma, retinoblastoma, glioblastoma, lung cancer, intestinal cancer, testicular cancer, stomach cancer, neuroblastoma, myxoma, myoma, lymphoma, endothelioma, osteoblastoma, osteoclastoma, osteosarcoma, chondrosarcoma, adenoma, breast cancer, prostate cancer, Kaposi's sarcoma and ovarian cancer); autoimmune disorders (such as, multiple sclerosis, Sjogren's syndrome, Hashimoto's thyroiditis, biliary cirrhosis, Behcet's disease, Crohn's disease, polymyositis, systemic lupus erythematosus and immune-related glomerulonephritis and rheumatoid arthritis) and viral infections (such as herpes viruses, pox viruses and adenoviruses), inflammation, graft v. host disease, acute graft rejection, and chronic graft rejection. In preferred embodiments, the polynucleotides or polypeptides, and/or agonists or antagonists of the invention are used to inhibit growth, progression, and/or metasis of cancers, in particular those listed above.

Additional diseases or conditions associated with increased cell survival that could be treated or detected by the polynucleotides or polypeptides, or agonists or antagonists of the invention, include, but are not limited to, progression, and/or metastases of malignancies and related disorders such as leukemia (including acute leukemias (e.g., acute lymphocytic leukemia, acute myelocytic leukemia (including myeloblastic, promyelocytic, myelomonocytic, monocytic, and erythroleukemia)) and chronic leukemias (e.g., chronic myelocytic (granulocytic) leukemia and chronic lymphocytic leukemia)), polycythemia vera, lymphomas (e.g., Hodgkin's disease and non-Hodgkin's disease), multiple myeloma, Waldenstrom's macroglobulinemia, heavy chain disease, and solid tumors including, but not limited to, sarcomas and carcinomas such as fibrosarcoma, myxosarcoma, liposarcoma, chondrosarcoma, osteogenic sarcoma, chordoma, angiosarcoma, endotheliosarcoma, lymphangiosarcoma, lymphangioendotheliosarcoma, synovioma, mesothelioma, Ewing's tumor, leiomyosarcoma, rhabdomyosarcoma, colon carcinoma, pancreatic cancer, breast cancer, ovarian cancer, prostate cancer, squamous cell carcinoma, basal cell carcinoma, adenocarcinoma, sweat gland carcinoma, sebaceous gland carcinoma, papillary carcinoma, papillary adenocarcinomas, cystadenocarcinoma, medullary

carcinoma, bronchogenic carcinoma, renal cell carcinoma, hepatoma, bile duct carcinoma, choriocarcinoma, seminoma, embryonal carcinoma, Wilm's tumor, cervical cancer, testicular tumor, lung carcinoma, small cell lung carcinoma, bladder carcinoma, epithelial carcinoma, glioma, astrocytoma, medulloblastoma, 5 craniopharyngioma, ependymoma, pinealoma, hemangioblastoma, acoustic neuroma, oligodendroglioma, meningioma, melanoma, neuroblastoma, and retinoblastoma.

Diseases associated with increased apoptosis that could be treated or detected by the polynucleotides or polypeptides, and/or agonists or antagonists of the invention, include AIDS; neurodegenerative disorders (such as Alzheimer's disease, 10 Parkinson's disease, Amyotrophic lateral sclerosis, Retinitis pigmentosa, Cerebellar degeneration and brain tumor or prior associated disease); autoimmune disorders (such as, multiple sclerosis, Sjogren's syndrome, Hashimoto's thyroiditis, biliary cirrhosis, Behcet's disease, Crohn's disease, polymyositis, systemic lupus erythematosus and immune-related glomerulonephritis and rheumatoid arthritis) 15 myelodysplastic syndromes (such as aplastic anemia), graft v. host disease, ischemic injury (such as that caused by myocardial infarction, stroke and reperfusion injury), liver injury (e.g., hepatitis related liver injury, ischemia/reperfusion injury, cholestasis (bile duct injury) and liver cancer); toxin-induced liver disease (such as that caused by alcohol), septic shock, cachexia and anorexia.

20

Wound Healing and Epithelial Cell Proliferation

In accordance with yet a further aspect of the present invention, there is provided a process for utilizing the polynucleotides or polypeptides, and/or agonists or antagonists of the invention, for therapeutic purposes, for example, to stimulate 25 epithelial cell proliferation and basal keratinocytes for the purpose of wound healing, and to stimulate hair follicle production and healing of dermal wounds.

Polynucleotides or polypeptides, as well as agonists or antagonists of the invention, may be clinically useful in stimulating wound healing including surgical wounds, excisional wounds, deep wounds involving damage of the dermis and epidermis, eye 30 tissue wounds, dental tissue wounds, oral cavity wounds, diabetic ulcers, dermal ulcers, cubitus ulcers, arterial ulcers, venous stasis ulcers, burns resulting from heat

exposure or chemicals, and other abnormal wound healing conditions such as uremia, malnutrition, vitamin deficiencies and complications associated with systemic treatment with steroids, radiation therapy and antineoplastic drugs and antimetabolites. Polynucleotides or polypeptides, and/or agonists or antagonists of the invention, could be used to promote dermal reestablishment subsequent to dermal loss

The polynucleotides or polypeptides, and/or agonists or antagonists of the invention, could be used to increase the adherence of skin grafts to a wound bed and to stimulate re-epithelialization from the wound bed. The following are a non-exhaustive list of grafts that polynucleotides or polypeptides, agonists or antagonists of the invention, could be used to increase adherence to a wound bed: autografts, artificial skin, allografts, autodermic graft, autoepidermic grafts, avacular grafts, Blair-Brown grafts, bone graft, brephoplastic grafts, cutis graft, delayed graft, dermic graft, epidermic graft, fascia graft, full thickness graft, heterologous graft, xenograft, homologous graft, hyperplastic graft, lamellar graft, mesh graft, mucosal graft, Ollier-Thiersch graft, omentopial graft, patch graft, pedicle graft, penetrating graft, split skin graft, thick split graft. The polynucleotides or polypeptides, and/or agonists or antagonists of the invention, can be used to promote skin strength and to improve the appearance of aged skin.

It is believed that the polynucleotides or polypeptides, and/or agonists or antagonists of the invention, will also produce changes in hepatocyte proliferation, and epithelial cell proliferation in the lung, breast, pancreas, stomach, small intestine, and large intestine. The polynucleotides or polypeptides, and/or agonists or antagonists of the invention, could promote proliferation of epithelial cells such as sebocytes, hair follicles, hepatocytes, type II pneumocytes, mucin-producing goblet cells, and other epithelial cells and their progenitors contained within the skin, lung, liver, and gastrointestinal tract. The polynucleotides or polypeptides, and/or agonists or antagonists of the invention, may promote proliferation of endothelial cells, keratinocytes, and basal keratinocytes.

The polynucleotides or polypeptides, and/or agonists or antagonists of the invention, could also be used to reduce the side effects of gut toxicity that result from

radiation, chemotherapy treatments or viral infections. The polynucleotides or polypeptides, and/or agonists or antagonists of the invention, may have a cytoprotective effect on the small intestine mucosa. The polynucleotides or polypeptides, and/or agonists or antagonists of the invention, may also stimulate
5 healing of mucositis (mouth ulcers) that result from chemotherapy and viral infections.

The polynucleotides or polypeptides, and/or agonists or antagonists of the invention, could further be used in full regeneration of skin in full and partial thickness skin defects, including burns, (i.e., repopulation of hair follicles, sweat
10 glands, and sebaceous glands), treatment of other skin defects such as psoriasis. The polynucleotides or polypeptides, and/or agonists or antagonists of the invention, could be used to treat epidermolysis bullosa, a defect in adherence of the epidermis to the underlying dermis which results in frequent, open and painful blisters by accelerating reepithelialization of these lesions. The polynucleotides or polypeptides, and/or
15 agonists or antagonists of the invention, could also be used to treat gastric and duodenal ulcers and help heal by scar formation of the mucosal lining and regeneration of glandular mucosa and duodenal mucosal lining more rapidly. Inflammatory bowel diseases, such as Crohn's disease and ulcerative colitis, are diseases which result in destruction of the mucosal surface of the small or large
20 intestine, respectively. Thus, the polynucleotides or polypeptides, and/or agonists or antagonists of the invention, could be used to promote the resurfacing of the mucosal surface to aid more rapid healing and to prevent progression of inflammatory bowel disease. Treatment with the polynucleotides or polypeptides, and/or agonists or antagonists of the invention, is expected to have a significant effect on the production
25 of mucus throughout the gastrointestinal tract and could be used to protect the intestinal mucosa from injurious substances that are ingested or following surgery. The polynucleotides or polypeptides, and/or agonists or antagonists of the invention, could be used to treat diseases associate with the under expression of the polynucleotides of the invention.

30 Moreover, the polynucleotides or polypeptides, and/or agonists or antagonists of the invention, could be used to prevent and heal damage to the lungs due to various

pathological states. A growth factor such as the polynucleotides or polypeptides, and/or agonists or antagonists of the invention, which could stimulate proliferation and differentiation and promote the repair of alveoli and bronchiolar epithelium to prevent or treat acute or chronic lung damage. For example, emphysema, which
5 results in the progressive loss of aveoli, and inhalation injuries, i.e., resulting from smoke inhalation and burns, that cause necrosis of the bronchiolar epithelium and alveoli could be effectively treated using the polynucleotides or polypeptides, and/or agonists or antagonists of the invention. Also, the polynucleotides or polypeptides, and/or agonists or antagonists of the invention, could be used to stimulate the
10 proliferation of and differentiation of type II pneumocytes, which may help treat or prevent disease such as hyaline membrane diseases, such as infant respiratory distress syndrome and bronchopulmonary displasia, in premature infants.

The polynucleotides or polypeptides, and/or agonists or antagonists of the invention, could stimulate the proliferation and differentiation of hepatocytes and,
15 thus, could be used to alleviate or treat liver diseases and pathologies such as fulminant liver failure caused by cirrhosis, liver damage caused by viral hepatitis and toxic substances (i.e., acetaminophen, carbon tetraholoride and other hepatotoxins known in the art).

In addition, the polynucleotides or polypeptides, and/or agonists or antagonists
20 of the invention, could be used treat or prevent the onset of diabetes mellitus. In patients with newly diagnosed Types I and II diabetes, where some islet cell function remains, the polynucleotides or polypeptides, and/or agonists or antagonists of the invention, could be used to maintain the islet function so as to alleviate, delay or prevent permanent manifestation of the disease. Also, the polynucleotides or
25 polypeptides, and/or agonists or antagonists of the invention, could be used as an auxiliary in islet cell transplantation to improve or promote islet cell function.

Neurological Diseases

Nervous system disorders, which can be treated with the compositions of the
30 invention (e.g., polypeptides, polynucleotides, and/or agonists or antagonists), include, but are not limited to, nervous system injuries, and diseases or disorders

which result in either a disconnection of axons, a diminution or degeneration of neurons, or demyelination. Nervous system lesions which may be treated in a patient (including human and non-human mammalian patients) according to the invention, include but are not limited to, the following lesions of either the central (including spinal cord, brain) or peripheral nervous systems: (1) ischemic lesions, in which a lack of oxygen in a portion of the nervous system results in neuronal injury or death, including cerebral infarction or ischemia, or spinal cord infarction or ischemia; (2) traumatic lesions, including lesions caused by physical injury or associated with surgery, for example, lesions which sever a portion of the nervous system, or compression injuries; (3) malignant lesions, in which a portion of the nervous system is destroyed or injured by malignant tissue which is either a nervous system associated malignancy or a malignancy derived from non-nervous system tissue; (4) infectious lesions, in which a portion of the nervous system is destroyed or injured as a result of infection, for example, by an abscess or associated with infection by human immunodeficiency virus, herpes zoster, or herpes simplex virus or with Lyme disease, tuberculosis, syphilis; (5) degenerative lesions, in which a portion of the nervous system is destroyed or injured as a result of a degenerative process including but not limited to degeneration associated with Parkinson's disease, Alzheimer's disease, Huntington's chorea, or amyotrophic lateral sclerosis (ALS); (6) lesions associated with nutritional diseases or disorders, in which a portion of the nervous system is destroyed or injured by a nutritional disorder or disorder of metabolism including but not limited to, vitamin B12 deficiency, folic acid deficiency, Wernicke disease, tobacco-alcohol amblyopia, Marchiafava-Bignami disease (primary degeneration of the corpus callosum), and alcoholic cerebellar degeneration; (7) neurological lesions associated with systemic diseases including, but not limited to, diabetes (diabetic neuropathy, Bell's palsy), systemic lupus erythematosus, carcinoma, or sarcoidosis; (8) lesions caused by toxic substances including alcohol, lead, or particular neurotoxins; and (9) demyelinated lesions in which a portion of the nervous system is destroyed or injured by a demyelinating disease including, but not limited to, multiple sclerosis, human immunodeficiency virus-associated myelopathy, transverse

myelopathy or various etiologies, progressive multifocal leukoencephalopathy, and central pontine myelinolysis.

In a preferred embodiment, the polypeptides, polynucleotides, or agonists or antagonists of the invention are used to protect neural cells from the damaging effects of cerebral hypoxia. According to this embodiment, the compositions of the invention are used to treat or prevent neural cell injury associated with cerebral hypoxia. In one aspect of this embodiment, the polypeptides, polynucleotides, or agonists or antagonists of the invention are used to treat or prevent neural cell injury associated with cerebral ischemia. In another aspect of this embodiment, the polypeptides, polynucleotides, or agonists or antagonists of the invention are used to treat or prevent neural cell injury associated with cerebral infarction. In another aspect of this embodiment, the polypeptides, polynucleotides, or agonists or antagonists of the invention are used to treat or prevent neural cell injury associated with a stroke. In a further aspect of this embodiment, the polypeptides, polynucleotides, or agonists or antagonists of the invention are used to treat or prevent neural cell injury associated with a heart attack.

The compositions of the invention which are useful for treating or preventing a nervous system disorder may be selected by testing for biological activity in promoting the survival or differentiation of neurons. For example, and not by way of limitation, compositions of the invention which elicit any of the following effects may be useful according to the invention: (1) increased survival time of neurons in culture; (2) increased sprouting of neurons in culture or *in vivo*; (3) increased production of a neuron-associated molecule in culture or *in vivo*, *e.g.*, choline acetyltransferase or acetylcholinesterase with respect to motor neurons; or (4) decreased symptoms of neuron dysfunction *in vivo*. Such effects may be measured by any method known in the art. In preferred, non-limiting embodiments, increased survival of neurons may routinely be measured using a method set forth herein or otherwise known in the art, such as, for example, the method set forth in Arakawa et al. (J. Neurosci. 10:3507-3515 (1990)); increased sprouting of neurons may be detected by methods known in the art, such as, for example, the methods set forth in Pestronk et al. (Exp. Neurol. 70:65-82 (1980)) or Brown et al. (Ann. Rev. Neurosci. 4:17-42 (1981));

increased production of neuron-associated molecules may be measured by bioassay, enzymatic assay, antibody binding, Northern blot assay, etc., using techniques known in the art and depending on the molecule to be measured; and motor neuron dysfunction may be measured by assessing the physical manifestation of motor neuron disorder, e.g., weakness, motor neuron conduction velocity, or functional disability.

In specific embodiments, motor neuron disorders that may be treated according to the invention include, but are not limited to, disorders such as infarction, infection, exposure to toxin, trauma, surgical damage, degenerative disease or malignancy that may affect motor neurons as well as other components of the nervous system, as well as disorders that selectively affect neurons such as amyotrophic lateral sclerosis, and including, but not limited to, progressive spinal muscular atrophy, progressive bulbar palsy, primary lateral sclerosis, infantile and juvenile muscular atrophy, progressive bulbar paralysis of childhood (Fazio-Londe syndrome), poliomyelitis and the post polio syndrome, and Hereditary Motorsensory Neuropathy (Charcot-Marie-Tooth Disease).

Infectious Disease

A polypeptide or polynucleotide and/or agonist or antagonist of the present invention can be used to treat or detect infectious agents. For example, by increasing the immune response, particularly increasing the proliferation and differentiation of B and/or T cells, infectious diseases may be treated. The immune response may be increased by either enhancing an existing immune response, or by initiating a new immune response. Alternatively, polypeptide or polynucleotide and/or agonist or antagonist of the present invention may also directly inhibit the infectious agent, without necessarily eliciting an immune response.

Viruses are one example of an infectious agent that can cause disease or symptoms that can be treated or detected by a polynucleotide or polypeptide and/or agonist or antagonist of the present invention. Examples of viruses, include, but are not limited to Examples of viruses, include, but are not limited to the following DNA and RNA viruses and viral families: Arbovirus, Adenoviridae, Arenaviridae,

- Arterivirus, Birnaviridae, Bunyaviridae, Caliciviridae, Circoviridae, Coronaviridae, Dengue, EBV, HIV, Flaviviridae, Hepadnaviridae (Hepatitis), Herpesviridae (such as, Cytomegalovirus, Herpes Simplex, Herpes Zoster), Mononegavirus (e.g., Paramyxoviridae, Morbillivirus, Rhabdoviridae), Orthomyxoviridae (e.g., Influenza A, Influenza B, and parainfluenza), Papiloma virus, Papovaviridae, Parvoviridae, Picornaviridae, Poxviridae (such as Smallpox or Vaccinia), Reoviridae (e.g., Rotavirus), Retroviridae (HTLV-I, HTLV-II, Lentivirus), and Togaviridae (e.g., Rubivirus). Viruses falling within these families can cause a variety of diseases or symptoms, including, but not limited to: arthritis, bronchiolitis, respiratory syncytial virus, encephalitis, eye infections (e.g., conjunctivitis, keratitis), chronic fatigue syndrome, hepatitis (A, B, C, E, Chronic Active, Delta), Japanese B encephalitis, Junin, Chikungunya, Rift Valley fever, yellow fever, meningitis, opportunistic infections (e.g., AIDS), pneumonia, Burkitt's Lymphoma, chickenpox, hemorrhagic fever, Measles, Mumps, Parainfluenza, Rabies, the common cold, Polio, leukemia, Rubella, sexually transmitted diseases, skin diseases (e.g., Kaposi's, warts), and viremia. polynucleotides or polypeptides, or agonists or antagonists of the invention, can be used to treat or detect any of these symptoms or diseases. In specific embodiments, polynucleotides, polypeptides, or agonists or antagonists of the invention are used to treat: meningitis, Dengue, EBV, and/or hepatitis (e.g., hepatitis B). In an additional specific embodiment polynucleotides, polypeptides, or agonists or antagonists of the invention are used to treat patients nonresponsive to one or more other commercially available hepatitis vaccines. In a further specific embodiment polynucleotides, polypeptides, or agonists or antagonists of the invention are used to treat AIDS.
- 25 Similarly, bacterial or fungal agents that can cause disease or symptoms and that can be treated or detected by a polynucleotide or polypeptide and/or agonist or antagonist of the present invention include, but not limited to, include, but not limited to, the following Gram-Negative and Gram-positive bacteria and bacterial families and fungi: Actinomycetales (e.g., Corynebacterium, Mycobacterium, Norcardia), Cryptococcus neoformans, Aspergillosis, Bacillaceae (e.g., Anthrax, Clostridium), Bacteroidaceae, Blastomycosis, Bordetella, Borrelia (e.g., Borrelia burgdorferi),

- Brucellosis, Candidiasis, Campylobacter, Coccidioidomycosis, Cryptococcosis, Dermatocycoses, E. coli (e.g., Enterotoxigenic E. coli and Enterohemorrhagic E. coli), Enterobacteriaceae (Klebsiella, Salmonella (e.g., Salmonella typhi, and Salmonella paratyphi), Serratia, Yersinia), Erysipelothrix, Helicobacter,
- 5 Legionellosis, Leptospirosis, Listeria, Mycoplasmatales, Mycobacterium leprae, Vibrio cholerae, Neisseriaceae (e.g., Acinetobacter, Gonorrhea, Meningococcal), Meisseria meningitidis, Pasteurellacea Infections (e.g., Actinobacillus, Haemophilus (e.g., Haemophilus influenza type B), Pasteurella), Pseudomonas, Rickettsiaceae, Chlamydiaceae, Syphilis, Shigella spp., Staphylococcal, Meningococcal,
- 10 Pneumococcal and Streptococcal (e.g., Streptococcus pneumoniae and Group B Streptococcus). These bacterial or fungal families can cause the following diseases or symptoms, including, but not limited to: bacteremia, endocarditis, eye infections (conjunctivitis, tuberculosis, uveitis), gingivitis, opportunistic infections (e.g., AIDS related infections), paronychia, prosthesis-related infections, Reiter's Disease,
- 15 respiratory tract infections, such as Whooping Cough or Empyema, sepsis, Lyme Disease, Cat-Scratch Disease, Dysentery, Paratyphoid Fever, food poisoning, Typhoid, pneumonia, Gonorrhea, meningitis (e.g., meningitis types A and B), Chlamydia, Syphilis, Diphtheria, Leprosy, Paratuberculosis, Tuberculosis, Lupus, Botulism, gangrene, tetanus, impetigo, Rheumatic Fever, Scarlet Fever, sexually
- 20 transmitted diseases, skin diseases (e.g., cellulitis, dermatocycoses), toxemia, urinary tract infections, wound infections. Polynucleotides or polypeptides, agonists or antagonists of the invention, can be used to treat or detect any of these symptoms or diseases. In specific embodiments, Polynucleotides, polypeptides, agonists or antagonists of the invention are used to treat: tetanus, Diphtheria, botulism, and/or
- 25 meningitis type B.

Moreover, parasitic agents causing disease or symptoms that can be treated or detected by a polynucleotide or polypeptide and/or agonist or antagonist of the present invention include, but not limited to, the following families or class:

Amebiasis, Babesiosis, Coccidiosis, Cryptosporidiosis, Dientamoebiasis, Dourine,

30 Ectoparasitic, Giardiasis, Helminthiasis, Leishmaniasis, Theileriasis, Toxoplasmosis, Trypanosomiasis, and Trichomonas and Sporozoans (e.g., Plasmodium virax,

Plasmodium falciparum, Plasmodium malariae and Plasmodium ovale). These parasites can cause a variety of diseases or symptoms, including, but not limited to: Scabies, Trombiculiasis, eye infections, intestinal disease (e.g., dysentery, giardiasis), liver disease, lung disease, opportunistic infections (e.g., AIDS related), malaria, pregnancy complications, and toxoplasmosis. polynucleotides or polypeptides, or agonists or antagonists of the invention, can be used to treat or detect any of these symptoms or diseases. In specific embodiments, polynucleotides, polypeptides, or agonists or antagonists of the invention are used to treat malaria.

Preferably, treatment using a polypeptide or polynucleotide and/or agonist or antagonist of the present invention could either be by administering an effective amount of a polypeptide to the patient, or by removing cells from the patient, supplying the cells with a polynucleotide of the present invention, and returning the engineered cells to the patient (ex vivo therapy). Moreover, the polypeptide or polynucleotide of the present invention can be used as an antigen in a vaccine to raise an immune response against infectious disease.

Regeneration

A polynucleotide or polypeptide and/or agonist or antagonist of the present invention can be used to differentiate, proliferate, and attract cells, leading to the regeneration of tissues. (See, Science 276:59-87 (1997).) The regeneration of tissues could be used to repair, replace, or protect tissue damaged by congenital defects, trauma (wounds, burns, incisions, or ulcers), age, disease (e.g. osteoporosis, osteoarthritis, periodontal disease, liver failure), surgery, including cosmetic plastic surgery, fibrosis, reperfusion injury, or systemic cytokine damage.

Tissues that could be regenerated using the present invention include organs (e.g., pancreas, liver, intestine, kidney, skin, endothelium), muscle (smooth, skeletal or cardiac), vasculature (including vascular and lymphatics), nervous, hematopoietic, and skeletal (bone, cartilage, tendon, and ligament) tissue. Preferably, regeneration occurs without or decreased scarring. Regeneration also may include angiogenesis.

Moreover, a polynucleotide or polypeptide and/or agonist or antagonist of the present invention may increase regeneration of tissues difficult to heal. For example,

increased tendon/ligament regeneration would quicken recovery time after damage. A polynucleotide or polypeptide and/or agonist or antagonist of the present invention could also be used prophylactically in an effort to avoid damage. Specific diseases that could be treated include of tendinitis, carpal tunnel syndrome, and other tendon or
5 ligament defects. A further example of tissue regeneration of non-healing wounds includes pressure ulcers, ulcers associated with vascular insufficiency, surgical, and traumatic wounds.

Similarly, nerve and brain tissue could also be regenerated by using a polynucleotide or polypeptide and/or agonist or antagonist of the present invention to
10 proliferate and differentiate nerve cells. Diseases that could be treated using this method include central and peripheral nervous system diseases, neuropathies, or mechanical and traumatic disorders (e.g., spinal cord disorders, head trauma, cerebrovascular disease, and stroke). Specifically, diseases associated with peripheral nerve injuries, peripheral neuropathy (e.g., resulting from chemotherapy or other
15 medical therapies), localized neuropathies, and central nervous system diseases (e.g., Alzheimer's disease, Parkinson's disease, Huntington's disease, amyotrophic lateral sclerosis, and Shy-Drager syndrome), could all be treated using the polynucleotide or polypeptide and/or agonist or antagonist of the present invention.

20 **Chemotaxis**

A polynucleotide or polypeptide and/or agonist or antagonist of the present invention may have chemotaxis activity. A chemotactic molecule attracts or mobilizes cells (e.g., monocytes, fibroblasts, neutrophils, T-cells, mast cells, eosinophils, epithelial and/or endothelial cells) to a particular site in the body, such as
25 inflammation, infection, or site of hyperproliferation. The mobilized cells can then fight off and/or heal the particular trauma or abnormality.

A polynucleotide or polypeptide and/or agonist or antagonist of the present invention may increase chemotactic activity of particular cells. These chemotactic molecules can then be used to treat inflammation, infection, hyperproliferative
30 disorders, or any immune system disorder by increasing the number of cells targeted to a particular location in the body. For example, chemotactic molecules can be used

to treat wounds and other trauma to tissues by attracting immune cells to the injured location. Chemotactic molecules of the present invention can also attract fibroblasts, which can be used to treat wounds.

It is also contemplated that a polynucleotide or polypeptide and/or agonist or antagonist of the present invention may inhibit chemotactic activity. These molecules could also be used to treat disorders. Thus, a polynucleotide or polypeptide and/or agonist or antagonist of the present invention could be used as an inhibitor of chemotaxis.

10 **Binding Activity**

A polypeptide of the present invention may be used to screen for molecules that bind to the polypeptide or for molecules to which the polypeptide binds. The binding of the polypeptide and the molecule may activate (agonist), increase, inhibit (antagonist), or decrease activity of the polypeptide or the molecule bound. Examples of such molecules include antibodies, oligonucleotides, proteins (e.g., receptors), or small molecules.

Preferably, the molecule is closely related to the natural ligand of the polypeptide, e.g., a fragment of the ligand, or a natural substrate, a ligand, a structural or functional mimetic. (See, Coligan et al., Current Protocols in Immunology 1(2):Chapter 5 (1991).) Similarly, the molecule can be closely related to the natural receptor to which the polypeptide binds, or at least, a fragment of the receptor capable of being bound by the polypeptide (e.g., active site). In either case, the molecule can be rationally designed using known techniques.

Preferably, the screening for these molecules involves producing appropriate cells which express the polypeptide, either as a secreted protein or on the cell membrane. Preferred cells include cells from mammals, yeast, *Drosophila*, or *E. coli*. Cells expressing the polypeptide (or cell membrane containing the expressed polypeptide) are then preferably contacted with a test compound potentially containing the molecule to observe binding, stimulation, or inhibition of activity of either the polypeptide or the molecule.

The assay may simply test binding of a candidate compound to the polypeptide, wherein binding is detected by a label, or in an assay involving competition with a labeled competitor. Further, the assay may test whether the candidate compound results in a signal generated by binding to the polypeptide.

5 Alternatively, the assay can be carried out using cell-free preparations, polypeptide/molecule affixed to a solid support, chemical libraries, or natural product mixtures. The assay may also simply comprise the steps of mixing a candidate compound with a solution containing a polypeptide, measuring polypeptide/molecule activity or binding, and comparing the polypeptide/molecule activity or binding to a
10 standard.

Preferably, an ELISA assay can measure polypeptide level or activity in a sample (e.g., biological sample) using a monoclonal or polyclonal antibody. The antibody can measure polypeptide level or activity by either binding, directly or indirectly, to the polypeptide or by competing with the polypeptide for a substrate.

15 Additionally, the receptor to which a polypeptide of the invention binds can be identified by numerous methods known to those of skill in the art, for example, ligand panning and FACS sorting (Coligan, et al., Current Protocols in Immun., 1(2), Chapter 5, (1991)). For example, expression cloning is employed wherein polyadenylated RNA is prepared from a cell responsive to the polypeptides, for
20 example, NIH3T3 cells which are known to contain multiple receptors for the FGF family proteins, and SC-3 cells, and a cDNA library created from this RNA is divided into pools and used to transfect COS cells or other cells that are not responsive to the polypeptides. Transfected cells which are grown on glass slides are exposed to the polypeptide of the present invention, after they have been labelled. The polypeptides
25 can be labeled by a variety of means including iodination or inclusion of a recognition site for a site-specific protein kinase.

Following fixation and incubation, the slides are subjected to auto-radiographic analysis. Positive pools are identified and sub-pools are prepared and re-transfected using an iterative sub-pooling and re-screening process, eventually
30 yielding a single clones that encodes the putative receptor.

As an alternative approach for receptor identification, the labeled polypeptides can be photoaffinity linked with cell membrane or extract preparations that express the receptor molecule. Cross-linked material is resolved by PAGE analysis and exposed to X-ray film. The labeled complex containing the receptors of the polypeptides can be excised, resolved into peptide fragments, and subjected to protein microsequencing. The amino acid sequence obtained from microsequencing would be used to design a set of degenerate oligonucleotide probes to screen a cDNA library to identify the genes encoding the putative receptors.

Moreover, the techniques of gene-shuffling, motif-shuffling, exon-shuffling, and/or codon-shuffling (collectively referred to as "DNA shuffling") may be employed to modulate the activities of polypeptides of the invention thereby effectively generating agonists and antagonists of polypeptides of the invention. See generally, U.S. Patent Nos. 5,605,793, 5,811,238, 5,830,721, 5,834,252, and 5,837,458, and Patten, P. A., et al., *Curr. Opinion Biotechnol.* 8:724-33 (1997); Harayama, S. *Trends Biotechnol.* 16(2):76-82 (1998); Hansson, L. O., et al., *J. Mol. Biol.* 287:265-76 (1999); and Lorenzo, M. M. and Blasco, R. *Biotechniques* 24(2):308-13 (1998) (each of these patents and publications are hereby incorporated by reference). In one embodiment, alteration of polynucleotides and corresponding polypeptides of the invention may be achieved by DNA shuffling. DNA shuffling involves the assembly of two or more DNA segments into a desired polynucleotide sequence of the invention molecule by homologous, or site-specific, recombination. In another embodiment, polynucleotides and corresponding polypeptides of the invention may be altered by being subjected to random mutagenesis by error-prone PCR, random nucleotide insertion or other methods prior to recombination. In another embodiment, one or more components, motifs, sections, parts, domains, fragments, etc., of the polypeptides of the invention may be recombined with one or more components, motifs, sections, parts, domains, fragments, etc. of one or more heterologous molecules. In preferred embodiments, the heterologous molecules are family members. In further preferred embodiments, the heterologous molecule is a growth factor such as, for example, platelet-derived growth factor (PDGF), insulin-like growth factor (IGF-I), transforming growth factor (TGF)-alpha, epidermal

growth factor (EGF), fibroblast growth factor (FGF), TGF-beta, bone morphogenetic protein (BMP)-2, BMP-4, BMP-5, BMP-6, BMP-7, activins A and B, decapentaplegic(dpp), 60A, OP-2, dorsalin, growth differentiation factors (GDFs), nodal, MIS, inhibin-alpha, TGF-beta1, TGF-beta2, TGF-beta3, TGF-beta5, and glial-
5 derived neurotrophic factor (GDNF).

Other preferred fragments are biologically active fragments of the polypeptides of the invention. Biologically active fragments are those exhibiting activity similar, but not necessarily identical, to an activity of the polypeptide. The biological activity of the fragments may include an improved desired activity, or a
10 decreased undesirable activity.

Additionally, this invention provides a method of screening compounds to identify those which modulate the action of the polypeptide of the present invention. An example of such an assay comprises combining a mammalian fibroblast cell, a the polypeptide of the present invention, the compound to be screened and 3[H]
15 thymidine under cell culture conditions where the fibroblast cell would normally proliferate. A control assay may be performed in the absence of the compound to be screened and compared to the amount of fibroblast proliferation in the presence of the compound to determine if the compound stimulates proliferation by determining the uptake of 3[H] thymidine in each case. The amount of fibroblast cell proliferation is
20 measured by liquid scintillation chromatography which measures the incorporation of 3[H] thymidine. Both agonist and antagonist compounds may be identified by this procedure.

In another method, a mammalian cell or membrane preparation expressing a receptor for a polypeptide of the present invention is incubated with a labeled
25 polypeptide of the present invention in the presence of the compound. The ability of the compound to enhance or block this interaction could then be measured. Alternatively, the response of a known second messenger system following interaction of a compound to be screened and the receptor is measured and the ability of the compound to bind to the receptor and elicit a second messenger response is
30 measured to determine if the compound is a potential agonist or antagonist. Such

second messenger systems include but are not limited to, cAMP guanylate cyclase, ion channels or phosphoinositide hydrolysis.

All of these above assays can be used as diagnostic or prognostic markers. The molecules discovered using these assays can be used to treat disease or to bring about a particular result in a patient (e.g., blood vessel growth) by activating or inhibiting the polypeptide/molecule. Moreover, the assays can discover agents which may inhibit or enhance the production of the polypeptides of the invention from suitably manipulated cells or tissues. Therefore, the invention includes a method of identifying compounds which bind to the polypeptides of the invention comprising the steps of: (a) incubating a candidate binding compound with the polypeptide; and (b) determining if binding has occurred. Moreover, the invention includes a method of identifying agonists/antagonists comprising the steps of: (a) incubating a candidate compound with the polypeptide, (b) assaying a biological activity, and (b) determining if a biological activity of the polypeptide has been altered.

Also, one could identify molecules bind a polypeptide of the invention experimentally by using the beta-pleated sheet regions contained in the polypeptide sequence of the protein. Accordingly, specific embodiments of the invention are directed to polynucleotides encoding polypeptides which comprise, or alternatively consist of, the amino acid sequence of each beta pleated sheet regions in a disclosed polypeptide sequence. Additional embodiments of the invention are directed to polynucleotides encoding polypeptides which comprise, or alternatively consist of, any combination or all of contained in the polypeptide sequences of the invention. Additional preferred embodiments of the invention are directed to polypeptides which comprise, or alternatively consist of, the amino acid sequence of each of the beta pleated sheet regions in one of the polypeptide sequences of the invention. Additional embodiments of the invention are directed to polypeptides which comprise, or alternatively consist of, any combination or all of the beta pleated sheet regions in one of the polypeptide sequences of the invention.

30 **Targeted Delivery**

In another embodiment, the invention provides a method of delivering

compositions to targeted cells expressing a receptor for a polypeptide of the invention, or cells expressing a cell bound form of a polypeptide of the invention.

As discussed herein, polypeptides or antibodies of the invention may be associated with heterologous polypeptides, heterologous nucleic acids, toxins, or prodrugs via hydrophobic, hydrophilic, ionic and/or covalent interactions. In one embodiment, the invention provides a method for the specific delivery of compositions of the invention to cells by administering polypeptides of the invention (including antibodies) that are associated with heterologous polypeptides or nucleic acids. In one example, the invention provides a method for delivering a therapeutic protein into the targeted cell. In another example, the invention provides a method for delivering a single stranded nucleic acid (e.g., antisense or ribozymes) or double stranded nucleic acid (e.g., DNA that can integrate into the cell's genome or replicate episomally and that can be transcribed) into the targeted cell.

In another embodiment, the invention provides a method for the specific destruction of cells (e.g., the destruction of tumor cells) by administering polypeptides of the invention (e.g., polypeptides of the invention or antibodies of the invention) in association with toxins or cytotoxic prodrugs.

By "toxin" is meant compounds that bind and activate endogenous cytotoxic effector systems, radioisotopes, holotoxins, modified toxins, catalytic subunits of toxins, or any molecules or enzymes not normally present in or on the surface of a cell that under defined conditions cause the cell's death. Toxins that may be used according to the methods of the invention include, but are not limited to, radioisotopes known in the art, compounds such as, for example, antibodies (or complement fixing containing portions thereof) that bind an inherent or induced endogenous cytotoxic effector system, thymidine kinase, endonuclease, RNase, alpha toxin, ricin, abrin, *Pseudomonas* exotoxin A, diphtheria toxin, saporin, momordin, gelonin, pokeweed antiviral protein, alpha-sarcin and cholera toxin. By "cytotoxic prodrug" is meant a non-toxic compound that is converted by an enzyme, normally present in the cell, into a cytotoxic compound. Cytotoxic prodrugs that may be used according to the methods of the invention include, but are not limited to, glutamyl derivatives of benzoic acid mustard alkylating agent, phosphate derivatives of etoposide or

mitomycin C, cytosine arabinoside, daunorubisin, and phenoxyacetamide derivatives of doxorubicin.

Drug Screening

5 Further contemplated is the use of the polypeptides of the present invention, or the polynucleotides encoding these polypeptides, to screen for molecules which modify the activities of the polypeptides of the present invention. Such a method would include contacting the polypeptide of the present invention with a selected compound(s) suspected of having antagonist or agonist activity, and assaying the
10 activity of these polypeptides following binding.

This invention is particularly useful for screening therapeutic compounds by using the polypeptides of the present invention, or binding fragments thereof, in any of a variety of drug screening techniques. The polypeptide or fragment employed in such a test may be affixed to a solid support, expressed on a cell surface, free in
15 solution, or located intracellularly. One method of drug screening utilizes eukaryotic or prokaryotic host cells which are stably transformed with recombinant nucleic acids expressing the polypeptide or fragment. Drugs are screened against such transformed cells in competitive binding assays. One may measure, for example, the formulation of complexes between the agent being tested and a polypeptide of the present
20 invention.

Thus, the present invention provides methods of screening for drugs or any other agents which affect activities mediated by the polypeptides of the present invention. These methods comprise contacting such an agent with a polypeptide of the present invention or a fragment thereof and assaying for the presence of a complex
25 between the agent and the polypeptide or a fragment thereof, by methods well known in the art. In such a competitive binding assay, the agents to screen are typically labeled. Following incubation, free agent is separated from that present in bound form, and the amount of free or uncomplexed label is a measure of the ability of a particular agent to bind to the polypeptides of the present invention.

30 Another technique for drug screening provides high throughput screening for compounds having suitable binding affinity to the polypeptides of the present

invention, and is described in great detail in European Patent Application 84/03564, published on September 13, 1984, which is incorporated herein by reference herein. Briefly stated, large numbers of different small peptide test compounds are synthesized on a solid substrate, such as plastic pins or some other surface. The peptide test compounds are reacted with polypeptides of the present invention and washed. Bound polypeptides are then detected by methods well known in the art. Purified polypeptides are coated directly onto plates for use in the aforementioned drug screening techniques. In addition, non-neutralizing antibodies may be used to capture the peptide and immobilize it on the solid support.

10 This invention also contemplates the use of competitive drug screening assays in which neutralizing antibodies capable of binding polypeptides of the present invention specifically compete with a test compound for binding to the polypeptides or fragments thereof. In this manner, the antibodies are used to detect the presence of any peptide which shares one or more antigenic epitopes with a polypeptide of the invention.

Antisense And Ribozyme (Antagonists)

In specific embodiments, antagonists according to the present invention are nucleic acids corresponding to the sequences contained in SEQ ID NO:X, or the complementary strand thereof, and/or to nucleotide sequences contained a deposited clone. In one embodiment, antisense sequence is generated internally by the organism, in another embodiment, the antisense sequence is separately administered (see, for example, O'Connor, Neurochem., 56:560 (1991). Oligodeoxynucleotides as Antisense Inhibitors of Gene Expression, CRC Press, Boca Raton, FL (1988).

25 Antisense technology can be used to control gene expression through antisense DNA or RNA, or through triple-helix formation. Antisense techniques are discussed for example, in Okano, Neurochem., 56:560 (1991); Oligodeoxynucleotides as Antisense Inhibitors of Gene Expression, CRC Press, Boca Raton, FL (1988). Triple helix formation is discussed in, for instance, Lee et al., Nucleic Acids Research, 6:3073

30 (1979); Cooney et al., Science, 241:456 (1988); and Dervan et al., Science, 251:1300

(1991). The methods are based on binding of a polynucleotide to a complementary DNA or RNA.

For example, the use of c-myc and c-myb antisense RNA constructs to inhibit the growth of the non-lymphocytic leukemia cell line HL-60 and other cell lines was previously described. (Wickstrom et al. (1988); Anfossi et al. (1989)). These experiments were performed in vitro by incubating cells with the oligoribonucleotide. A similar procedure for in vivo use is described in WO 91/15580. Briefly, a pair of oligonucleotides for a given antisense RNA is produced as follows: A sequence complimentary to the first 15 bases of the open reading frame is flanked by an EcoRI site on the 5' end and a HindIII site on the 3' end. Next, the pair of oligonucleotides is heated at 90°C for one minute and then annealed in 2X ligation buffer (20mM TRIS HCl pH 7.5, 10mM MgCl₂, 10mM dithiothreitol (DTT) and 0.2 mM ATP) and then ligated to the EcoRI/Hind III site of the retroviral vector PMV7 (WO 91/15580).

For example, the 5' coding portion of a polynucleotide that encodes the mature polypeptide of the present invention may be used to design an antisense RNA oligonucleotide of from about 10 to 40 base pairs in length. A DNA oligonucleotide is designed to be complementary to a region of the gene involved in transcription thereby preventing transcription and the production of the receptor. The antisense RNA oligonucleotide hybridizes to the mRNA in vivo and blocks translation of the mRNA molecule into receptor polypeptide.

In one embodiment, the antisense nucleic acid of the invention is produced intracellularly by transcription from an exogenous sequence. For example, a vector or a portion thereof, is transcribed, producing an antisense nucleic acid (RNA) of the invention. Such a vector would contain a sequence encoding the antisense nucleic acid of the invention. Such a vector can remain episomal or become chromosomally integrated, as long as it can be transcribed to produce the desired antisense RNA. Such vectors can be constructed by recombinant DNA technology methods standard in the art. Vectors can be plasmid, viral, or others known in the art, used for replication and expression in vertebrate cells. Expression of the sequence encoding a polypeptide of the invention, or fragments thereof, can be by any promoter known in the art to act in vertebrate, preferably human cells. Such promoters can be inducible

or constitutive. Such promoters include, but are not limited to, the SV40 early promoter region (Bernoist and Chambon, *Nature*, 29:304-310 (1981), the promoter contained in the 3' long terminal repeat of Rous sarcoma virus (Yamamoto et al., *Cell*, 22:787-797 (1980), the herpes thymidine promoter (Wagner et al., *Proc. Natl. Acad. Sci. U.S.A.*, 78:1441-1445 (1981), the regulatory sequences of the metallothionein gene (Brinster et al., *Nature*, 296:39-42 (1982)), etc.

The antisense nucleic acids of the invention comprise a sequence complementary to at least a portion of an RNA transcript of a gene of interest. However, absolute complementarity, although preferred, is not required. A sequence
10 "complementary to at least a portion of an RNA," referred to herein, means a sequence having sufficient complementarity to be able to hybridize with the RNA, forming a stable duplex; in the case of double stranded antisense nucleic acids of the invention, a single strand of the duplex DNA may thus be tested, or triplex formation may be assayed. The ability to hybridize will depend on both the degree of
15 complementarity and the length of the antisense nucleic acid. Generally, the larger the hybridizing nucleic acid, the more base mismatches with a RNA sequence of the invention it may contain and still form a stable duplex (or triplex as the case may be). One skilled in the art can ascertain a tolerable degree of mismatch by use of standard procedures to determine the melting point of the hybridized complex.

20 Oligonucleotides that are complementary to the 5' end of the message, *e.g.*, the 5' untranslated sequence up to and including the AUG initiation codon, should work most efficiently at inhibiting translation. However, sequences complementary to the 3' untranslated sequences of mRNAs have been shown to be effective at inhibiting translation of mRNAs as well. See generally, Wagner, R., *Nature*,
25 372:333-335 (1994). Thus, oligonucleotides complementary to either the 5' - or 3' - non-translated, non-coding regions of a polynucleotide sequence of the invention could be used in an antisense approach to inhibit translation of endogenous mRNA. Oligonucleotides complementary to the 5' untranslated region of the mRNA should include the complement of the AUG start codon. Antisense oligonucleotides
30 complementary to mRNA coding regions are less efficient inhibitors of translation but could be used in accordance with the invention. Whether designed to hybridize to the

5' -, 3' - or coding region of mRNA, antisense nucleic acids should be at least six nucleotides in length, and are preferably oligonucleotides ranging from 6 to about 50 nucleotides in length. In specific aspects the oligonucleotide is at least 10 nucleotides, at least 17 nucleotides, at least 25 nucleotides or at least 50 nucleotides.

5 The polynucleotides of the invention can be DNA or RNA or chimeric mixtures or derivatives or modified versions thereof, single-stranded or double-stranded. The oligonucleotide can be modified at the base moiety, sugar moiety, or phosphate backbone, for example, to improve stability of the molecule, hybridization, etc. The oligonucleotide may include other appended groups such as peptides (e.g.,
10 for targeting host cell receptors in vivo), or agents facilitating transport across the cell membrane (see, e.g., Letsinger et al., Proc. Natl. Acad. Sci. U.S.A. 86:6553-6556 (1989); Lemaitre et al., Proc. Natl. Acad. Sci., 84:648-652 (1987); PCT Publication NO: WO88/09810, published December 15, 1988) or the blood-brain barrier (see, e.g., PCT Publication NO: WO89/10134, published April 25, 1988), hybridization-
15 triggered cleavage agents. (See, e.g., Krol et al., BioTechniques, 6:958-976 (1988)) or intercalating agents. (See, e.g., Zon, Pharm. Res., 5:539-549 (1988)). To this end, the oligonucleotide may be conjugated to another molecule, e.g., a peptide, hybridization triggered cross-linking agent, transport agent, hybridization-triggered cleavage agent, etc.

20 The antisense oligonucleotide may comprise at least one modified base moiety which is selected from the group including, but not limited to, 5-fluorouracil, 5-bromouracil, 5-chlorouracil, 5-iodouracil, hypoxanthine, xantine, 4-acetylcytosine, 5-(carboxyhydroxymethyl) uracil, 5-carboxymethylaminomethyl-2-thiouridine, 5-carboxymethylaminomethyluracil, dihydrouracil, beta-D-galactosylqueosine,
25 inosine, N6-isopentenyladenine, 1-methylguanine, 1-methylinosine, 2,2-dimethylguanine, 2-methyladenine, 2-methylguanine, 3-methylcytosine, 5-methylcytosine, N6-adenine, 7-methylguanine, 5-methylaminomethyluracil, 5-methoxymethylaminomethyl-2-thiouracil, beta-D-mannosylqueosine, 5'-methoxycarboxymethyluracil, 5-methoxyuracil, 2-methylthio-N6-
30 isopentenyladenine, uracil-5-oxyacetic acid (v), wybutoxosine, pseudouracil, queosine, 2-thiocytosine, 5-methyl-2-thiouracil, 2-thiouracil, 4-thiouracil,

5-methyluracil, uracil-5-oxyacetic acid methylester, uracil-5-oxyacetic acid (v),
5-methyl-2-thiouracil, 3-(3-amino-3-N-2-carboxypropyl) uracil, (acp3)w, and
2,6-diaminopurine.

5 The antisense oligonucleotide may also comprise at least one modified sugar
moiety selected from the group including, but not limited to, arabinose,
2-fluoroarabinose, xylulose, and hexose.

In yet another embodiment, the antisense oligonucleotide comprises at least
one modified phosphate backbone selected from the group including, but not limited
to, a phosphorothioate, a phosphorodithioate, a phosphoramidothioate, a
10 phosphoramidate, a phosphordiamidate, a methylphosphonate, an alkyl
phosphotriester, and a formacetal or analog thereof.

In yet another embodiment, the antisense oligonucleotide is an a-anomeric
oligonucleotide. An a-anomeric oligonucleotide forms specific double-stranded
hybrids with complementary RNA in which, contrary to the usual b-units, the strands
15 run parallel to each other (Gautier et al., Nucl. Acids Res., 15:6625-6641 (1987)).
The oligonucleotide is a 2-O-methylribonucleotide (Inoue et al., Nucl. Acids Res.,
15:6131-6148 (1987)), or a chimeric RNA-DNA analogue (Inoue et al., FEBS Lett.
215:327-330 (1987)).

Polynucleotides of the invention may be synthesized by standard methods
20 known in the art, e.g. by use of an automated DNA synthesizer (such as are
commercially available from Biosearch, Applied Biosystems, etc.). As examples,
phosphorothioate oligonucleotides may be synthesized by the method of Stein et al.
(Nucl. Acids Res., 16:3209 (1988)), methylphosphonate oligonucleotides can be
prepared by use of controlled pore glass polymer supports (Sarin et al., Proc. Natl.
25 Acad. Sci. U.S.A., 85:7448-7451 (1988)), etc.

While antisense nucleotides complementary to the coding region sequence of
the invention could be used, those complementary to the transcribed untranslated
region are most preferred.

Potential antagonists according to the invention also include catalytic RNA, or
30 a ribozyme (See, e.g., PCT International Publication WO 90/11364, published
October 4, 1990; Sarver et al, Science, 247:1222-1225 (1990). While ribozymes that

cleave mRNA at site specific recognition sequences can be used to destroy mRNAs corresponding to the polynucleotides of the invention, the use of hammerhead ribozymes is preferred. Hammerhead ribozymes cleave mRNAs at locations dictated by flanking regions that form complementary base pairs with the target mRNA. The sole requirement is that the target mRNA have the following sequence of two bases: 5'-UG-3'. The construction and production of hammerhead ribozymes is well known in the art and is described more fully in Haseloff and Gerlach, *Nature*, 334:585-591 (1988). There are numerous potential hammerhead ribozyme cleavage sites within each nucleotide sequence disclosed in the sequence listing. Preferably, the ribozyme is engineered so that the cleavage recognition site is located near the 5' end of the mRNA corresponding to the polynucleotides of the invention; i.e., to increase efficiency and minimize the intracellular accumulation of non-functional mRNA transcripts.

As in the antisense approach, the ribozymes of the invention can be composed of modified oligonucleotides (e.g. for improved stability, targeting, etc.) and should be delivered to cells which express the polynucleotides of the invention in vivo. DNA constructs encoding the ribozyme may be introduced into the cell in the same manner as described above for the introduction of antisense encoding DNA. A preferred method of delivery involves using a DNA construct "encoding" the ribozyme under the control of a strong constitutive promoter, such as, for example, pol III or pol II promoter, so that transfected cells will produce sufficient quantities of the ribozyme to destroy endogenous messages and inhibit translation. Since ribozymes unlike antisense molecules, are catalytic, a lower intracellular concentration is required for efficiency.

Antagonist/agonist compounds may be employed to inhibit the cell growth and proliferation effects of the polypeptides of the present invention on neoplastic cells and tissues, i.e. stimulation of angiogenesis of tumors, and, therefore, retard or prevent abnormal cellular growth and proliferation, for example, in tumor formation or growth.

The antagonist/agonist may also be employed to prevent hyper-vascular diseases, and prevent the proliferation of epithelial lens cells after extracapsular

cataract surgery. Prevention of the mitogenic activity of the polypeptides of the present invention may also be desirous in cases such as restenosis after balloon angioplasty.

5 The antagonist/agonist may also be employed to prevent the growth of scar tissue during wound healing.

The antagonist/agonist may also be employed to treat the diseases described herein.

Thus, the invention provides a method of treating disorders or diseases, including but not limited to the disorders or diseases listed throughout this application, associated
10 with overexpression of a polynucleotide of the present invention by administering to a patient (a) an antisense molecule directed to the polynucleotide of the present invention, and/or (b) a ribozyme directed to the polynucleotide of the present invention

15 **Other Activities**

The polypeptide of the present invention, as a result of the ability to stimulate vascular endothelial cell growth, may be employed in treatment for stimulating re-vascularization of ischemic tissues due to various disease conditions such as thrombosis, arteriosclerosis, and other cardiovascular conditions. These polypeptide
20 may also be employed to stimulate angiogenesis and limb regeneration, as discussed above.

The polypeptide may also be employed for treating wounds due to injuries, burns, post-operative tissue repair, and ulcers since they are mitogenic to various cells of different origins, such as fibroblast cells and skeletal muscle cells, and therefore,
25 facilitate the repair or replacement of damaged or diseased tissue.

The polypeptide of the present invention may also be employed stimulate neuronal growth and to treat and prevent neuronal damage which occurs in certain neuronal disorders or neuro-degenerative conditions such as Alzheimer's disease, Parkinson's disease, and AIDS-related complex. The polypeptide of the invention
30 may have the ability to stimulate chondrocyte growth, therefore, they may be

employed to enhance bone and periodontal regeneration and aid in tissue transplants or bone grafts.

The polypeptide of the present invention may be also be employed to prevent skin aging due to sunburn by stimulating keratinocyte growth.

5 The polypeptide of the invention may also be employed for preventing hair loss, since FGF family members activate hair-forming cells and promotes melanocyte growth. Along the same lines, the polypeptides of the present invention may be employed to stimulate growth and differentiation of hematopoietic cells and bone marrow cells when used in combination with other cytokines.

10 The polypeptide of the invention may also be employed to maintain organs before transplantation or for supporting cell culture of primary tissues.

The polypeptide of the present invention may also be employed for inducing tissue of mesodermal origin to differentiate in early embryos.

15 The polypeptide or polynucleotides and/or agonist or antagonists of the present invention may also increase or decrease the differentiation or proliferation of embryonic stem cells, besides, as discussed above, hematopoietic lineage.

20 The polypeptide or polynucleotides and/or agonist or antagonists of the present invention may also be used to modulate mammalian characteristics, such as body height, weight, hair color, eye color, skin, percentage of adipose tissue, pigmentation, size, and shape (e.g., cosmetic surgery). Similarly, polypeptides or polynucleotides and/or agonist or antagonists of the present invention may be used to modulate mammalian metabolism affecting catabolism, anabolism, processing, utilization, and storage of energy.

25 Polypeptide or polynucleotides and/or agonist or antagonists of the present invention may be used to change a mammal's mental state or physical state by influencing biorhythms, circadian rhythms, depression (including depressive disorders), tendency for violence, tolerance for pain, reproductive capabilities (preferably by Activin or Inhibin-like activity), hormonal or endocrine levels, appetite, libido, memory, stress, or other cognitive qualities.

30 Polypeptide or polynucleotides and/or agonist or antagonists of the present invention may also be used as a food additive or preservative, such as to increase or

decrease storage capabilities, fat content, lipid, protein, carbohydrate, vitamins, minerals, cofactors or other nutritional components.

Other Preferred Embodiments

5 Other preferred embodiments of the claimed invention include an isolated nucleic acid molecule comprising a nucleotide sequence which is at least 95% identical to a sequence of at least about 50 contiguous nucleotides in the nucleotide sequence of SEQ ID NO:X wherein X is any integer as defined in Table 1.

10 Also preferred is a nucleic acid molecule wherein said sequence of contiguous nucleotides is included in the nucleotide sequence of SEQ ID NO:X in the range of positions beginning with the nucleotide at about the position of the 5' Nucleotide of the Clone Sequence and ending with the nucleotide at about the position of the 3' Nucleotide of the Clone Sequence as defined for SEQ ID NO:X in Table 1.

15 Also preferred is a nucleic acid molecule wherein said sequence of contiguous nucleotides is included in the nucleotide sequence of SEQ ID NO:X in the range of positions beginning with the nucleotide at about the position of the 5' Nucleotide of the Start Codon and ending with the nucleotide at about the position of the 3' Nucleotide of the Clone Sequence as defined for SEQ ID NO:X in Table 1.

20 Similarly preferred is a nucleic acid molecule wherein said sequence of contiguous nucleotides is included in the nucleotide sequence of SEQ ID NO:X in the range of positions beginning with the nucleotide at about the position of the 5' Nucleotide of the First Amino Acid of the Signal Peptide and ending with the nucleotide at about the position of the 3' Nucleotide of the Clone Sequence as defined for SEQ ID NO:X in Table 1.

25 Also preferred is an isolated nucleic acid molecule comprising a nucleotide sequence which is at least 95% identical to a sequence of at least about 150 contiguous nucleotides in the nucleotide sequence of SEQ ID NO:X.

30 Further preferred is an isolated nucleic acid molecule comprising a nucleotide sequence which is at least 95% identical to a sequence of at least about 500 contiguous nucleotides in the nucleotide sequence of SEQ ID NO:X.

A further preferred embodiment is a nucleic acid molecule comprising a nucleotide sequence which is at least 95% identical to the nucleotide sequence of SEQ ID NO:X beginning with the nucleotide at about the position of the 5' Nucleotide of the First Amino Acid of the Signal Peptide and ending with the nucleotide at about the position of the 3' Nucleotide of the Clone Sequence as defined for SEQ ID NO:X in Table 1.

A further preferred embodiment is an isolated nucleic acid molecule comprising a nucleotide sequence which is at least 95% identical to the complete nucleotide sequence of SEQ ID NO:X.

Also preferred is an isolated nucleic acid molecule which hybridizes under stringent hybridization conditions to a nucleic acid molecule, wherein said nucleic acid molecule which hybridizes does not hybridize under stringent hybridization conditions to a nucleic acid molecule having a nucleotide sequence consisting of only A residues or of only T residues.

Also preferred is a composition of matter comprising a DNA molecule which comprises a human cDNA clone identified by a cDNA Clone Identifier in Table 1, which DNA molecule is contained in the material deposited with the American Type Culture Collection and given the ATCC Deposit Number shown in Table 1 for said cDNA Clone Identifier.

Also preferred is an isolated nucleic acid molecule comprising a nucleotide sequence which is at least 95% identical to a sequence of at least 50 contiguous nucleotides in the nucleotide sequence of a human cDNA clone identified by a cDNA Clone Identifier in Table 1, which DNA molecule is contained in the deposit given the ATCC Deposit Number shown in Table 1.

Also preferred is an isolated nucleic acid molecule, wherein said sequence of at least 50 contiguous nucleotides is included in the nucleotide sequence of the complete open reading frame sequence encoded by said human cDNA clone.

Also preferred is an isolated nucleic acid molecule comprising a nucleotide sequence which is at least 95% identical to sequence of at least 150 contiguous nucleotides in the nucleotide sequence encoded by said human cDNA clone.

A further preferred embodiment is an isolated nucleic acid molecule comprising a nucleotide sequence which is at least 95% identical to sequence of at least 500 contiguous nucleotides in the nucleotide sequence encoded by said human cDNA clone.

5 A further preferred embodiment is an isolated nucleic acid molecule comprising a nucleotide sequence which is at least 95% identical to the complete nucleotide sequence encoded by said human cDNA clone.

A further preferred embodiment is a method for detecting in a biological sample a nucleic acid molecule comprising a nucleotide sequence which is at least
10 95% identical to a sequence of at least 50 contiguous nucleotides in a sequence selected from the group consisting of: a nucleotide sequence of SEQ ID NO:X wherein X is any integer as defined in Table 1; and a nucleotide sequence encoded by a human cDNA clone identified by a cDNA Clone Identifier in Table 1 and contained in the deposit with the ATCC Deposit Number shown for said cDNA clone in Table
15 1; which method comprises a step of comparing a nucleotide sequence of at least one nucleic acid molecule in said sample with a sequence selected from said group and determining whether the sequence of said nucleic acid molecule in said sample is at least 95% identical to said selected sequence.

Also preferred is the above method wherein said step of comparing sequences
20 comprises determining the extent of nucleic acid hybridization between nucleic acid molecules in said sample and a nucleic acid molecule comprising said sequence selected from said group. Similarly, also preferred is the above method wherein said step of comparing sequences is performed by comparing the nucleotide sequence determined from a nucleic acid molecule in said sample with said sequence selected
25 from said group. The nucleic acid molecules can comprise DNA molecules or RNA molecules.

A further preferred embodiment is a method for identifying the species, tissue or cell type of a biological sample which method comprises a step of detecting nucleic acid molecules in said sample, if any, comprising a nucleotide sequence that is at least
30 95% identical to a sequence of at least 50 contiguous nucleotides in a sequence selected from the group consisting of: a nucleotide sequence of SEQ ID NO:X

wherein X is any integer as defined in Table 1; and a nucleotide sequence encoded by a human cDNA clone identified by a cDNA Clone Identifier in Table 1 and contained in the deposit with the ATCC Deposit Number shown for said cDNA clone in Table 1.

5 The method for identifying the species, tissue or cell type of a biological sample can comprise a step of detecting nucleic acid molecules comprising a nucleotide sequence in a panel of at least two nucleotide sequences, wherein at least one sequence in said panel is at least 95% identical to a sequence of at least 50 contiguous nucleotides in a sequence selected from said group.

10 Also preferred is a method for diagnosing in a subject a pathological condition associated with abnormal structure or expression of a gene encoding a secreted protein identified in Table 1, which method comprises a step of detecting in a biological sample obtained from said subject nucleic acid molecules, if any, comprising a nucleotide sequence that is at least 95% identical to a sequence of at
15 least 50 contiguous nucleotides in a sequence selected from the group consisting of: a nucleotide sequence of SEQ ID NO:X wherein X is any integer as defined in Table 1; and a nucleotide sequence encoded by a human cDNA clone identified by a cDNA Clone Identifier in Table 1 and contained in the deposit with the ATCC Deposit Number shown for said cDNA clone in Table 1.

20 The method for diagnosing a pathological condition can comprise a step of detecting nucleic acid molecules comprising a nucleotide sequence in a panel of at least two nucleotide sequences, wherein at least one sequence in said panel is at least 95% identical to a sequence of at least 50 contiguous nucleotides in a sequence selected from said group.

25 Also preferred is a composition of matter comprising isolated nucleic acid molecules wherein the nucleotide sequences of said nucleic acid molecules comprise a panel of at least two nucleotide sequences, wherein at least one sequence in said panel is at least 95% identical to a sequence of at least 50 contiguous nucleotides in a sequence selected from the group consisting of: a nucleotide sequence of SEQ ID
30 NO:X wherein X is any integer as defined in Table 1; and a nucleotide sequence encoded by a human cDNA clone identified by a cDNA Clone Identifier in Table 1

and contained in the deposit with the ATCC Deposit Number shown for said cDNA clone in Table 1. The nucleic acid molecules can comprise DNA molecules or RNA molecules.

Also preferred is an isolated polypeptide comprising an amino acid sequence
5 at least 90% identical to a sequence of at least about 10 contiguous amino acids in the amino acid sequence of SEQ ID NO:Y wherein Y is any integer as defined in Table 1.

Also preferred is a polypeptide, wherein said sequence of contiguous amino acids is included in the amino acid sequence of SEQ ID NO:Y in the range of positions beginning with the residue at about the position of the First Amino Acid of
10 the Secreted Portion and ending with the residue at about the Last Amino Acid of the Open Reading Frame as set forth for SEQ ID NO:Y in Table 1.

Also preferred is an isolated polypeptide comprising an amino acid sequence at least 95% identical to a sequence of at least about 30 contiguous amino acids in the amino acid sequence of SEQ ID NO:Y.

Further preferred is an isolated polypeptide comprising an amino acid
15 sequence at least 95% identical to a sequence of at least about 100 contiguous amino acids in the amino acid sequence of SEQ ID NO:Y.

Further preferred is an isolated polypeptide comprising an amino acid sequence at least 95% identical to the complete amino acid sequence of SEQ ID
20 NO:Y.

Further preferred is an isolated polypeptide comprising an amino acid sequence at least 90% identical to a sequence of at least about 10 contiguous amino acids in the complete amino acid sequence of a secreted protein encoded by a human cDNA clone identified by a cDNA Clone Identifier in Table 1 and contained in the
25 deposit with the ATCC Deposit Number shown for said cDNA clone in Table 1.

Also preferred is a polypeptide wherein said sequence of contiguous amino acids is included in the amino acid sequence of a secreted portion of the secreted protein encoded by a human cDNA clone identified by a cDNA Clone Identifier in Table 1 and contained in the deposit with the ATCC Deposit Number shown for said
30 cDNA clone in Table 1.

Also preferred is an isolated polypeptide comprising an amino acid sequence at least 95% identical to a sequence of at least about 30 contiguous amino acids in the amino acid sequence of the secreted portion of the protein encoded by a human cDNA clone identified by a cDNA Clone Identifier in Table 1 and contained in the deposit
5 with the ATCC Deposit Number shown for said cDNA clone in Table 1.

Also preferred is an isolated polypeptide comprising an amino acid sequence at least 95% identical to a sequence of at least about 100 contiguous amino acids in the amino acid sequence of the secreted portion of the protein encoded by a human cDNA clone identified by a cDNA Clone Identifier in Table 1 and contained in the
10 deposit with the ATCC Deposit Number shown for said cDNA clone in Table 1.

Also preferred is an isolated polypeptide comprising an amino acid sequence at least 95% identical to the amino acid sequence of the secreted portion of the protein encoded by a human cDNA clone identified by a cDNA Clone Identifier in Table 1 and contained in the deposit with the ATCC Deposit Number shown for said cDNA
15 clone in Table 1.

Further preferred is an isolated antibody which binds specifically to a polypeptide comprising an amino acid sequence that is at least 90% identical to a sequence of at least 10 contiguous amino acids in a sequence selected from the group consisting of: an amino acid sequence of SEQ ID NO:Y wherein Y is any integer as
20 defined in Table 1; and a complete amino acid sequence of a protein encoded by a human cDNA clone identified by a cDNA Clone Identifier in Table 1 and contained in the deposit with the ATCC Deposit Number shown for said cDNA clone in Table 1.

Further preferred is a method for detecting in a biological sample a
25 polypeptide comprising an amino acid sequence which is at least 90% identical to a sequence of at least 10 contiguous amino acids in a sequence selected from the group consisting of: an amino acid sequence of SEQ ID NO:Y wherein Y is any integer as defined in Table 1; and a complete amino acid sequence of a protein encoded by a human cDNA clone identified by a cDNA Clone Identifier in Table 1 and contained
30 in the deposit with the ATCC Deposit Number shown for said cDNA clone in Table 1; which method comprises a step of comparing an amino acid sequence of at least

one polypeptide molecule in said sample with a sequence selected from said group and determining whether the sequence of said polypeptide molecule in said sample is at least 90% identical to said sequence of at least 10 contiguous amino acids.

Also preferred is the above method wherein said step of comparing an amino acid sequence of at least one polypeptide molecule in said sample with a sequence selected from said group comprises determining the extent of specific binding of polypeptides in said sample to an antibody which binds specifically to a polypeptide comprising an amino acid sequence that is at least 90% identical to a sequence of at least 10 contiguous amino acids in a sequence selected from the group consisting of: an amino acid sequence of SEQ ID NO:Y wherein Y is any integer as defined in Table 1; and a complete amino acid sequence of a protein encoded by a human cDNA clone identified by a cDNA Clone Identifier in Table 1 and contained in the deposit with the ATCC Deposit Number shown for said cDNA clone in Table 1.

Also preferred is the above method wherein said step of comparing sequences is performed by comparing the amino acid sequence determined from a polypeptide molecule in said sample with said sequence selected from said group.

Also preferred is a method for identifying the species, tissue or cell type of a biological sample which method comprises a step of detecting polypeptide molecules in said sample, if any, comprising an amino acid sequence that is at least 90% identical to a sequence of at least 10 contiguous amino acids in a sequence selected from the group consisting of: an amino acid sequence of SEQ ID NO:Y wherein Y is any integer as defined in Table 1; and a complete amino acid sequence of a secreted protein encoded by a human cDNA clone identified by a cDNA Clone Identifier in Table 1 and contained in the deposit with the ATCC Deposit Number shown for said cDNA clone in Table 1.

Also preferred is the above method for identifying the species, tissue or cell type of a biological sample, which method comprises a step of detecting polypeptide molecules comprising an amino acid sequence in a panel of at least two amino acid sequences, wherein at least one sequence in said panel is at least 90% identical to a sequence of at least 10 contiguous amino acids in a sequence selected from the above group.

Also preferred is a method for diagnosing in a subject a pathological condition associated with abnormal structure or expression of a gene encoding a secreted protein identified in Table 1, which method comprises a step of detecting in a biological sample obtained from said subject polypeptide molecules comprising an amino acid sequence in a panel of at least two amino acid sequences, wherein at least one sequence in said panel is at least 90% identical to a sequence of at least 10 contiguous amino acids in a sequence selected from the group consisting of: an amino acid sequence of SEQ ID NO:Y wherein Y is any integer as defined in Table 1; and a complete amino acid sequence of a secreted protein encoded by a human cDNA clone identified by a cDNA Clone Identifier in Table 1 and contained in the deposit with the ATCC Deposit Number shown for said cDNA clone in Table 1.

In any of these methods, the step of detecting said polypeptide molecules includes using an antibody.

Also preferred is an isolated nucleic acid molecule comprising a nucleotide sequence which is at least 95% identical to a nucleotide sequence encoding a polypeptide wherein said polypeptide comprises an amino acid sequence that is at least 90% identical to a sequence of at least 10 contiguous amino acids in a sequence selected from the group consisting of: an amino acid sequence of SEQ ID NO:Y wherein Y is any integer as defined in Table 1; and a complete amino acid sequence of a secreted protein encoded by a human cDNA clone identified by a cDNA Clone Identifier in Table 1 and contained in the deposit with the ATCC Deposit Number shown for said cDNA clone in Table 1.

Also preferred is an isolated nucleic acid molecule, wherein said nucleotide sequence encoding a polypeptide has been optimized for expression of said polypeptide in a prokaryotic host.

Also preferred is an isolated nucleic acid molecule, wherein said polypeptide comprises an amino acid sequence selected from the group consisting of: an amino acid sequence of SEQ ID NO:Y wherein Y is any integer as defined in Table 1; and a complete amino acid sequence of a secreted protein encoded by a human cDNA clone identified by a cDNA Clone Identifier in Table 1 and contained in the deposit with the ATCC Deposit Number shown for said cDNA clone in Table 1.

Further preferred is a method of making a recombinant vector comprising inserting any of the above isolated nucleic acid molecule into a vector. Also preferred is the recombinant vector produced by this method. Also preferred is a method of making a recombinant host cell comprising introducing the vector into a host cell, as
5 well as the recombinant host cell produced by this method.

Also preferred is a method of making an isolated polypeptide comprising culturing this recombinant host cell under conditions such that said polypeptide is expressed and recovering said polypeptide. Also preferred is this method of making an isolated polypeptide, wherein said recombinant host cell is a eukaryotic cell and
10 said polypeptide is a secreted portion of a human secreted protein comprising an amino acid sequence selected from the group consisting of: an amino acid sequence of SEQ ID NO:Y beginning with the residue at the position of the First Amino Acid of the Secreted Portion of SEQ ID NO:Y wherein Y is an integer set forth in Table 1 and said position of the First Amino Acid of the Secreted Portion of SEQ ID NO:Y is
15 defined in Table 1; and an amino acid sequence of a secreted portion of a protein encoded by a human cDNA clone identified by a cDNA Clone Identifier in Table 1 and contained in the deposit with the ATCC Deposit Number shown for said cDNA clone in Table 1. The isolated polypeptide produced by this method is also preferred.

Also preferred is a method of treatment of an individual in need of an
20 increased level of a secreted protein activity, which method comprises administering to such an individual a pharmaceutical composition comprising an amount of an isolated polypeptide, polynucleotide, or antibody of the claimed invention effective to increase the level of said protein activity in said individual.

The above-recited applications have uses in a wide variety of hosts. Such
25 hosts include, but are not limited to, human, murine, rabbit, goat, guinea pig, camel, horse, mouse, rat, hamster, pig, micro-pig, chicken, goat, cow, sheep, dog, cat, non-human primate, and human. In specific embodiments, the host is a mouse, rabbit, goat, guinea pig, chicken, rat, hamster, pig, sheep, dog or cat. In preferred embodiments, the host is a mammal. In most preferred embodiments, the host is a
30 human.

In specific embodiments of the invention, for each "Contig ID" listed in the fourth column of Table 2, preferably excluded are one or more polynucleotides comprising, or alternatively consisting of, a nucleotide sequence referenced in the fifth column of Table 2 and described by the general formula of a-b, whereas a and b are uniquely determined for the corresponding SEQ ID NO:X referred to in column 3 of Table 2. Further specific embodiments are directed to polynucleotide sequences excluding one, two, three, four, or more of the specific polynucleotide sequences referred to in the fifth column of Table 2. In no way is this listing meant to encompass all of the sequences which may be excluded by the general formula, it is just a representative example. All references available through these accessions are hereby incorporated by reference in their entirety.

TABLE 2

Gene No.	cDNA Clone ID	NT SEQ ID NO: X	Contig ID	Public Accession Numbers
4	HCE1G78	14	761204	R13943, R15390, R18788, R40003, R42130, R42130, R40003, H14886, AA074234
9	HCEEM33	19	821322	R23514, R43444, R43444, R60880, H24284, AA460540, AA460541
13	HCDCE51	23	813504	R46859, H08961
14	HCDDDB15	24	841041	T63315, T63934, R51869, R54262, R79539, H09632, H10762, H13764, H14630, H28449, H29036, H93696, H94377, N30248, N39604, N46750, N49809, N57117, N57144, W67707, W68166, AA010772, AA069124, AA085654, AA086136, AA134120, AA130292, AA148300, AA148301, AA151837, AA165188, AA190685, AA190831, AA530878, AA533129, AA554441, AA558166, AA577995, AA578080, AA730891, AA736746, AA765008, AA804389, AA825715, AA902725, AA932141, AA935282, AA946804, AA953261, AA989042, AI074021, D80545, D80546, D80821, D80809, D80822, D80988, D81381, F21272, F21575, AA488191, AA488246, AA707276, AA723413, AA775418, AA860931, AA897608, AA907930, AI095666
19	HCE1N56	29	821704	R42843, R42843, H08744, H10211, H58176, H58177, AA126915, AA128107, AA235131
26	HCE3D89	36	648612	H08731, H40434, H42581, H43056, H45987, H46052
27	HCE3N23	37	810211	R12617, R24970, R45231, R49868, R49869,

				R54204, R45231, R61154, R61870, R71727, R71726, R72040, R72039, R73431, H11117, H11222, H20519, H20608, H26600, H26639, H41309, H42232, H42305, H43070, H43094, H43154, H45150, R84953, R85207, R85995, H50076, N36397, N44272, AA084608
33	HCEDJ05	43	634525	R17521, R41888, R41888, H14380, H14430, H17901, N48190, N53337, AA010177, AA010178, AA233935, AA234206, AA253377, AA253376
35	HCEDO17	45	824312	T66294, T66346, T75467, R26105, R26316, R34588, R39238, H05085, H05544, H12764, H12808, R90842, H61362
39	HCE3J43	49	766758	R09825, R12412, H26132, H41135, R85332, R85331, R86698, R86697, R94798, H99492, N28306, N31819, N42488, N47626, N47879, N49254, N49356, N57481, N73093, W02321, AA004631, AA004232, AA029604, AA079478, AA079477, AA112602, AA161481, AA160953, AA164232, AA164233, AA191097, AA250764, AA429479
44	HCEFH31	54	801890	T81066, W73595, W73495, AA037024, AA149490, AA149494, AA235464
46	HCDCY76	56	837972	R26141, R26355, W23737, AA662446
47	HCE3B04	57	831151	T86722, N22895, N29813, N45545, N48695, N49753, N51450, N52681, N59844, N72208, W19735, W19616, W31671, N89992, N90029, AA148623, AA255954, AA256117, AA291263, AA908587, AA983273, AI014575, N55976, AA642349

Having generally described the invention, the same will be more readily understood by reference to the following examples, which are provided by way of illustration and are not intended as limiting.

5

Examples

Example 1: Isolation of a Selected cDNA Clone From the Deposited Sample

Each cDNA clone in a cited ATCC deposit is contained in a plasmid vector. Table 1 identifies the vectors used to construct the cDNA library from which each clone was isolated. In many cases, the vector used to construct the library is a phage vector from which a plasmid has been excised. The table immediately below correlates the related plasmid for each phage vector used in constructing the cDNA library. For example, where a particular clone is identified in Table 1 as being isolated in the vector "Lambda Zap," the corresponding deposited clone is in "pBluescript."

	<u>Vector Used to Construct Library</u>	<u>Corresponding Deposited</u>
	<u>Plasmid</u>	
	Lambda Zap	pBluescript (pBS)
20	Uni-Zap XR	pBluescript (pBS)
	Zap Express	pBK
	lafmid BA	plafmid BA
	pSport1	pSport1
	pCMVSPORT 2.0	pCMVSPORT 2.0
25	pCMVSPORT 3.0	pCMVSPORT 3.0
	pCR [®] 2.1	pCR [®] 2.1

Vectors Lambda Zap (U.S. Patent Nos. 5,128,256 and 5,286,636), Uni-Zap XR (U.S. Patent Nos. 5,128, 256 and 5,286,636), Zap Express (U.S. Patent Nos. 5,128,256 and 5,286,636), pBluescript (pBS) (Short, J. M. et al., Nucleic Acids Res. 16:7583-7600 (1988); Altling-Mees, M. A. and Short, J. M., Nucleic Acids Res. 17:9494 (1989)) and pBK (Altling-Mees, M. A. et al., Strategies 5:58-61 (1992)) are

30

commercially available from Stratagene Cloning Systems, Inc., 11011 N. Torrey Pines Road, La Jolla, CA, 92037. pBS contains an ampicillin resistance gene and pBK contains a neomycin resistance gene. Both can be transformed into E. coli strain XL-1 Blue, also available from Stratagene. pBS comes in 4 forms SK+, SK-, KS+ and KS. The S and K refers to the orientation of the polylinker to the T7 and T3 primer sequences which flank the polylinker region ("S" is for SacI and "K" is for KpnI which are the first sites on each respective end of the linker). "+" or "-" refer to the orientation of the fl origin of replication ("ori"), such that in one orientation, single stranded rescue initiated from the fl ori generates sense strand DNA and in the other, antisense.

Vectors pSport1, pCMVSPORT 2.0 and pCMVSPORT 3.0, were obtained from Life Technologies, Inc., P. O. Box 6009, Gaithersburg, MD 20897. All Sport vectors contain an ampicillin resistance gene and may be transformed into E. coli strain DH10B, also available from Life Technologies. (See, for instance, Gruber, C. E., et al., Focus 15:59 (1993).) Vector lacmid BA (Bento Soares, Columbia University, NY) contains an ampicillin resistance gene and can be transformed into E. coli strain XL-1 Blue. Vector pCR[®]2.1, which is available from Invitrogen, 1600 Faraday Avenue, Carlsbad, CA 92008, contains an ampicillin resistance gene and may be transformed into E. coli strain DH10B, available from Life Technologies. (See, for instance, Clark, J. M., Nuc. Acids Res. 16:9677-9686 (1988) and Mead, D. et al., Bio/Technology 9: (1991).) Preferably, a polynucleotide of the present invention does not comprise the phage vector sequences identified for the particular clone in Table 1, as well as the corresponding plasmid vector sequences designated above.

The deposited material in the sample assigned the ATCC Deposit Number cited in Table 1 for any given cDNA clone also may contain one or more additional plasmids, each comprising a cDNA clone different from that given clone. Thus, deposits sharing the same ATCC Deposit Number contain at least a plasmid for each cDNA clone identified in Table 1. Typically, each ATCC deposit sample cited in Table 1 comprises a mixture of approximately equal amounts (by weight) of about 50 plasmid DNAs, each containing a different cDNA clone; but such a deposit sample

may include plasmids for more or less than 50 cDNA clones, up to about 500 cDNA clones.

Two approaches can be used to isolate a particular clone from the deposited sample of plasmid DNAs cited for that clone in Table 1. First, a plasmid is directly
5 isolated by screening the clones using a polynucleotide probe corresponding to SEQ ID NO:X.

Particularly, a specific polynucleotide with 30-40 nucleotides is synthesized using an Applied Biosystems DNA synthesizer according to the sequence reported. The oligonucleotide is labeled, for instance, with ^{32}P - γ -ATP using T4 polynucleotide
10 kinase and purified according to routine methods. (E.g., Maniatis et al., Molecular Cloning: A Laboratory Manual, Cold Spring Harbor Press, Cold Spring, NY (1982).) The plasmid mixture is transformed into a suitable host, as indicated above (such as XL-1 Blue (Stratagene)) using techniques known to those of skill in the art, such as those provided by the vector supplier or in related publications or patents cited above.
15 The transformants are plated on 1.5% agar plates (containing the appropriate selection agent, e.g., ampicillin) to a density of about 150 transformants (colonies) per plate. These plates are screened using Nylon membranes according to routine methods for bacterial colony screening (e.g., Sambrook et al., Molecular Cloning: A Laboratory Manual, 2nd Edit., (1989), Cold Spring Harbor Laboratory Press, pages 1.93 to
20 1.104), or other techniques known to those of skill in the art.

Alternatively, two primers of 17-20 nucleotides derived from both ends of the SEQ ID NO:X (i.e., within the region of SEQ ID NO:X bounded by the 5' NT and the 3' NT of the clone defined in Table 1) are synthesized and used to amplify the desired cDNA using the deposited cDNA plasmid as a template. The polymerase
25 chain reaction is carried out under routine conditions, for instance, in 25 μl of reaction mixture with 0.5 μg of the above cDNA template. A convenient reaction mixture is 1.5-5 mM MgCl_2 , 0.01% (w/v) gelatin, 20 μM each of dATP, dCTP, dGTP, dTTP, 25 pmol of each primer and 0.25 Unit of Taq polymerase. Thirty five cycles of PCR (denaturation at 94 degree C for 1 min; annealing at 55 degree C for 1 min; elongation
30 at 72 degree C for 1 min) are performed with a Perkin-Elmer Cetus automated thermal cycler. The amplified product is analyzed by agarose gel electrophoresis and

the DNA band with expected molecular weight is excised and purified. The PCR product is verified to be the selected sequence by subcloning and sequencing the DNA product.

Several methods are available for the identification of the 5' or 3' non-coding portions of a gene which may not be present in the deposited clone. These methods include but are not limited to, filter probing, clone enrichment using specific probes, and protocols similar or identical to 5' and 3' "RACE" protocols which are well known in the art. For instance, a method similar to 5' RACE is available for generating the missing 5' end of a desired full-length transcript. (Fromont-Racine et al., Nucleic Acids Res. 21(7):1683-1684 (1993).)

Briefly, a specific RNA oligonucleotide is ligated to the 5' ends of a population of RNA presumably containing full-length gene RNA transcripts. A primer set containing a primer specific to the ligated RNA oligonucleotide and a primer specific to a known sequence of the gene of interest is used to PCR amplify the 5' portion of the desired full-length gene. This amplified product may then be sequenced and used to generate the full length gene.

This above method starts with total RNA isolated from the desired source, although poly-A+ RNA can be used. The RNA preparation can then be treated with phosphatase if necessary to eliminate 5' phosphate groups on degraded or damaged RNA which may interfere with the later RNA ligase step. The phosphatase should then be inactivated and the RNA treated with tobacco acid pyrophosphatase in order to remove the cap structure present at the 5' ends of messenger RNAs. This reaction leaves a 5' phosphate group at the 5' end of the cap cleaved RNA which can then be ligated to an RNA oligonucleotide using T4 RNA ligase.

This modified RNA preparation is used as a template for first strand cDNA synthesis using a gene specific oligonucleotide. The first strand synthesis reaction is used as a template for PCR amplification of the desired 5' end using a primer specific to the ligated RNA oligonucleotide and a primer specific to the known sequence of the gene of interest. The resultant product is then sequenced and analyzed to confirm that the 5' end sequence belongs to the desired gene.

Example 2: Isolation of Genomic Clones Corresponding to a Polynucleotide

A human genomic P1 library (Genomic Systems, Inc.) is screened by PCR using primers selected for the cDNA sequence corresponding to SEQ ID NO:X., according to the method described in Example 1. (See also, Sambrook.)

5

Example 3: Tissue Distribution of Polypeptide

Tissue distribution of mRNA expression of polynucleotides of the present invention is determined using protocols for Northern blot analysis, described by, among others, Sambrook et al. For example, a cDNA probe produced by the method described in Example 1 is labeled with P³² using the rediprime™ DNA labeling system (Amersham Life Science), according to manufacturer's instructions. After labeling, the probe is purified using CHROMA SPIN-100™ column (Clontech Laboratories, Inc.), according to manufacturer's protocol number PT1200-1. The purified labeled probe is then used to examine various human tissues for mRNA expression.

10
15

Multiple Tissue Northern (MTN) blots containing various human tissues (H) or human immune system tissues (IM) (Clontech) are examined with the labeled probe using ExpressHyb™ hybridization solution (Clontech) according to manufacturer's protocol number PT1190-1. Following hybridization and washing, the blots are mounted and exposed to film at -70 degree C overnight, and the films developed according to standard procedures.

20

Example 4: Chromosomal Mapping of the Polynucleotides

An oligonucleotide primer set is designed according to the sequence at the 5' end of SEQ ID NO:X. This primer preferably spans about 100 nucleotides. This primer set is then used in a polymerase chain reaction under the following set of conditions : 30 seconds, 95 degree C; 1 minute, 56 degree C; 1 minute, 70 degree C. This cycle is repeated 32 times followed by one 5 minute cycle at 70 degree C. Human, mouse, and hamster DNA is used as template in addition to a somatic cell hybrid panel containing individual chromosomes or chromosome fragments (Bios, Inc). The reactions is analyzed on either 8% polyacrylamide gels or 3.5 % agarose

25
30

gels. Chromosome mapping is determined by the presence of an approximately 100 bp PCR fragment in the particular somatic cell hybrid.

Example 5: Bacterial Expression of a Polypeptide

5 A polynucleotide encoding a polypeptide of the present invention is amplified using PCR oligonucleotide primers corresponding to the 5' and 3' ends of the DNA sequence, as outlined in Example 1, to synthesize insertion fragments. The primers used to amplify the cDNA insert should preferably contain restriction sites, such as BamHI and XbaI, at the 5' end of the primers in order to clone the amplified product
10 into the expression vector. For example, BamHI and XbaI correspond to the restriction enzyme sites on the bacterial expression vector pQE-9. (Qiagen, Inc., Chatsworth, CA). This plasmid vector encodes antibiotic resistance (Amp^r), a bacterial origin of replication (ori), an IPTG-regulatable promoter/operator (P/O), a ribosome binding site (RBS), a 6-histidine tag (6-His), and restriction enzyme cloning
15 sites.

The pQE-9 vector is digested with BamHI and XbaI and the amplified fragment is ligated into the pQE-9 vector maintaining the reading frame initiated at the bacterial RBS. The ligation mixture is then used to transform the E. coli strain M15/rep4 (Qiagen, Inc.) which contains multiple copies of the plasmid pREP4, which
20 expresses the lacI repressor and also confers kanamycin resistance (Kan^r). Transformants are identified by their ability to grow on LB plates and ampicillin/kanamycin resistant colonies are selected. Plasmid DNA is isolated and confirmed by restriction analysis.

Clones containing the desired constructs are grown overnight (O/N) in liquid
25 culture in LB media supplemented with both Amp (100 ug/ml) and Kan (25 ug/ml). The O/N culture is used to inoculate a large culture at a ratio of 1:100 to 1:250. The cells are grown to an optical density 600 (O.D.⁶⁰⁰) of between 0.4 and 0.6. IPTG (Isopropyl-B-D-thiogalacto pyranoside) is then added to a final concentration of 1 mM. IPTG induces by inactivating the lacI repressor, clearing the P/O leading to
30 increased gene expression.

Cells are grown for an extra 3 to 4 hours. Cells are then harvested by centrifugation (20 mins at 6000Xg). The cell pellet is solubilized in the chaotropic agent 6 Molar Guanidine HCl by stirring for 3-4 hours at 4 degree C. The cell debris is removed by centrifugation, and the supernatant containing the polypeptide is loaded
5 onto a nickel-nitrilo-tri-acetic acid ("Ni-NTA") affinity resin column (available from QIAGEN, Inc., *supra*). Proteins with a 6 x His tag bind to the Ni-NTA resin with high affinity and can be purified in a simple one-step procedure (for details see: The QIAexpressionist (1995) QIAGEN, Inc., *supra*).

Briefly, the supernatant is loaded onto the column in 6 M guanidine-HCl, pH
10 8, the column is first washed with 10 volumes of 6 M guanidine-HCl, pH 8, then washed with 10 volumes of 6 M guanidine-HCl pH 6, and finally the polypeptide is eluted with 6 M guanidine-HCl, pH 5.

The purified protein is then renatured by dialyzing it against phosphate-buffered saline (PBS) or 50 mM Na-acetate, pH 6 buffer plus 200 mM NaCl.
15 Alternatively, the protein can be successfully refolded while immobilized on the Ni-NTA column. The recommended conditions are as follows: renature using a linear 6M-1M urea gradient in 500 mM NaCl, 20% glycerol, 20 mM Tris/HCl pH 7.4, containing protease inhibitors. The renaturation should be performed over a period of 1.5 hours or more. After renaturation the proteins are eluted by the addition of 250
20 mM imidazole. Imidazole is removed by a final dialyzing step against PBS or 50 mM sodium acetate pH 6 buffer plus 200 mM NaCl. The purified protein is stored at 4 degree C or frozen at -80 degree C.

In addition to the above expression vector, the present invention further includes an expression vector comprising phage operator and promoter elements
25 operatively linked to a polynucleotide of the present invention, called pHE4a. (ATCC Accession Number 209645, deposited on February 25, 1998.) This vector contains:
1) a neomycinphosphotransferase gene as a selection marker, 2) an E. coli origin of replication, 3) a T5 phage promoter sequence, 4) two lac operator sequences, 5) a Shine-Delgarno sequence, and 6) the lactose operon repressor gene (lacIq). The
30 origin of replication (oriC) is derived from pUC19 (LTI, Gaithersburg, MD). The promoter sequence and operator sequences are made synthetically.

DNA can be inserted into the pHEa by restricting the vector with NdeI and XbaI, BamHI, XhoI. or Asp718, running the restricted product on a gel, and isolating the larger fragment (the stuffer fragment should be about 310 base pairs). The DNA insert is generated according to the PCR protocol described in Example 1, using PCR
5 primers having restriction sites for NdeI (5' primer) and XbaI, BamHI, XhoI, or Asp718 (3' primer). The PCR insert is gel purified and restricted with compatible enzymes. The insert and vector are ligated according to standard protocols.

The engineered vector could easily be substituted in the above protocol to express protein in a bacterial system.

10

Example 6: Purification of a Polypeptide from an Inclusion Body

The following alternative method can be used to purify a polypeptide expressed in *E coli* when it is present in the form of inclusion bodies. Unless otherwise specified, all of the following steps are conducted at 4-10 degree C.

15 Upon completion of the production phase of the *E. coli* fermentation, the cell culture is cooled to 4-10 degree C and the cells harvested by continuous centrifugation at 15,000 rpm (Heraeus Sepatech). On the basis of the expected yield of protein per unit weight of cell paste and the amount of purified protein required, an appropriate amount of cell paste, by weight, is suspended in a buffer solution
20 containing 100 mM Tris, 50 mM EDTA, pH 7.4. The cells are dispersed to a homogeneous suspension using a high shear mixer.

The cells are then lysed by passing the solution through a microfluidizer (Microfluidics, Corp. or APV Gaulin, Inc.) twice at 4000-6000 psi. The homogenate is then mixed with NaCl solution to a final concentration of 0.5 M NaCl, followed by
25 centrifugation at 7000 xg for 15 min. The resultant pellet is washed again using 0.5M NaCl, 100 mM Tris, 50 mM EDTA, pH 7.4.

The resulting washed inclusion bodies are solubilized with 1.5 M guanidine hydrochloride (GuHCl) for 2-4 hours. After 7000 xg centrifugation for 15 min., the pellet is discarded and the polypeptide containing supernatant is incubated at 4 degree
30 C overnight to allow further GuHCl extraction.

Following high speed centrifugation (30,000 xg) to remove insoluble particles, the GuHCl solubilized protein is refolded by quickly mixing the GuHCl extract with 20 volumes of buffer containing 50 mM sodium, pH 4.5, 150 mM NaCl, 2 mM EDTA by vigorous stirring. The refolded diluted protein solution is kept at 4 degree C
5 without mixing for 12 hours prior to further purification steps.

To clarify the refolded polypeptide solution, a previously prepared tangential filtration unit equipped with 0.16 um membrane filter with appropriate surface area (e.g., Filtron), equilibrated with 40 mM sodium acetate, pH 6.0 is employed. The filtered sample is loaded onto a cation exchange resin (e.g., Poros HS-50, Perseptive
10 Biosystems). The column is washed with 40 mM sodium acetate, pH 6.0 and eluted with 250 mM, 500 mM, 1000 mM, and 1500 mM NaCl in the same buffer, in a stepwise manner. The absorbance at 280 nm of the effluent is continuously monitored. Fractions are collected and further analyzed by SDS-PAGE.

Fractions containing the polypeptide are then pooled and mixed with 4
15 volumes of water. The diluted sample is then loaded onto a previously prepared set of tandem columns of strong anion (Poros HQ-50, Perseptive Biosystems) and weak anion (Poros CM-20, Perseptive Biosystems) exchange resins. The columns are equilibrated with 40 mM sodium acetate, pH 6.0. Both columns are washed with 40 mM sodium acetate, pH 6.0, 200 mM NaCl. The CM-20 column is then eluted using
20 a 10 column volume linear gradient ranging from 0.2 M NaCl, 50 mM sodium acetate, pH 6.0 to 1.0 M NaCl, 50 mM sodium acetate, pH 6.5. Fractions are collected under constant A_{280} monitoring of the effluent. Fractions containing the polypeptide (determined, for instance, by 16% SDS-PAGE) are then pooled.

The resultant polypeptide should exhibit greater than 95% purity after the
25 above refolding and purification steps. No major contaminant bands should be observed from Commassie blue stained 16% SDS-PAGE gel when 5 ug of purified protein is loaded. The purified protein can also be tested for endotoxin/LPS contamination, and typically the LPS content is less than 0.1 ng/ml according to LAL assays.

Example 7: Cloning and Expression of a Polypeptide in a Baculovirus Expression System

In this example, the plasmid shuttle vector pA2 is used to insert a polynucleotide into a baculovirus to express a polypeptide. This expression vector
5 contains the strong polyhedrin promoter of the *Autographa californica* nuclear polyhedrosis virus (AcMNPV) followed by convenient restriction sites such as BamHI, Xba I and Asp718. The polyadenylation site of the simian virus 40 ("SV40") is used for efficient polyadenylation. For easy selection of recombinant virus, the plasmid contains the beta-galactosidase gene from *E. coli* under control of a weak
10 *Drosophila* promoter in the same orientation, followed by the polyadenylation signal of the polyhedrin gene. The inserted genes are flanked on both sides by viral sequences for cell-mediated homologous recombination with wild-type viral DNA to generate a viable virus that express the cloned polynucleotide.

Many other baculovirus vectors can be used in place of the vector above, such
15 as pAc373, pVL941, and pAcIM1, as one skilled in the art would readily appreciate, as long as the construct provides appropriately located signals for transcription, translation, secretion and the like, including a signal peptide and an in-frame AUG as required. Such vectors are described, for instance, in Luckow et al., *Virology* 170:31-39 (1989).

20 Specifically, the cDNA sequence contained in the deposited clone, including the AUG initiation codon and the naturally associated leader sequence identified in Table 1, is amplified using the PCR protocol described in Example 1. If the naturally occurring signal sequence is used to produce the secreted protein, the pA2 vector does not need a second signal peptide. Alternatively, the vector can be modified (pA2 GP)
25 to include a baculovirus leader sequence, using the standard methods described in Summers et al., "A Manual of Methods for Baculovirus Vectors and Insect Cell Culture Procedures," Texas Agricultural Experimental Station Bulletin No. 1555 (1987).

The amplified fragment is isolated from a 1% agarose gel using a
30 commercially available kit ("Geneclean," BIO 101 Inc., La Jolla, Ca.). The fragment

then is digested with appropriate restriction enzymes and again purified on a 1% agarose gel.

The plasmid is digested with the corresponding restriction enzymes and optionally, can be dephosphorylated using calf intestinal phosphatase, using routine procedures known in the art. The DNA is then isolated from a 1% agarose gel using a commercially available kit ("GeneClean" BIO 101 Inc., La Jolla, Ca.).

The fragment and the dephosphorylated plasmid are ligated together with T4 DNA ligase. *E. coli* HB101 or other suitable *E. coli* hosts such as XL-1 Blue (Stratagene Cloning Systems, La Jolla, CA) cells are transformed with the ligation mixture and spread on culture plates. Bacteria containing the plasmid are identified by digesting DNA from individual colonies and analyzing the digestion product by gel electrophoresis. The sequence of the cloned fragment is confirmed by DNA sequencing.

Five ug of a plasmid containing the polynucleotide is co-transfected with 1.0 ug of a commercially available linearized baculovirus DNA ("BaculoGold™ baculovirus DNA", Pharmingen, San Diego, CA), using the lipofection method described by Felgner et al., Proc. Natl. Acad. Sci. USA 84:7413-7417 (1987). One ug of BaculoGold™ virus DNA and 5 ug of the plasmid are mixed in a sterile well of a microtiter plate containing 50 ul of serum-free Grace's medium (Life Technologies Inc., Gaithersburg, MD). Afterwards, 10 ul Lipofectin plus 90 ul Grace's medium are added, mixed and incubated for 15 minutes at room temperature. Then the transfection mixture is added drop-wise to Sf9 insect cells (ATCC CRL 1711) seeded in a 35 mm tissue culture plate with 1 ml Grace's medium without serum. The plate is then incubated for 5 hours at 27 degrees C. The transfection solution is then removed from the plate and 1 ml of Grace's insect medium supplemented with 10% fetal calf serum is added. Cultivation is then continued at 27 degrees C for four days.

After four days the supernatant is collected and a plaque assay is performed, as described by Summers and Smith, *supra*. An agarose gel with "Blue Gal" (Life Technologies Inc., Gaithersburg) is used to allow easy identification and isolation of gal-expressing clones, which produce blue-stained plaques. (A detailed description of a "plaque assay" of this type can also be found in the user's guide for insect cell

culture and baculovirology distributed by Life Technologies Inc., Gaithersburg, page 9-10.) After appropriate incubation, blue stained plaques are picked with the tip of a micropipettor (e.g., Eppendorf). The agar containing the recombinant viruses is then resuspended in a microcentrifuge tube containing 200 ul of Grace's medium and the suspension containing the recombinant baculovirus is used to infect Sf9 cells seeded in 35 mm dishes. Four days later the supernatants of these culture dishes are harvested and then they are stored at 4 degree C.

To verify the expression of the polypeptide, Sf9 cells are grown in Grace's medium supplemented with 10% heat-inactivated FBS. The cells are infected with the recombinant baculovirus containing the polynucleotide at a multiplicity of infection ("MOI") of about 2. If radiolabeled proteins are desired, 6 hours later the medium is removed and is replaced with SF900 II medium minus methionine and cysteine (available from Life Technologies Inc., Rockville, MD). After 42 hours, 5 uCi of ^{35}S -methionine and 5 uCi ^{35}S -cysteine (available from Amersham) are added. The cells are further incubated for 16 hours and then are harvested by centrifugation. The proteins in the supernatant as well as the intracellular proteins are analyzed by SDS-PAGE followed by autoradiography (if radiolabeled).

Microsequencing of the amino acid sequence of the amino terminus of purified protein may be used to determine the amino terminal sequence of the produced protein.

Example 8: Expression of a Polypeptide in Mammalian Cells

The polypeptide of the present invention can be expressed in a mammalian cell. A typical mammalian expression vector contains a promoter element, which mediates the initiation of transcription of mRNA, a protein coding sequence, and signals required for the termination of transcription and polyadenylation of the transcript. Additional elements include enhancers, Kozak sequences and intervening sequences flanked by donor and acceptor sites for RNA splicing. Highly efficient transcription is achieved with the early and late promoters from SV40, the long terminal repeats (LTRs) from Retroviruses, e.g., RSV, HTLVI, HIVI and the early

promoter of the cytomegalovirus (CMV). However, cellular elements can also be used (e.g., the human actin promoter).

Suitable expression vectors for use in practicing the present invention include, for example, vectors such as pSVL and pMSG (Pharmacia, Uppsala, Sweden),
5 pRSVcat (ATCC 37152), pSV2dhfr (ATCC 37146), pBC12MI (ATCC 67109), pCMVSPORT 2.0, and pCMVSPORT 3.0. Mammalian host cells that could be used include, human Hela, 293, H9 and Jurkat cells, mouse NIH3T3 and C127 cells, Cos 1, Cos 7 and CV1, quail QC1-3 cells, mouse L cells and Chinese hamster ovary (CHO) cells.

10 Alternatively, the polypeptide can be expressed in stable cell lines containing the polynucleotide integrated into a chromosome. The co-transfection with a selectable marker such as dhfr, gpt, neomycin, hygromycin allows the identification and isolation of the transfected cells.

The transfected gene can also be amplified to express large amounts of the
15 encoded protein. The DHFR (dihydrofolate reductase) marker is useful in developing cell lines that carry several hundred or even several thousand copies of the gene of interest. (See, e.g., Alt, F. W., et al., J. Biol. Chem. 253:1357-1370 (1978); Hamlin, J. L. and Ma, C., Biochem. et Biophys. Acta, 1097:107-143 (1990); Page, M. J. and Sydenham, M. A., Biotechnology 9:64-68 (1991).) Another useful selection marker
20 is the enzyme glutamine synthase (GS) (Murphy et al., Biochem J. 227:277-279 (1991); Bebbington et al., Bio/Technology 10:169-175 (1992). Using these markers, the mammalian cells are grown in selective medium and the cells with the highest resistance are selected. These cell lines contain the amplified gene(s) integrated into a chromosome. Chinese hamster ovary (CHO) and NSO cells are often used for the
25 production of proteins.

Derivatives of the plasmid pSV2-dhfr (ATCC Accession No. 37146), the expression vectors pC4 (ATCC Accession No. 209646) and pC6 (ATCC Accession No. 209647) contain the strong promoter (LTR) of the Rous Sarcoma Virus (Cullen et al., Molecular and Cellular Biology, 438-447 (March, 1985)) plus a fragment of the
30 CMV-enhancer (Boshart et al., Cell 41:521-530 (1985).) Multiple cloning sites, e.g., with the restriction enzyme cleavage sites BamHI, XbaI and Asp718, facilitate the

cloning of the gene of interest. The vectors also contain the 3' intron, the polyadenylation and termination signal of the rat preproinsulin gene, and the mouse DHFR gene under control of the SV40 early promoter.

Specifically, the plasmid pC6, for example, is digested with appropriate
5 restriction enzymes and then dephosphorylated using calf intestinal phosphates by procedures known in the art. The vector is then isolated from a 1% agarose gel.

A polynucleotide of the present invention is amplified according to the protocol outlined in Example 1. If the naturally occurring signal sequence is used to produce the secreted protein, the vector does not need a second signal peptide.
10 Alternatively, if the naturally occurring signal sequence is not used, the vector can be modified to include a heterologous signal sequence. (See, e.g., WO 96/34891.)

The amplified fragment is isolated from a 1% agarose gel using a commercially available kit ("Geneclean," BIO 101 Inc., La Jolla, Ca.). The fragment then is digested with appropriate restriction enzymes and again purified on a 1%
15 agarose gel.

The amplified fragment is then digested with the same restriction enzyme and purified on a 1% agarose gel. The isolated fragment and the dephosphorylated vector are then ligated with T4 DNA ligase. *E. coli* HB101 or XL-1 Blue cells are then transformed and bacteria are identified that contain the fragment inserted into plasmid
20 pC6 using, for instance, restriction enzyme analysis.

Chinese hamster ovary cells lacking an active DHFR gene is used for transfection. Five µg of the expression plasmid pC6 a pC4 is cotransfected with 0.5 µg of the plasmid pSVneo using lipofectin (Felgner et al., *supra*). The plasmid pSV2-neo contains a dominant selectable marker, the *neo* gene from Tn5 encoding an
25 enzyme that confers resistance to a group of antibiotics including G418. The cells are seeded in alpha minus MEM supplemented with 1 mg/ml G418. After 2 days, the cells are trypsinized and seeded in hybridoma cloning plates (Greiner, Germany) in alpha minus MEM supplemented with 10, 25, or 50 ng/ml of methotrexate plus 1 mg/ml G418. After about 10-14 days single clones are trypsinized and then seeded in
30 6-well petri dishes or 10 ml flasks using different concentrations of methotrexate (50 nM, 100 nM, 200 nM, 400 nM, 800 nM). Clones growing at the highest

concentrations of methotrexate are then transferred to new 6-well plates containing even higher concentrations of methotrexate (1 uM, 2 uM, 5 uM, 10 mM, 20 mM). The same procedure is repeated until clones are obtained which grow at a concentration of 100 - 200 uM. Expression of the desired gene product is analyzed,
5 for instance, by SDS-PAGE and Western blot or by reversed phase HPLC analysis.

Example 9: Protein Fusions

The polypeptides of the present invention are preferably fused to other proteins. These fusion proteins can be used for a variety of applications. For
10 example, fusion of the present polypeptides to His-tag, HA-tag, protein A, IgG domains, and maltose binding protein facilitates purification. (See Example 5; see also EP A 394,827; Traunecker, et al., Nature 331:84-86 (1988).) Similarly, fusion to IgG-1, IgG-3, and albumin increases the halflife time in vivo. Nuclear localization signals fused to the polypeptides of the present invention can target the protein to a
15 specific subcellular localization, while covalent heterodimer or homodimers can increase or decrease the activity of a fusion protein. Fusion proteins can also create chimeric molecules having more than one function. Finally, fusion proteins can increase solubility and/or stability of the fused protein compared to the non-fused protein. All of the types of fusion proteins described above can be made by
20 modifying the following protocol, which outlines the fusion of a polypeptide to an IgG molecule, or the protocol described in Example 5.

Briefly, the human Fc portion of the IgG molecule can be PCR amplified, using primers that span the 5' and 3' ends of the sequence described below. These primers also should have convenient restriction enzyme sites that will facilitate
25 cloning into an expression vector, preferably a mammalian expression vector.

For example, if pC4 (Accession No. 209646) is used, the human Fc portion can be ligated into the BamHI cloning site. Note that the 3' BamHI site should be destroyed. Next, the vector containing the human Fc portion is re-restricted with BamHI, linearizing the vector, and a polynucleotide of the present invention, isolated
30 by the PCR protocol described in Example 1, is ligated into this BamHI site. Note

that the polynucleotide is cloned without a stop codon, otherwise a fusion protein will not be produced.

If the naturally occurring signal sequence is used to produce the secreted protein, pC4 does not need a second signal peptide. Alternatively, if the naturally occurring signal sequence is not used, the vector can be modified to include a heterologous signal sequence. (See, e.g., WO 96/34891.)

Human IgG Fc region:

GGGATCCGGAGCCCAAATCTTCTGACAAAACCTCACACATGCCCACCGTGCCAGCACCTGAATTCGAGG
 10 GTGCACCGTCAGTCTTCTTCTCCCCCAAACCCAAAGGACACCCTCATGATCTCCCGGACTCCTGAGG
 TCACATGCGTGGTGGTGGACGTAAGCCACGAAGACCCTGAGGTCAAGTTCAACTGGTACGTGGACGGCG
 TGGAGGTGCATAATGCCAAGACAAAGCCGCGGGAGGAGCAGTACAACAGCACGTACCGTGTGGTCAGCG
 TCCTCACCGTCTGACACCAGGACTGGCTGAATGGCAAGGAGTACAAGTGCAAGGTCTCCAACAAAGCCC
 TCCCAACCCCCATCGAGAAAACCATCTCCAAAGCCAAAGGGCAGCCCCGAGAACCACAGGTGTACACCC
 15 TGCCCCCATCCCGGGATGAGCTGACCAAGAACCAGGTCAGCCTGACCTGCCTGGTCAAAGGCTTCTATC
 CAAGCGACATCGCCGTGGAGTGGGAGAGCAATGGGCAGCCGGAGAACAACCTACAAGACCACGCCTCCCCG
 TGCTGGACTCCGACGGCTCCTTCTTCTCTACAGCAAGCTCACCGTGACAAGAGCAGGTGGCAGCAGG
 GGAACGTCTTCTCATGCTCCGTGATGCATGAGGCTCTGCACAACCACTACACGCAGAAGAGCCTCTCCC
 TGTCTCCGGGTAAATGAGTGCACGGCCGCGACTCTAGAGGAT (SEQ ID NO:1)

20

Example 10: Production of an Antibody from a Polypeptide

The antibodies of the present invention can be prepared by a variety of methods. (See, Current Protocols, Chapter 2.) As one example of such methods, cells expressing a polypeptide of the present invention is administered to an animal to induce the production of sera containing polyclonal antibodies. In a preferred method, a preparation of the secreted protein is prepared and purified to render it substantially free of natural contaminants. Such a preparation is then introduced into an animal in order to produce polyclonal antisera of greater specific activity.

In the most preferred method, the antibodies of the present invention are monoclonal antibodies (or protein binding fragments thereof). Such monoclonal antibodies can be prepared using hybridoma technology. (Köhler et al., Nature 256:495 (1975); Köhler et al., Eur. J. Immunol. 6:511 (1976); Köhler et al., Eur. J. Immunol. 6:292 (1976); Hammerling et al., in: Monoclonal Antibodies and T-Cell Hybridomas, Elsevier, N.Y., pp. 563-681 (1981).) In general, such procedures

30

involve immunizing an animal (preferably a mouse) with polypeptide or, more preferably, with a secreted polypeptide-expressing cell. Such cells may be cultured in any suitable tissue culture medium; however, it is preferable to culture cells in Earle's modified Eagle's medium supplemented with 10% fetal bovine serum (inactivated at
5 about 56 degrees C), and supplemented with about 10 g/l of nonessential amino acids, about 1,000 U/ml of penicillin, and about 100 ug/ml of streptomycin.

The splenocytes of such mice are extracted and fused with a suitable myeloma cell line. Any suitable myeloma cell line may be employed in accordance with the present invention; however, it is preferable to employ the parent myeloma cell line
10 (SP2O), available from the ATCC. After fusion, the resulting hybridoma cells are selectively maintained in HAT medium, and then cloned by limiting dilution as described by Wands et al. (Gastroenterology 80:225-232 (1981).) The hybridoma cells obtained through such a selection are then assayed to identify clones which secrete antibodies capable of binding the polypeptide.

15 Alternatively, additional antibodies capable of binding to the polypeptide can be produced in a two-step procedure using anti-idiotypic antibodies. Such a method makes use of the fact that antibodies are themselves antigens, and therefore, it is possible to obtain an antibody which binds to a second antibody. In accordance with this method, protein specific antibodies are used to immunize an animal, preferably a
20 mouse. The splenocytes of such an animal are then used to produce hybridoma cells, and the hybridoma cells are screened to identify clones which produce an antibody whose ability to bind to the protein-specific antibody can be blocked by the polypeptide. Such antibodies comprise anti-idiotypic antibodies to the protein-specific antibody and can be used to immunize an animal to induce formation of
25 further protein-specific antibodies.

It will be appreciated that Fab and F(ab')₂ and other fragments of the antibodies of the present invention may be used according to the methods disclosed herein. Such fragments are typically produced by proteolytic cleavage, using
enzymes such as papain (to produce Fab fragments) or pepsin (to produce F(ab')₂
30 fragments). Alternatively, secreted protein-binding fragments can be produced

through the application of recombinant DNA technology or through synthetic chemistry.

For in vivo use of antibodies in humans, it may be preferable to use "humanized" chimeric monoclonal antibodies. Such antibodies can be produced using genetic constructs derived from hybridoma cells producing the monoclonal antibodies described above. Methods for producing chimeric antibodies are known in the art. (See, for review, Morrison, Science 229:1202 (1985); Oi et al., BioTechniques 4:214 (1986); Cabilly et al., U.S. Patent No. 4,816,567; Taniguchi et al., EP 171496; Morrison et al., EP 173494; Neuberger et al., WO 8601533; Robinson et al., WO 8702671; Boulianne et al., Nature 312:643 (1984); Neuberger et al., Nature 314:268 (1985).)

Example 11: Production Of Secreted Protein For High-Throughput Screening Assays

The following protocol produces a supernatant containing a polypeptide to be tested. This supernatant can then be used in the Screening Assays described in Examples 13-20.

First, dilute Poly-D-Lysine (644 587 Boehringer-Mannheim) stock solution (1mg/ml in PBS) 1:20 in PBS (w/o calcium or magnesium 17-516F Biowhittaker) for a working solution of 50ug/ml. Add 200 ul of this solution to each well (24 well plates) and incubate at RT for 20 minutes. Be sure to distribute the solution over each well (note: a 12-channel pipetter may be used with tips on every other channel). Aspirate off the Poly-D-Lysine solution and rinse with 1ml PBS (Phosphate Buffered Saline). The PBS should remain in the well until just prior to plating the cells and plates may be poly-lysine coated in advance for up to two weeks.

Plate 293T cells (do not carry cells past P+20) at 2×10^5 cells/well in .5ml DMEM(Dulbecco's Modified Eagle Medium)(with 4.5 G/L glucose and L-glutamine (12-604F Biowhittaker))/10% heat inactivated FBS(14-503F Biowhittaker)/1x Penstrep(17-602E Biowhittaker). Let the cells grow overnight.

The next day, mix together in a sterile solution basin: 300 ul Lipofectamine (18324-012 Gibco/BRL) and 5ml Optimem I (31985070 Gibco/BRL)/96-well plate.

With a small volume multi-channel pipetter, aliquot approximately 2ug of an expression vector containing a polynucleotide insert, produced by the methods described in Examples 8 or 9, into an appropriately labeled 96-well round bottom plate. With a multi-channel pipetter, add 50ul of the Lipofectamine/Optimem I mixture to each well. Pipette up and down gently to mix. Incubate at RT 15-45 minutes. After about 20 minutes, use a multi-channel pipetter to add 150ul Optimem I to each well. As a control, one plate of vector DNA lacking an insert should be transfected with each set of transfections.

Preferably, the transfection should be performed by tag-teaming the following tasks. By tag-teaming, hands on time is cut in half, and the cells do not spend too much time on PBS. First, person A aspirates off the media from four 24-well plates of cells, and then person B rinses each well with .5-1ml PBS. Person A then aspirates off PBS rinse, and person B, using a 12-channel pipetter with tips on every other channel, adds the 200ul of DNA/Lipofectamine/Optimem I complex to the odd wells first, then to the even wells, to each row on the 24-well plates. Incubate at 37 degrees C for 6 hours.

While cells are incubating, prepare appropriate media, either 1%BSA in DMEM with 1x penstrep, or CHO-5 media (116.6 mg/L of CaCl₂ (anhyd); 0.00130 mg/L CuSO₄·5H₂O; 0.050 mg/L of Fe(NO₃)₃·9H₂O; 0.417 mg/L of FeSO₄·7H₂O; 311.80 mg/L of KCl; 28.64 mg/L of MgCl₂; 48.84 mg/L of MgSO₄; 6995.50 mg/L of NaCl; 2400.0 mg/L of NaHCO₃; 62.50 mg/L of NaH₂PO₄·H₂O; 71.02 mg/L of Na₂HPO₄; .4320 mg/L of ZnSO₄·7H₂O; .002 mg/L of Arachidonic Acid ; 1.022 mg/L of Cholesterol; .070 mg/L of DL-alpha-Tocopherol-Acetate; 0.0520 mg/L of Linoleic Acid; 0.010 mg/L of Linolenic Acid; 0.010 mg/L of Myristic Acid; 0.010 mg/L of Oleic Acid; 0.010 mg/L of Palmitric Acid; 0.010 mg/L of Palmitic Acid; 100 mg/L of Pluronic F-68; 0.010 mg/L of Stearic Acid; 2.20 mg/L of Tween 80; 4551 mg/L of D-Glucose; 130.85 mg/ml of L- Alanine; 147.50 mg/ml of L-Arginine-HCL; 7.50 mg/ml of L-Asparagine-H₂O; 6.65 mg/ml of L-Aspartic Acid; 29.56 mg/ml of L-Cystine-2HCL-H₂O; 31.29 mg/ml of L-Cystine-2HCL; 7.35 mg/ml of L-Glutamic Acid; 365.0 mg/ml of L-Glutamine; 18.75 mg/ml of Glycine; 52.48 mg/ml of L-Histidine-HCL-H₂O; 106.97 mg/ml of L-Isoleucine; 111.45 mg/ml of L-Leucine; 163.75 mg/ml of L-

- Lysine HCL; 32.34 mg/ml of L-Methionine; 68.48 mg/ml of L-Phenylalanine; 40.0 mg/ml of L-Proline; 26.25 mg/ml of L-Serine; 101.05 mg/ml of L-Threonine; 19.22 mg/ml of L-Tryptophan; 91.79 mg/ml of L-Tyrosine-2Na-2H₂O; 99.65 mg/ml of L-Valine; 0.0035 mg/L of Biotin; 3.24 mg/L of D-Ca Pantothenate; 11.78 mg/L of
- 5 Choline Chloride; 4.65 mg/L of Folic Acid; 15.60 mg/L of i-Inositol; 3.02 mg/L of Niacinamide; 3.00 mg/L of Pyridoxal HCL; 0.031 mg/L of Pyridoxine HCL; 0.319 mg/L of Riboflavin; 3.17 mg/L of Thiamine HCL; 0.365 mg/L of Thymidine; and 0.680 mg/L of Vitamin B₁₂; 25 mM of HEPES Buffer; 2.39 mg/L of Na Hypoxanthine; 0.105 mg/L of Lipoic Acid; 0.081 mg/L of Sodium Putrescine-2HCL;
- 10 55.0 mg/L of Sodium Pyruvate; 0.0067 mg/L of Sodium Selenite; 20uM of Ethanolamine; 0.122 mg/L of Ferric Citrate; 41.70 mg/L of Methyl-B-Cyclodextrin complexed with Linoleic Acid; 33.33 mg/L of Methyl-B-Cyclodextrin complexed with Oleic Acid; and 10 mg/L of Methyl-B-Cyclodextrin complexed with Retinal) with 2mm glutamine and 1x penstrep. (BSA (81-068-3 Bayer) 100gm dissolved in 1L
- 15 DMEM for a 10% BSA stock solution). Filter the media and collect 50 ul for endotoxin assay in 15ml polystyrene conical.

The transfection reaction is terminated, preferably by tag-teaming, at the end of the incubation period. Person A aspirates off the transfection media, while person B adds 1.5ml appropriate media to each well. Incubate at 37 degrees C for 45 or 72

20 hours depending on the media used: 1%BSA for 45 hours or CHO-5 for 72 hours.

On day four, using a 300ul multichannel pipetter, aliquot 600ul in one 1ml deep well plate and the remaining supernatant into a 2ml deep well. The supernatants from each well can then be used in the assays described in Examples 13-20.

It is specifically understood that when activity is obtained in any of the assays

25 described below using a supernatant, the activity originates from either the polypeptide directly (e.g., as a secreted protein) or by the polypeptide inducing expression of other proteins, which are then secreted into the supernatant. Thus, the invention further provides a method of identifying the protein in the supernatant characterized by an activity in a particular assay.

30

Example 12: Construction of GAS Reporter Construct

One signal transduction pathway involved in the differentiation and proliferation of cells is called the Jaks-STATs pathway. Activated proteins in the Jaks-STATs pathway bind to gamma activation site "GAS" elements or interferon-sensitive responsive element ("ISRE"), located in the promoter of many genes. The binding of a protein to these elements alter the expression of the associated gene.

GAS and ISRE elements are recognized by a class of transcription factors called Signal Transducers and Activators of Transcription, or "STATs." There are six members of the STATs family. Stat1 and Stat3 are present in many cell types, as is Stat2 (as response to IFN-alpha is widespread). Stat4 is more restricted and is not in many cell types though it has been found in T helper class I, cells after treatment with IL-12. Stat5 was originally called mammary growth factor, but has been found at higher concentrations in other cells including myeloid cells. It can be activated in tissue culture cells by many cytokines.

The STATs are activated to translocate from the cytoplasm to the nucleus upon tyrosine phosphorylation by a set of kinases known as the Janus Kinase ("Jaks") family. Jaks represent a distinct family of soluble tyrosine kinases and include Tyk2, Jak1, Jak2, and Jak3. These kinases display significant sequence similarity and are generally catalytically inactive in resting cells.

The Jaks are activated by a wide range of receptors summarized in the Table below. (Adapted from review by Schidler and Darnell, Ann. Rev. Biochem. 64:621-51 (1995).) A cytokine receptor family, capable of activating Jaks, is divided into two groups: (a) Class 1 includes receptors for IL-2, IL-3, IL-4, IL-6, IL-7, IL-9, IL-11, IL-12, IL-15, Epo, PRL, GH, G-CSF, GM-CSF, LIF, CNTF, and thrombopoietin; and (b) Class 2 includes IFN-a, IFN-g, and IL-10. The Class 1 receptors share a conserved cysteine motif (a set of four conserved cysteines and one tryptophan) and a WSXWS motif (a membrane proximal region encoding Trp-Ser-Xxx-Trp-Ser (SEQ ID NO:2)).

Thus, on binding of a ligand to a receptor, Jaks are activated, which in turn activate STATs, which then translocate and bind to GAS elements. This entire process is encompassed in the Jaks-STATs signal transduction pathway.

Therefore, activation of the Jaks-STATs pathway, reflected by the binding of the GAS or the ISRE element, can be used to indicate proteins involved in the

proliferation and differentiation of cells. For example, growth factors and cytokines are known to activate the Jaks-STATs pathway. (See Table below.) Thus, by using GAS elements linked to reporter molecules, activators of the Jaks-STATs pathway can be identified.

	<u>Ligand</u>	<u>tyk2</u>	<u>JAKs</u> <u>Jak1</u>	<u>Jak2</u>	<u>Jak3</u>	<u>STATS</u>	<u>GAS(elements) or ISRE</u>
	<u>IFN family</u>						
5	IFN-a/B	+	+	-	-	1,2,3	ISRE
	IFN-g		+	+	-	1	GAS (IRF1>Lys6>IFP)
	Il-10	+	?	?	-	1,3	
	<u>gp130 family</u>						
10	IL-6 (Pleiotrophic)	+	+	+	?	1,3	GAS (IRF1>Lys6>IFP)
	Il-11(Pleiotrophic)	?	+	?	?	1,3	
	OnM(Pleiotrophic)	?	+	+	?	1,3	
	LIF(Pleiotrophic)	?	+	+	?	1,3	
	CNTF(Pleiotrophic)	-/+	+	+	?	1,3	
15	G-CSF(Pleiotrophic)	?	+	?	?	1,3	
	IL-12(Pleiotrophic)	+	-	+	+	1,3	
	<u>g-C family</u>						
20	IL-2 (lymphocytes)	-	+	-	+	1,3,5	GAS
	IL-4 (lymph/myeloid)	-	+	-	+	6	GAS (IRF1 = IFP >>Ly6)(IgH)
	IL-7 (lymphocytes)	-	+	-	+	5	GAS
	IL-9 (lymphocytes)	-	+	-	+	5	GAS
	IL-13 (lymphocyte)	-	+	?	?	6	GAS
	IL-15	?	+	?	+	5	GAS
25	<u>gp140 family</u>						
	IL-3 (myeloid)	-	-	+	-	5	GAS (IRF1>IFP>>Ly6)
	IL-5 (myeloid)	-	-	+	-	5	GAS
	GM-CSF (myeloid)	-	-	+	-	5	GAS
30	<u>Growth hormone family</u>						
	GH	?	-	+	-	5	
	PRL	?	+/-	+	-	1,3,5	
	EPO	?	-	+	-	5	GAS(B-CAS>IRF1=IFP>>Ly6)
35	<u>Receptor Tyrosine Kinases</u>						
	EGF	?	+	+	-	1,3	GAS (IRF1)
	PDGF	?	+	+	-	1,3	
	CSF-1	?	+	+	-	1,3	GAS (not IRF1)
40							

To construct a synthetic GAS containing promoter element, which is used in the Biological Assays described in Examples 13-14, a PCR based strategy is employed to generate a GAS-SV40 promoter sequence. The 5' primer contains four tandem copies of the GAS binding site found in the IRF1 promoter and previously demonstrated to bind STATs upon induction with a range of cytokines (Rothman et al., Immunity 1:457-468 (1994).), although other GAS or ISRE elements can be used instead. The 5' primer also contains 18bp of sequence complementary to the SV40 early promoter sequence and is flanked with an XhoI site. The sequence of the 5' primer is:

5' : GCGCCTCGAGATTTCCTCCGAAATCTAGATTTCCTCCGAAATGATTTCCTCCGAAATGATTTCCTCCGAAATATCTGCCATCTCAATTAG : 3' (SEQ ID NO:3)

The downstream primer is complementary to the SV40 promoter and is flanked with a Hind III site: 5' : GCGGCAAGCTTTTTTGCAAAGCCTAGGC : 3' (SEQ ID NO:4)

PCR amplification is performed using the SV40 promoter template present in the B-gal:promoter plasmid obtained from Clontech. The resulting PCR fragment is digested with XhoI/Hind III and subcloned into BLSK2-. (Stratagene.) Sequencing with forward and reverse primers confirms that the insert contains the following sequence:

5' : CTCGAGATTTCCTCCGAAATCTAGATTTCCTCCGAAATGATTTCCTCCGAAATGATTTCCTCCGAAATATCTGCCATCTCAATTAGTCAGCAACCATAGTCCCGCCCCTAACTCCGCCCATCCCGCCCCTAACTCCGCCCAGTTCCGCCCATTCTCCGCCCCATGGCTGACTAATTTTTTTTATTTATGCAGAGGCCGAGGCCGCCTCGGCCTCTGAGCTATTCCAGAAGTAGTGAGGAGGCTTTTTTGGAGGCCTAGGCTTTTGCAAAAAGCTT : 3' (SEQ ID NO:5)

With this GAS promoter element linked to the SV40 promoter, a GAS:SEAP2 reporter construct is next engineered. Here, the reporter molecule is a secreted alkaline phosphatase, or "SEAP." Clearly, however, any reporter molecule can be instead of SEAP, in this or in any of the other Examples. Well known reporter molecules that can be used instead of SEAP include chloramphenicol

acetyltransferase (CAT), luciferase, alkaline phosphatase, B-galactosidase, green fluorescent protein (GFP), or any protein detectable by an antibody.

The above sequence confirmed synthetic GAS-SV40 promoter element is subcloned into the pSEAP-Promoter vector obtained from Clontech using HindIII and XhoI, effectively replacing the SV40 promoter with the amplified GAS:SV40 promoter element, to create the GAS-SEAP vector. However, this vector does not contain a neomycin resistance gene, and therefore, is not preferred for mammalian expression systems.

Thus, in order to generate mammalian stable cell lines expressing the GAS-SEAP reporter, the GAS-SEAP cassette is removed from the GAS-SEAP vector using SalI and NotI, and inserted into a backbone vector containing the neomycin resistance gene, such as pGFP-1 (Clontech), using these restriction sites in the multiple cloning site, to create the GAS-SEAP/Neo vector. Once this vector is transfected into mammalian cells, this vector can then be used as a reporter molecule for GAS binding as described in Examples 13-14.

Other constructs can be made using the above description and replacing GAS with a different promoter sequence. For example, construction of reporter molecules containing NFK-B and EGR promoter sequences are described in Examples 15 and 16. However, many other promoters can be substituted using the protocols described in these Examples. For instance, SRE, IL-2, NFAT, or Osteocalcin promoters can be substituted, alone or in combination (e.g., GAS/NF-KB/EGR, GAS/NF-KB, IL-2/NFAT, or NF-KB/GAS). Similarly, other cell lines can be used to test reporter construct activity, such as HELA (epithelial), HUVEC (endothelial), Reh (B-cell), Saos-2 (osteoblast), HUVAC (aortic), or Cardiomyocyte.

25

Example 13: High-Throughput Screening Assay for T-cell Activity.

The following protocol is used to assess T-cell activity by identifying factors, and determining whether supernate containing a polypeptide of the invention proliferates and/or differentiates T-cells. T-cell activity is assessed using the GAS/SEAP/Neo construct produced in Example 12. Thus, factors that increase SEAP activity indicate the ability to activate the Jaks-STATS signal transduction pathway.

30

The T-cell used in this assay is Jurkat T-cells (ATCC Accession No. TIB-152), although Molt-3 cells (ATCC Accession No. CRL-1552) and Molt-4 cells (ATCC Accession No. CRL-1582) cells can also be used.

5 Jurkat T-cells are lymphoblastic CD4+ Th1 helper cells. In order to generate stable cell lines, approximately 2 million Jurkat cells are transfected with the GAS-SEAP/neo vector using DMRIE-C (Life Technologies)(transfection procedure described below). The transfected cells are seeded to a density of approximately 20,000 cells per well and transfectants resistant to 1 mg/ml gentamicin selected. Resistant colonies are expanded and then tested for their response to increasing
10 concentrations of interferon gamma. The dose response of a selected clone is demonstrated.

Specifically, the following protocol will yield sufficient cells for 75 wells containing 200 ul of cells. Thus, it is either scaled up, or performed in multiple to generate sufficient cells for multiple 96 well plates. Jurkat cells are maintained in
15 RPMI + 10% serum with 1%Pen-Strep. Combine 2.5 mls of OPTI-MEM (Life Technologies) with 10 ug of plasmid DNA in a T25 flask. Add 2.5 ml OPTI-MEM containing 50 ul of DMRIE-C and incubate at room temperature for 15-45 mins.

During the incubation period, count cell concentration, spin down the required number of cells (10^7 per transfection), and resuspend in OPTI-MEM to a final
20 concentration of 10^7 cells/ml. Then add 1ml of 1×10^7 cells in OPTI-MEM to T25 flask and incubate at 37 degrees C for 6 hrs. After the incubation, add 10 ml of RPMI + 15% serum.

The Jurkat:GAS-SEAP stable reporter lines are maintained in RPMI + 10% serum, 1 mg/ml Gentamicin, and 1% Pen-Strep. These cells are treated with
25 supernatants containing polypeptides of the invention and/or induced polypeptides of the invention as produced by the protocol described in Example 11.

On the day of treatment with the supernatant, the cells should be washed and resuspended in fresh RPMI + 10% serum to a density of 500,000 cells per ml. The exact number of cells required will depend on the number of supernatants being
30 screened. For one 96 well plate, approximately 10 million cells (for 10 plates, 100 million cells) are required.

Transfer the cells to a triangular reservoir boat, in order to dispense the cells into a 96 well dish, using a 12 channel pipette. Using a 12 channel pipette, transfer 200 ul of cells into each well (therefore adding 100, 000 cells per well).

After all the plates have been seeded, 50 ul of the supernatants are transferred
5 directly from the 96 well plate containing the supernatants into each well using a 12 channel pipette. In addition, a dose of exogenous interferon gamma (0.1, 1.0, 10 ng) is added to wells H9, H10, and H11 to serve as additional positive controls for the assay.

The 96 well dishes containing Jurkat cells treated with supernatants are placed
10 in an incubator for 48 hrs (note: this time is variable between 48-72 hrs). 35 ul samples from each well are then transferred to an opaque 96 well plate using a 12 channel pipette. The opaque plates should be covered (using sellophene covers) and stored at -20 degrees C until SEAP assays are performed according to Example 17. The plates containing the remaining treated cells are placed at 4 degrees C and serve
15 as a source of material for repeating the assay on a specific well if desired.

As a positive control, 100 Unit/ml interferon gamma can be used which is known to activate Jurkat T cells. Over 30 fold induction is typically observed in the positive control wells.

The above protocol may be used in the generation of both transient, as well as,
20 stable transfected cells, which would be apparent to those of skill in the art.

Example 14: High-Throughput Screening Assay Identifying Myeloid Activity

The following protocol is used to assess myeloid activity by determining whether polypeptides of the invention proliferates and/or differentiates myeloid cells.
25 Myeloid cell activity is assessed using the GAS/SEAP/Neo construct produced in Example 12. Thus, factors that increase SEAP activity indicate the ability to activate the Jaks-STATS signal transduction pathway. The myeloid cell used in this assay is U937, a pre-monocyte cell line, although TF-1, HL60, or KG1 can be used.

To transiently transfect U937 cells with the GAS/SEAP/Neo construct
30 produced in Example 12, a DEAE-Dextran method (Kharbanda et. al., 1994, Cell Growth & Differentiation, 5:259-265) is used. First, harvest 2×10^7 U937 cells and

wash with PBS. The U937 cells are usually grown in RPMI 1640 medium containing 10% heat-inactivated fetal bovine serum (FBS) supplemented with 100 units/ml penicillin and 100 mg/ml streptomycin.

Next, suspend the cells in 1 ml of 20 mM Tris-HCl (pH 7.4) buffer containing
5 0.5 mg/ml DEAE-Dextran, 8 ug GAS-SEAP2 plasmid DNA, 140 mM NaCl, 5 mM KCl, 375 uM Na₂HPO₄·7H₂O, 1 mM MgCl₂, and 675 uM CaCl₂. Incubate at 37 degrees C for 45 min.

Wash the cells with RPMI 1640 medium containing 10% FBS and then resuspend in 10 ml complete medium and incubate at 37 degrees C for 36 hr.

10 The GAS-SEAP/U937 stable cells are obtained by growing the cells in 400 ug/ml G418. The G418-free medium is used for routine growth but every one to two months, the cells should be re-grown in 400 ug/ml G418 for couple of passages.

These cells are tested by harvesting 1×10^8 cells (this is enough for ten 96-well plates assay) and wash with PBS. Suspend the cells in 200 ml above described
15 growth medium, with a final density of 5×10^5 cells/ml. Plate 200 ul cells per well in the 96-well plate (or 1×10^5 cells/well).

Add 50 ul of the supernatant prepared by the protocol described in Example 11. Incubate at 37 degrees C for 48 to 72 hr. As a positive control, 100 Unit/ml interferon gamma can be used which is known to activate U937 cells. Over 30 fold
20 induction is typically observed in the positive control wells. SEAP assay the supernatant according to the protocol described in Example 17.

Example 15: High-Throughput Screening Assay Identifying Neuronal Activity.

When cells undergo differentiation and proliferation, a group of genes are
25 activated through many different signal transduction pathways. One of these genes, EGR1 (early growth response gene 1), is induced in various tissues and cell types upon activation. The promoter of EGR1 is responsible for such induction. Using the EGR1 promoter linked to reporter molecules, activation of cells can be assessed.

Particularly, the following protocol is used to assess neuronal activity in PC12
30 cell lines. PC12 cells (rat pheochromocytoma cells) are known to proliferate and/or differentiate by activation with a number of mitogens, such as TPA (tetradecanoyl

phorbol acetate), NGF (nerve growth factor), and EGF (epidermal growth factor). The EGR1 gene expression is activated during this treatment. Thus, by stably transfecting PC12 cells with a construct containing an EGR promoter linked to SEAP reporter, activation of PC12 cells can be assessed.

5 The EGR/SEAP reporter construct can be assembled by the following protocol. The EGR-1 promoter sequence (-633 to +1)(Sakamoto K et al., Oncogene 6:867-871 (1991)) can be PCR amplified from human genomic DNA using the following primers:

5' GCGCTCGAGGGATGACAGCGATAGAACCCCGG -3' (SEQ ID NO:6)

10 5' GCGAAGCTTCGCGACTCCCCGGATCCGCCTC-3' (SEQ ID NO:7)

Using the GAS:SEAP/Neo vector produced in Example 12, EGR1 amplified product can then be inserted into this vector. Linearize the GAS:SEAP/Neo vector using restriction enzymes XhoI/HindIII, removing the GAS/SV40 stuffer. Restrict the EGR1 amplified product with these same enzymes. Ligate the vector and the EGR1
15 promoter.

To prepare 96 well-plates for cell culture, two mls of a coating solution (1:30 dilution of collagen type I (Upstate Biotech Inc. Cat#08-115) in 30% ethanol (filter sterilized)) is added per one 10 cm plate or 50 ml per well of the 96-well plate, and allowed to air dry for 2 hr.

20 PC12 cells are routinely grown in RPMI-1640 medium (Bio Whittaker) containing 10% horse serum (JRH BIOSCIENCES, Cat. # 12449-78P), 5% heat-inactivated fetal bovine serum (FBS) supplemented with 100 units/ml penicillin and 100 ug/ml streptomycin on a precoated 10 cm tissue culture dish. One to four split is done every three to four days. Cells are removed from the plates by scraping and
25 resuspended with pipetting up and down for more than 15 times.

Transfect the EGR/SEAP/Neo construct into PC12 using the Lipofectamine protocol described in Example 11. EGR-SEAP/PC12 stable cells are obtained by growing the cells in 300 ug/ml G418. The G418-free medium is used for routine growth but every one to two months, the cells should be re-grown in 300 ug/ml G418
30 for couple of passages.

To assay for neuronal activity, a 10 cm plate with cells around 70 to 80% confluent is screened by removing the old medium. Wash the cells once with PBS (Phosphate buffered saline). Then starve the cells in low serum medium (RPMI-1640 containing 1% horse serum and 0.5% FBS with antibiotics) overnight.

5 The next morning, remove the medium and wash the cells with PBS. Scrape off the cells from the plate, suspend the cells well in 2 ml low serum medium. Count the cell number and add more low serum medium to reach final cell density as 5×10^5 cells/ml.

 Add 200 ul of the cell suspension to each well of 96-well plate (equivalent to
10 1×10^5 cells/well). Add 50 ul supernatant produced by Example 11, 37°C for 48 to 72 hr. As a positive control, a growth factor known to activate PC12 cells through EGR can be used, such as 50 ng/ul of Neuronal Growth Factor (NGF). Over fifty-fold induction of SEAP is typically seen in the positive control wells. SEAP assay the supernatant according to Example 17.

15

Example 16: High-Throughput Screening Assay for T-cell Activity

 NF-KB (Nuclear Factor KB) is a transcription factor activated by a wide variety of agents including the inflammatory cytokines IL-1 and TNF, CD30 and CD40, lymphotoxin-alpha and lymphotoxin-beta, by exposure to LPS or thrombin,
20 and by expression of certain viral gene products. As a transcription factor, NF-KB regulates the expression of genes involved in immune cell activation, control of apoptosis (NF- KB appears to shield cells from apoptosis), B and T-cell development, anti-viral and antimicrobial responses, and multiple stress responses.

 In non-stimulated conditions, NF- KB is retained in the cytoplasm with I-KB
25 (Inhibitor KB). However, upon stimulation, I- KB is phosphorylated and degraded, causing NF- KB to shuttle to the nucleus, thereby activating transcription of target genes. Target genes activated by NF- KB include IL-2, IL-6, GM-CSF, ICAM-1 and class 1 MHC.

 Due to its central role and ability to respond to a range of stimuli, reporter
30 constructs utilizing the NF-KB promoter element are used to screen the supernatants produced in Example 11. Activators or inhibitors of NF-KB would be useful in

treating diseases. For example, inhibitors of NF-KB could be used to treat those diseases related to the acute or chronic activation of NF-KB, such as rheumatoid arthritis.

To construct a vector containing the NF-KB promoter element, a PCR based strategy is employed. The upstream primer contains four tandem copies of the NF-KB binding site (GGGGACTTTCCC) (SEQ ID NO:8), 18 bp of sequence complementary to the 5' end of the SV40 early promoter sequence, and is flanked with an XhoI site:

5':GCGGCCTCGAGGGGACTTTCCCGGGACTTTCCGGGGACTTTCCGGGAC
TTTCCATCCTGCCATCTCAATTAG:3' (SEQ ID NO:9)

The downstream primer is complementary to the 3' end of the SV40 promoter and is flanked with a Hind III site:

5':GCGGCAAGCTTTTTGCAAAGCCTAGGC:3' (SEQ ID NO:4)

PCR amplification is performed using the SV40 promoter template present in the pB-gal:promoter plasmid obtained from Clontech. The resulting PCR fragment is digested with XhoI and Hind III and subcloned into BLSK2-. (Stratagene) Sequencing with the T7 and T3 primers confirms the insert contains the following sequence:

5' : CTCGAGGGGACTTTCCCGGGGACTTTCCCGGGGACTTTCCGGGACTTTCCATCTGCCATCTCAATTA
GTCAGCAACCATAGTCCCGCCCCCTAACTCCGCCCATCCCGCCCCCTAACTCCGCCCAGTTCCGCCCATTC
TCCGCCCCATGGCTGACTAATTTTTTTTATTTATGCAGAGGCCGAGGCCGCTCGGCCTCTGAGCTATT
CCAGAAGTAGTGAGGAGGCTTTTTTGGAGGCCTAGGCTTTTGCAAAAAGCTT:3' (SEQ ID
NO:10)

Next, replace the SV40 minimal promoter element present in the pSEAP2-promoter plasmid (Clontech) with this NF-KB/SV40 fragment using XhoI and HindIII. However, this vector does not contain a neomycin resistance gene, and therefore, is not preferred for mammalian expression systems.

In order to generate stable mammalian cell lines, the NF-KB/SV40/SEAP cassette is removed from the above NF-KB/SEAP vector using restriction enzymes SalI and NotI, and inserted into a vector containing neomycin resistance. Particularly,

the NF-KB/SV40/SEAP cassette was inserted into pGFP-1 (Clontech), replacing the GFP gene, after restricting pGFP-1 with Sall and NotI.

Once NF-KB/SV40/SEAP/Neo vector is created, stable Jurkat T-cells are created and maintained according to the protocol described in Example 13. Similarly, the method for assaying supernatants with these stable Jurkat T-cells is also described in Example 13. As a positive control, exogenous TNF alpha (0.1, 1, 10 ng) is added to wells H9, H10, and H11, with a 5-10 fold activation typically observed.

Example 17: Assay for SEAP Activity

As a reporter molecule for the assays described in Examples 13-16, SEAP activity is assayed using the Tropix Phospho-light Kit (Cat. BP-400) according to the following general procedure. The Tropix Phospho-light Kit supplies the Dilution, Assay, and Reaction Buffers used below.

Prime a dispenser with the 2.5x Dilution Buffer and dispense 15 ul of 2.5x dilution buffer into Optiplates containing 35 ul of a supernatant. Seal the plates with a plastic sealer and incubate at 65 degree C for 30 min. Separate the Optiplates to avoid uneven heating.

Cool the samples to room temperature for 15 minutes. Empty the dispenser and prime with the Assay Buffer. Add 50 ml Assay Buffer and incubate at room temperature 5 min. Empty the dispenser and prime with the Reaction Buffer (see the table below). Add 50 ul Reaction Buffer and incubate at room temperature for 20 minutes. Since the intensity of the chemiluminescent signal is time dependent, and it takes about 10 minutes to read 5 plates on luminometer, one should treat 5 plates at each time and start the second set 10 minutes later.

Read the relative light unit in the luminometer. Set H12 as blank, and print the results. An increase in chemiluminescence indicates reporter activity.

Reaction Buffer Formulation:

# of plates	Rxn buffer diluent (ml)	CSPD (ml)
10	60	3
11	65	3.25
12	70	3.5
13	75	3.75

14	80	4
15	85	4.25
16	90	4.5
17	95	4.75
18	100	5
19	105	5.25
20	110	5.5
21	115	5.75
22	120	6
23	125	6.25
24	130	6.5
25	135	6.75
26	140	7
27	145	7.25
28	150	7.5
29	155	7.75
30	160	8
31	165	8.25
32	170	8.5
33	175	8.75
34	180	9
35	185	9.25
36	190	9.5
37	195	9.75
38	200	10
39	205	10.25
40	210	10.5
41	215	10.75
42	220	11
43	225	11.25
44	230	11.5
45	235	11.75
46	240	12
47	245	12.25
48	250	12.5
49	255	12.75
50	260	13

Example 18: High-Throughput Screening Assay Identifying Changes in Small Molecule Concentration and Membrane Permeability

Binding of a ligand to a receptor is known to alter intracellular levels of small molecules, such as calcium, potassium, sodium, and pH, as well as alter membrane potential. These alterations can be measured in an assay to identify supernatants which bind to receptors of a particular cell. Although the following protocol describes an assay for calcium, this protocol can easily be modified to detect changes in potassium, sodium, pH, membrane potential, or any other small molecule which is detectable by a fluorescent probe.

The following assay uses Fluorometric Imaging Plate Reader ("FLIPR") to measure changes in fluorescent molecules (Molecular Probes) that bind small molecules. Clearly, any fluorescent molecule detecting a small molecule can be used instead of the calcium fluorescent molecule, fluo-4 (Molecular Probes, Inc.; catalog no. F-14202), used here.

For adherent cells, seed the cells at 10,000 -20,000 cells/well in a Co-star black 96-well plate with clear bottom. The plate is incubated in a CO₂ incubator for 20 hours. The adherent cells are washed two times in Biotek washer with 200 ul of HBSS (Hank's Balanced Salt Solution) leaving 100 ul of buffer after the final wash.

A stock solution of 1 mg/ml fluo-4 is made in 10% pluronic acid DMSO. To load the cells with fluo-4, 50 ul of 12 ug/ml fluo-4 is added to each well. The plate is incubated at 37 degrees C in a CO₂ incubator for 60 min. The plate is washed four times in the Biotek washer with HBSS leaving 100 ul of buffer.

For non-adherent cells, the cells are spun down from culture media. Cells are re-suspended to 2-5x10⁶ cells/ml with HBSS in a 50-ml conical tube. 4 ul of 1 mg/ml fluo-4 solution in 10% pluronic acid DMSO is added to each ml of cell suspension. The tube is then placed in a 37 degrees C water bath for 30-60 min. The cells are washed twice with HBSS, resuspended to 1x10⁶ cells/ml, and dispensed into a microplate, 100 ul/well. The plate is centrifuged at 1000 rpm for 5 min. The plate is then washed once in Denley CellWash with 200 ul, followed by an aspiration step to 100 ul final volume.

For a non-cell based assay, each well contains a fluorescent molecule, such as fluo-4. The supernatant is added to the well, and a change in fluorescence is detected.

To measure the fluorescence of intracellular calcium, the FLIPR is set for the following parameters: (1) System gain is 300-800 mW; (2) Exposure time is 0.4 second; (3) Camera F/stop is F/2; (4) Excitation is 488 nm; (5) Emission is 530 nm; and (6) Sample addition is 50 ul. Increased emission at 530 nm indicates an extracellular signaling event which has resulted in an increase in the intracellular Ca⁺⁺ concentration.

Example 19: High-Throughput Screening Assay Identifying Tyrosine Kinase Activity

The Protein Tyrosine Kinases (PTK) represent a diverse group of transmembrane and cytoplasmic kinases. Within the Receptor Protein Tyrosine Kinase (RPTK) group are receptors for a range of mitogenic and metabolic growth factors including the PDGF, FGF, EGF, NGF, HGF and Insulin receptor subfamilies. In addition there are a large family of RPTKs for which the corresponding ligand is unknown. Ligands for RPTKs include mainly secreted small proteins, but also membrane-bound and extracellular matrix proteins.

Activation of RPTK by ligands involves ligand-mediated receptor dimerization, resulting in transphosphorylation of the receptor subunits and activation of the cytoplasmic tyrosine kinases. The cytoplasmic tyrosine kinases include receptor associated tyrosine kinases of the src-family (e.g., src, yes, lck, lyn, fyn) and non-receptor linked and cytosolic protein tyrosine kinases, such as the Jak family, members of which mediate signal transduction triggered by the cytokine superfamily of receptors (e.g., the Interleukins, Interferons, GM-CSF, and Leptin).

Because of the wide range of known factors capable of stimulating tyrosine kinase activity, the identification of novel human secreted proteins capable of activating tyrosine kinase signal transduction pathways are of interest. Therefore, the following protocol is designed to identify those novel human secreted proteins capable of activating the tyrosine kinase signal transduction pathways.

Seed target cells (e.g., primary keratinocytes) at a density of approximately 25,000 cells per well in a 96 well Loprodyne Silent Screen Plates purchased from Nalge Nunc (Naperville, IL). The plates are sterilized with two 30 minute rinses with 100% ethanol, rinsed with water and dried overnight. Some plates are coated for 2 hr with 100 ml of cell culture grade type I collagen (50 mg/ml), gelatin (2%) or polylysine (50 mg/ml), all of which can be purchased from Sigma Chemicals (St. Louis, MO) or 10% Matrigel purchased from Becton Dickinson (Bedford, MA), or calf serum, rinsed with PBS and stored at 4 degree C. Cell growth on these plates is assayed by seeding 5,000 cells/well in growth medium and indirect quantitation of cell number through use of alamarBlue as described by the manufacturer Alamar

Biosciences, Inc. (Sacramento, CA) after 48 hr. Falcon plate covers #3071 from Becton Dickinson (Bedford, MA) are used to cover the Loprodyne Silent Screen Plates. Falcon Microtest III cell culture plates can also be used in some proliferation experiments.

5 To prepare extracts, A431 cells are seeded onto the nylon membranes of Loprodyne plates (20,000/200ml/well) and cultured overnight in complete medium. Cells are quiesced by incubation in serum-free basal medium for 24 hr. After 5-20 minutes treatment with EGF (60ng/ml) or 50 ul of the supernatant produced in Example 11, the medium was removed and 100 ml of extraction buffer ((20 mM
10 HEPES pH 7.5, 0.15 M NaCl, 1% Triton X-100, 0.1% SDS, 2 mM Na₃VO₄, 2 mM Na₄P₂O₇ and a cocktail of protease inhibitors (# 1836170) obtained from Boehringer Mannheim (Indianapolis, IN) is added to each well and the plate is shaken on a rotating shaker for 5 minutes at 4 degrees C. The plate is then placed in a vacuum transfer manifold and the extract filtered through the 0.45 mm membrane
15 bottoms of each well using house vacuum. Extracts are collected in a 96-well catch/assay plate in the bottom of the vacuum manifold and immediately placed on ice. To obtain extracts clarified by centrifugation, the content of each well, after detergent solubilization for 5 minutes, is removed and centrifuged for 15 minutes at 4 degrees C at 16,000 x g.

20 Test the filtered extracts for levels of tyrosine kinase activity. Although many methods of detecting tyrosine kinase activity are known, one method is described here.

 Generally, the tyrosine kinase activity of a supernatant is evaluated by determining its ability to phosphorylate a tyrosine residue on a specific substrate (a
25 biotinylated peptide). Biotinylated peptides that can be used for this purpose include PSK1 (corresponding to amino acids 6-20 of the cell division kinase cdc2-p34) and PSK2 (corresponding to amino acids 1-17 of gastrin). Both peptides are substrates for a range of tyrosine kinases and are available from Boehringer Mannheim.

 The tyrosine kinase reaction is set up by adding the following components in
30 order. First, add 10ul of 5uM Biotinylated Peptide, then 10ul ATP/Mg₂⁺ (5mM ATP/50mM MgCl₂), then 10ul of 5x Assay Buffer (40mM imidazole hydrochloride,

pH7.3, 40 mM beta-glycerophosphate, 1mM EGTA, 100mM MgCl₂, 5 mM MnCl₂, 0.5 mg/ml BSA), then 5ul of Sodium Vanadate(1mM), and then 5ul of water. Mix the components gently and preincubate the reaction mix at 30 degrees C for 2 min. Initial the reaction by adding 10ul of the control enzyme or the filtered supernatant.

- 5 The tyrosine kinase assay reaction is then terminated by adding 10 ul of 120mM EDTA and place the reactions on ice.

Tyrosine kinase activity is determined by transferring 50 ul aliquot of reaction mixture to a microtiter plate (MTP) module and incubating at 37 degrees C for 20 min. This allows the streptavidin coated 96 well plate to associate with the
10 biotinylated peptide. Wash the MTP module with 300ul/well of PBS four times. Next add 75 ul of anti-phosphotyrosine antibody conjugated to horse radish peroxidase(anti-P-Tyr-POD(0.5u/ml)) to each well and incubate at 37 degrees C for one hour. Wash the well as above.

- 15 Next add 100ul of peroxidase substrate solution (Boehringer Mannheim) and incubate at room temperature for at least 5 mins (up to 30 min). Measure the absorbance of the sample at 405 nm by using ELISA reader. The level of bound peroxidase activity is quantitated using an ELISA reader and reflects the level of tyrosine kinase activity.

20 **Example 20: High-Throughput Screening Assay Identifying Phosphorylation Activity**

- As a potential alternative and/or compliment to the assay of protein tyrosine kinase activity described in Example 19, an assay which detects activation (phosphorylation) of major intracellular signal transduction intermediates can also be
25 used. For example, as described below one particular assay can detect tyrosine phosphorylation of the Erk-1 and Erk-2 kinases. However, phosphorylation of other molecules, such as Raf, JNK, p38 MAP, Map kinase kinase (MEK), MEK kinase, Src, Muscle specific kinase (MuSK), IRAK, Tec, and Janus, as well as any other phosphoserine, phosphotyrosine, or phosphothreonine molecule, can be detected by
30 substituting these molecules for Erk-1 or Erk-2 in the following assay.

Specifically, assay plates are made by coating the wells of a 96-well ELISA plate with 0.1ml of protein G (1ug/ml) for 2 hr at room temp, (RT). The plates are then rinsed with PBS and blocked with 3% BSA/PBS for 1 hr at RT. The protein G plates are then treated with 2 commercial monoclonal antibodies (100ng/well) against
5 Erk-1 and Erk-2 (1 hr at RT) (Santa Cruz Biotechnology). (To detect other molecules, this step can easily be modified by substituting a monoclonal antibody detecting any of the above described molecules.) After 3-5 rinses with PBS, the plates are stored at 4 degrees C until use.

A431 cells are seeded at 20,000/well in a 96-well Loprodyne filterplate and
10 cultured overnight in growth medium. The cells are then starved for 48 hr in basal medium (DMEM) and then treated with EGF (6ng/well) or 50 ul of the supernatants obtained in Example 11 for 5-20 minutes. The cells are then solubilized and extracts filtered directly into the assay plate.

After incubation with the extract for 1 hr at RT, the wells are again rinsed. As
15 a positive control, a commercial preparation of MAP kinase (10ng/well) is used in place of A431 extract. Plates are then treated with a commercial polyclonal (rabbit) antibody (1ug/ml) which specifically recognizes the phosphorylated epitope of the Erk-1 and Erk-2 kinases (1 hr at RT). This antibody is biotinylated by standard procedures. The bound polyclonal antibody is then quantitated by successive
20 incubations with Europium-streptavidin and Europium fluorescence enhancing reagent in the Wallac DELFIA instrument (time-resolved fluorescence). An increased fluorescent signal over background indicates a phosphorylation.

Example 21: Method of Determining Alterations in a Gene Corresponding to a
25 **Polynucleotide**

RNA isolated from entire families or individual patients presenting with a phenotype of interest (such as a disease) is be isolated. cDNA is then generated from these RNA samples using protocols known in the art. (See, Sambrook.) The cDNA is then used as a template for PCR, employing primers surrounding regions of interest
30 in SEQ ID NO:X. Suggested PCR conditions consist of 35 cycles at 95 degrees C for

30 seconds; 60-120 seconds at 52-58 degrees C; and 60-120 seconds at 70 degrees C, using buffer solutions described in Sidransky et al., Science 252:706 (1991).

PCR products are then sequenced using primers labeled at their 5' end with T4 polynucleotide kinase, employing SequiTherm Polymerase. (Epicentre
5 Technologies). The intron-exon borders of selected exons is also determined and genomic PCR products analyzed to confirm the results. PCR products harboring suspected mutations is then cloned and sequenced to validate the results of the direct sequencing.

PCR products is cloned into T-tailed vectors as described in Holton et al.,
10 Nucleic Acids Research, 19:1156 (1991) and sequenced with T7 polymerase (United States Biochemical). Affected individuals are identified by mutations not present in unaffected individuals.

Genomic rearrangements are also observed as a method of determining alterations in a gene corresponding to a polynucleotide. Genomic clones isolated
15 according to Example 2 are nick-translated with digoxigenindeoxy-uridine 5'-triphosphate (Boehringer Mannheim), and FISH performed as described in Johnson et al., Methods Cell Biol. 35:73-99 (1991). Hybridization with the labeled probe is carried out using a vast excess of human cot-1 DNA for specific hybridization to the corresponding genomic locus.

20 Chromosomes are counterstained with 4,6-diamino-2-phenylidole and propidium iodide, producing a combination of C- and R-bands. Aligned images for precise mapping are obtained using a triple-band filter set (Chroma Technology, Brattleboro, VT) in combination with a cooled charge-coupled device camera (Photometrics, Tucson, AZ) and variable excitation wavelength filters. (Johnson et
25 al., Genet. Anal. Tech. Appl., 8:75 (1991).) Image collection, analysis and chromosomal fractional length measurements are performed using the ISee Graphical Program System. (Inovision Corporation, Durham, NC.) Chromosome alterations of the genomic region hybridized by the probe are identified as insertions, deletions, and translocations. These alterations are used as a diagnostic marker for an associated
30 disease.

Example 22: Method of Detecting Abnormal Levels of a Polypeptide in a Biological Sample

5 A polypeptide of the present invention can be detected in a biological sample, and if an increased or decreased level of the polypeptide is detected, this polypeptide is a marker for a particular phenotype. Methods of detection are numerous, and thus, it is understood that one skilled in the art can modify the following assay to fit their particular needs.

10 For example, antibody-sandwich ELISAs are used to detect polypeptides in a sample, preferably a biological sample. Wells of a microtiter plate are coated with specific antibodies, at a final concentration of 0.2 to 10 ug/ml. The antibodies are either monoclonal or polyclonal and are produced by the method described in Example 10. The wells are blocked so that non-specific binding of the polypeptide to the well is reduced.

15 The coated wells are then incubated for > 2 hours at RT with a sample containing the polypeptide. Preferably, serial dilutions of the sample should be used to validate results. The plates are then washed three times with deionized or distilled water to remove unbounded polypeptide.

20 Next, 50 ul of specific antibody-alkaline phosphatase conjugate, at a concentration of 25-400 ng, is added and incubated for 2 hours at room temperature. The plates are again washed three times with deionized or distilled water to remove unbounded conjugate.

25 Add 75 ul of 4-methylumbelliferyl phosphate (MUP) or p-nitrophenyl phosphate (NPP) substrate solution to each well and incubate 1 hour at room temperature. Measure the reaction by a microtiter plate reader. Prepare a standard curve, using serial dilutions of a control sample, and plot polypeptide concentration on the X-axis (log scale) and fluorescence or absorbance of the Y-axis (linear scale). Interpolate the concentration of the polypeptide in the sample using the standard curve.

Example 23: Formulation

The invention also provides methods of treatment and/or prevention of diseases or disorders (such as, for example, any one or more of the diseases or disorders disclosed herein) by administration to a subject of an effective amount of a
5 Therapeutic. By therapeutic is meant a polynucleotides or polypeptides of the invention (including fragments and variants), agonists or antagonists thereof, and/or antibodies thereto, in combination with a pharmaceutically acceptable carrier type (e.g., a sterile carrier).

The Therapeutic will be formulated and dosed in a fashion consistent with
10 good medical practice, taking into account the clinical condition of the individual patient (especially the side effects of treatment with the Therapeutic alone), the site of delivery, the method of administration, the scheduling of administration, and other factors known to practitioners. The "effective amount" for purposes herein is thus determined by such considerations.

As a general proposition, the total pharmaceutically effective amount of the
15 Therapeutic administered parenterally per dose will be in the range of about 1 ug/kg/day to 10 mg/kg/day of patient body weight, although, as noted above, this will be subject to therapeutic discretion. More preferably, this dose is at least 0.01 mg/kg/day, and most preferably for humans between about 0.01 and 1 mg/kg/day for
20 the hormone. If given continuously, the Therapeutic is typically administered at a dose rate of about 1 ug/kg/hour to about 50 ug/kg/hour, either by 1-4 injections per day or by continuous subcutaneous infusions, for example, using a mini-pump. An intravenous bag solution may also be employed. The length of treatment needed to observe changes and the interval following treatment for responses to occur appears
25 to vary depending on the desired effect.

Therapeutics can be are administered orally, rectally, parenterally, intracistemally, intravaginally, intraperitoneally, topically (as by powders, ointments, gels, drops or transdermal patch), buccally, or as an oral or nasal spray.

"Pharmaceutically acceptable carrier" refers to a non-toxic solid, semisolid or liquid
30 filler, diluent, encapsulating material or formulation auxiliary of any. The term "parenteral" as used herein refers to modes of administration which include

intravenous, intramuscular, intraperitoneal, intrasternal, subcutaneous and intraarticular injection and infusion.

Therapeutics of the invention are also suitably administered by sustained-release systems. Suitable examples of sustained-release Therapeutics are administered orally, rectally, parenterally, intracisternally, intravaginally, intraperitoneally, topically (as by powders, ointments, gels, drops or transdermal patch), buccally, or as an oral or nasal spray. "Pharmaceutically acceptable carrier" refers to a non-toxic solid, semisolid or liquid filler, diluent, encapsulating material or formulation auxiliary of any type. The term "parenteral" as used herein refers to modes of administration which include intravenous, intramuscular, intraperitoneal, intrasternal, subcutaneous and intraarticular injection and infusion.

Therapeutics of the invention are also suitably administered by sustained-release systems. Suitable examples of sustained-release Therapeutics include suitable polymeric materials (such as, for example, semi-permeable polymer matrices in the form of shaped articles, e.g., films, or microcapsules), suitable hydrophobic materials (for example as an emulsion in an acceptable oil) or ion exchange resins, and sparingly soluble derivatives (such as, for example, a sparingly soluble salt).

Sustained-release matrices include polylactides (U.S. Pat. No. 3,773,919, EP 58,481), copolymers of L-glutamic acid and gamma-ethyl-L-glutamate (Sidman et al., Biopolymers 22:547-556 (1983)), poly (2-hydroxyethyl methacrylate) (Langer et al., J. Biomed. Mater. Res. 15:167-277 (1981), and Langer, Chem. Tech. 12:98-105 (1982)), ethylene vinyl acetate (Langer et al., Id.) or poly-D-(-)-3-hydroxybutyric acid (EP 133,988).

Sustained-release Therapeutics also include liposomally entrapped Therapeutics of the invention (see generally, Langer, *Science* 249:1527-1533 (1990); Treat et al., in *Liposomes in the Therapy of Infectious Disease and Cancer*, Lopez-Berestein and Fidler (eds.), Liss, New York, pp. 317-327 and 353-365 (1989)). Liposomes containing the Therapeutic are prepared by methods known per se: DE 3,218,121; Epstein et al., Proc. Natl. Acad. Sci. (USA) 82:3688-3692 (1985); Hwang et al., Proc. Natl. Acad. Sci. (USA) 77:4030-4034 (1980); EP 52,322; EP 36,676; EP 88,046; EP 143,949; EP 142,641; Japanese Pat. Appl. 83-118008; U.S. Pat. Nos.

4,485,045 and 4,544,545; and EP 102,324. Ordinarily, the liposomes are of the small (about 200-800 Angstroms) unilamellar type in which the lipid content is greater than about 30 mol. percent cholesterol, the selected proportion being adjusted for the optimal Therapeutic.

5 In yet an additional embodiment, the Therapeutics of the invention are delivered by way of a pump (*see* Langer, *supra*; Sefton, CRC Crit. Ref. Biomed. Eng. 14:201 (1987); Buchwald et al., Surgery 88:507 (1980); Saudek et al., N. Engl. J. Med. 321:574 (1989)).

Other controlled release systems are discussed in the review by Langer
10 (*Science* 249:1527-1533 (1990)).

For parenteral administration, in one embodiment, the Therapeutic is formulated generally by mixing it at the desired degree of purity, in a unit dosage injectable form (solution, suspension, or emulsion), with a pharmaceutically acceptable carrier, i.e., one that is non-toxic to recipients at the dosages and
15 concentrations employed and is compatible with other ingredients of the formulation. For example, the formulation preferably does not include oxidizing agents and other compounds that are known to be deleterious to the Therapeutic.

Generally, the formulations are prepared by contacting the Therapeutic uniformly and intimately with liquid carriers or finely divided solid carriers or both.
20 Then, if necessary, the product is shaped into the desired formulation. Preferably the carrier is a parenteral carrier, more preferably a solution that is isotonic with the blood of the recipient. Examples of such carrier vehicles include water, saline, Ringer's solution, and dextrose solution. Non-aqueous vehicles such as fixed oils and ethyl oleate are also useful herein, as well as liposomes.

25 The carrier suitably contains minor amounts of additives such as substances that enhance isotonicity and chemical stability. Such materials are non-toxic to recipients at the dosages and concentrations employed, and include buffers such as phosphate, citrate, succinate, acetic acid, and other organic acids or their salts; antioxidants such as ascorbic acid; low molecular weight (less than about ten
30 residues) polypeptides, e.g., polyarginine or tripeptides; proteins, such as serum albumin, gelatin, or immunoglobulins; hydrophilic polymers such as

polyvinylpyrrolidone; amino acids, such as glycine, glutamic acid, aspartic acid, or arginine; monosaccharides, disaccharides, and other carbohydrates including cellulose or its derivatives, glucose, manose, or dextrans; chelating agents such as EDTA; sugar alcohols such as mannitol or sorbitol; counterions such as sodium; and/or nonionic
5 surfactants such as polysorbates, poloxamers, or PEG.

The Therapeutic is typically formulated in such vehicles at a concentration of about 0.1 mg/ml to 100 mg/ml, preferably 1-10 mg/ml, at a pH of about 3 to 8. It will be understood that the use of certain of the foregoing excipients, carriers, or stabilizers will result in the formation of polypeptide salts.

10 Any pharmaceutical used for therapeutic administration can be sterile. Sterility is readily accomplished by filtration through sterile filtration membranes (e.g., 0.2 micron membranes). Therapeutics generally are placed into a container having a sterile access port, for example, an intravenous solution bag or vial having a stopper pierceable by a hypodermic injection needle.

15 Therapeutics ordinarily will be stored in unit or multi-dose containers, for example, sealed ampoules or vials, as an aqueous solution or as a lyophilized formulation for reconstitution. As an example of a lyophilized formulation, 10-ml vials are filled with 5 ml of sterile-filtered 1% (w/v) aqueous Therapeutic solution, and the resulting mixture is lyophilized. The infusion solution is prepared by
20 reconstituting the lyophilized Therapeutic using bacteriostatic Water-for-Injection.

The invention also provides a pharmaceutical pack or kit comprising one or more containers filled with one or more of the ingredients of the Therapeutics of the invention. Associated with such container(s) can be a notice in the form prescribed by a governmental agency regulating the manufacture, use or sale of pharmaceuticals or
25 biological products, which notice reflects approval by the agency of manufacture, use or sale for human administration. In addition, the Therapeutics may be employed in conjunction with other therapeutic compounds.

The Therapeutics of the invention may be administered alone or in combination with adjuvants. Adjuvants that may be administered with the
30 Therapeutics of the invention include, but are not limited to, alum, alum plus deoxycholate (ImmunoAg), MTP-PE (Biocine Corp.), QS21 (Genentech, Inc.), BCG,

and MPL. In a specific embodiment, Therapeutics of the invention are administered in combination with alum. In another specific embodiment, Therapeutics of the invention are administered in combination with QS-21. Further adjuvants that may be administered with the Therapeutics of the invention include, but are not limited to, 5 Monophosphoryl lipid immunomodulator, AdjuVax 100a, QS-21, QS-18, CRL1005, Aluminum salts, MF-59, and Virosomal adjuvant technology. Vaccines that may be administered with the Therapeutics of the invention include, but are not limited to, vaccines directed toward protection against MMR (measles, mumps, rubella), polio, varicella, tetanus/diphtheria, hepatitis A, hepatitis B, haemophilus influenzae B, 10 whooping cough, pneumonia, influenza, Lyme's Disease, rotavirus, cholera, yellow fever, Japanese encephalitis, poliomyelitis, rabies, typhoid fever, and pertussis. Combinations may be administered either concomitantly, e.g., as an admixture, separately but simultaneously or concurrently; or sequentially. This includes presentations in which the combined agents are administered together as a therapeutic 15 mixture, and also procedures in which the combined agents are administered separately but simultaneously, e.g., as through separate intravenous lines into the same individual. Administration "in combination" further includes the separate administration of one of the compounds or agents given first, followed by the second.

The Therapeutics of the invention may be administered alone or in 20 combination with other therapeutic agents. Therapeutic agents that may be administered in combination with the Therapeutics of the invention, include but not limited to, other members of the TNF family, chemotherapeutic agents, antibiotics, steroidal and non-steroidal anti-inflammatories, conventional immunotherapeutic agents, cytokines and/or growth factors. Combinations may be administered either 25 concomitantly, e.g., as an admixture, separately but simultaneously or concurrently; or sequentially. This includes presentations in which the combined agents are administered together as a therapeutic mixture, and also procedures in which the combined agents are administered separately but simultaneously, e.g., as through separate intravenous lines into the same individual. Administration "in combination" 30 further includes the separate administration of one of the compounds or agents given first, followed by the second.

In one embodiment, the Therapeutics of the invention are administered in combination with members of the TNF family. TNF, TNF-related or TNF-like molecules that may be administered with the Therapeutics of the invention include, but are not limited to, soluble forms of TNF-alpha, lymphotoxin-alpha (LT-alpha, also known as TNF-beta), LT-beta (found in complex heterotrimer LT-alpha2-beta), OPGL, FasL, CD27L, CD30L, CD40L, 4-1BBL, DcR3, OX40L, TNF-gamma (International Publication No. WO 96/14328), AIM-I (International Publication No. WO 97/33899), endokine-alpha (International Publication No. WO 98/07880), TR6 (International Publication No. WO 98/30694), OPG, and neutrokin-alpha (International Publication No. WO 98/18921, OX40, and nerve growth factor (NGF), and soluble forms of Fas, CD30, CD27, CD40 and 4-1BB, TR2 (International Publication No. WO 96/34095), DR3 (International Publication No. WO 97/33904), DR4 (International Publication No. WO 98/32856), TR5 (International Publication No. WO 98/30693), TR6 (International Publication No. WO 98/30694), TR7 (International Publication No. WO 98/41629), TRANK, TR9 (International Publication No. WO 98/56892), TR10 (International Publication No. WO 98/54202), 312C2 (International Publication No. WO 98/06842), and TR12, and soluble forms CD154, CD70, and CD153.

In certain embodiments, Therapeutics of the invention are administered in combination with antiretroviral agents, nucleoside reverse transcriptase inhibitors, non-nucleoside reverse transcriptase inhibitors, and/or protease inhibitors. Nucleoside reverse transcriptase inhibitors that may be administered in combination with the Therapeutics of the invention, include, but are not limited to, RETROVIR™ (zidovudine/AZT), VIDEX™ (didanosine/ddI), HIVID™ (zalcitabine/ddC), ZERIT™ (stavudine/d4T), EPIVIR™ (lamivudine/3TC), and COMBIVIR™ (zidovudine/lamivudine). Non-nucleoside reverse transcriptase inhibitors that may be administered in combination with the Therapeutics of the invention, include, but are not limited to, VIRAMUNE™ (nevirapine), RESCRIPTOR™ (delavirdine), and SUSTIVA™ (efavirenz). Protease inhibitors that may be administered in combination with the Therapeutics of the invention, include, but are not limited to, CRIVAN™ (indinavir), NORVIR™ (ritonavir), INVIRASE™ (saquinavir), and

VIRACEPT™ (nelfinavir). In a specific embodiment, antiretroviral agents, nucleoside reverse transcriptase inhibitors, non-nucleoside reverse transcriptase inhibitors, and/or protease inhibitors may be used in any combination with Therapeutics of the invention to treat AIDS and/or to prevent or treat HIV infection.

- 5 In other embodiments, Therapeutics of the invention may be administered in combination with anti-opportunistic infection agents. Anti-opportunistic agents that may be administered in combination with the Therapeutics of the invention, include, but are not limited to, TRIMETHOPRIM-SULFAMETHOXAZOLE™, DAPSONE™, PENTAMIDINE™, ATOVAQUONE™, ISONIAZID™,
- 10 RIFAMPIN™, PYRAZINAMIDE™, ETHAMBUTOL™, RIFABUTIN™, CLARITHROMYCIN™, AZITHROMYCIN™, GANCICLOVIR™, FOSCARNET™, CIDOFOVIR™, FLUCONAZOLE™, ITRACONAZOLE™, KETOCONAZOLE™, ACYCLOVIR™, FAMCICOLVIR™, PYRIMETHAMINE™, LEUCOVORIN™, NEUPOGEN™ (filgrastim/G-CSF), and LEUKINE™
- 15 (sargramostim/GM-CSF). In a specific embodiment, Therapeutics of the invention are used in any combination with TRIMETHOPRIM-SULFAMETHOXAZOLE™, DAPSONE™, PENTAMIDINE™, and/or ATOVAQUONE™ to prophylactically treat or prevent an opportunistic *Pneumocystis carinii* pneumonia infection. In another specific embodiment, Therapeutics of the invention are used in any
- 20 combination with ISONIAZID™, RIFAMPIN™, PYRAZINAMIDE™, and/or ETHAMBUTOL™ to prophylactically treat or prevent an opportunistic *Mycobacterium avium* complex infection. In another specific embodiment, Therapeutics of the invention are used in any combination with RIFABUTIN™, CLARITHROMYCIN™, and/or AZITHROMYCIN™ to prophylactically treat or
- 25 prevent an opportunistic *Mycobacterium tuberculosis* infection. In another specific embodiment, Therapeutics of the invention are used in any combination with GANCICLOVIR™, FOSCARNET™, and/or CIDOFOVIR™ to prophylactically treat or prevent an opportunistic cytomegalovirus infection. In another specific embodiment, Therapeutics of the invention are used in any combination with
- 30 FLUCONAZOLE™, ITRACONAZOLE™, and/or KETOCONAZOLE™ to

prophylactically treat or prevent an opportunistic fungal infection. In another specific embodiment, Therapeutics of the invention are used in any combination with ACYCLOVIR™ and/or FAMCICOLVIR™ to prophylactically treat or prevent an opportunistic herpes simplex virus type I and/or type II infection. In another specific
5 embodiment, Therapeutics of the invention are used in any combination with PYRIMETHAMINE™ and/or LEUCOVORIN™ to prophylactically treat or prevent an opportunistic *Toxoplasma gondii* infection. In another specific embodiment, Therapeutics of the invention are used in any combination with LEUCOVORIN™ and/or NEUPOGEN™ to prophylactically treat or prevent an opportunistic bacterial
10 infection.

In a further embodiment, the Therapeutics of the invention are administered in combination with an antiviral agent. Antiviral agents that may be administered with the Therapeutics of the invention include, but are not limited to, acyclovir, ribavirin, amantadine, and remantidine.

15 In a further embodiment, the Therapeutics of the invention are administered in combination with an antibiotic agent. Antibiotic agents that may be administered with the Therapeutics of the invention include, but are not limited to, amoxicillin, beta-lactamases, aminoglycosides, beta-lactam (glycopeptide), beta-lactamases, Clindamycin, chloramphenicol, cephalosporins, ciprofloxacin, ciprofloxacin,
20 erythromycin, fluoroquinolones, macrolides, metronidazole, penicillins, quinolones, rifampin, streptomycin, sulfonamide, tetracyclines, trimethoprim, trimethoprim-sulfamthoxazole, and vancomycin.

Conventional nonspecific immunosuppressive agents, that may be administered in combination with the Therapeutics of the invention include, but are
25 not limited to, steroids, cyclosporine, cyclosporine analogs, cyclophosphamide methylprednisone, prednisone, azathioprine, FK-506, 15-deoxyspergualin, and other immunosuppressive agents that act by suppressing the function of responding T cells.

In specific embodiments, Therapeutics of the invention are administered in combination with immunosuppressants. Immunosuppressants preparations that may
30 be administered with the Therapeutics of the invention include, but are not limited to, ORTHOCLONE™ (OKT3), SANDIMMUNET™/NEORAL™/SANGDYA™

(cyclosporin), PROGRAF™ (tacrolimus), CELLCEPT™ (mycophenolate), Azathioprine, glucorticosteroids, and RAPAMUNE™ (sirolimus). In a specific embodiment, immunosuppressants may be used to prevent rejection of organ or bone marrow transplantation.

5 In an additional embodiment, Therapeutics of the invention are administered alone or in combination with one or more intravenous immune globulin preparations. Intravenous immune globulin preparations that may be administered with the Therapeutics of the invention include, but not limited to, GAMMAR™, IVEEGAM™, SANDOGLOBULIN™, GAMMAGARD S/D™, and GAMIMUNE™.

10 In a specific embodiment, Therapeutics of the invention are administered in combination with intravenous immune globulin preparations in transplantation therapy (e.g., bone marrow transplant).

In an additional embodiment, the Therapeutics of the invention are administered alone or in combination with an anti-inflammatory agent. Anti-
15 inflammatory agents that may be administered with the Therapeutics of the invention include, but are not limited to, glucocorticoids and the nonsteroidal anti-inflammatories, aminoarylcarboxylic acid derivatives, arylacetic acid derivatives, arylbutyric acid derivatives, arylcarboxylic acids, arylpropionic acid derivatives, pyrazoles, pyrazolones, salicylic acid derivatives, thiazinecarboxamides, e-
20 acetamidocaproic acid, S-adenosylmethionine, 3-amino-4-hydroxybutyric acid, amixetrine, bendazac, benzydamine, bucolome, difenpiramide, ditazol, emorfazone, guaiazulene, nabumetone, nimesulide, orgotein, oxaceprol, paranyline, perisoxal, pifoxime, proquazone, proxazole, and tenidap.

In another embodiment, compositions of the invention are administered in
25 combination with a chemotherapeutic agent. Chemotherapeutic agents that may be administered with the Therapeutics of the invention include, but are not limited to, antibiotic derivatives (e.g., doxorubicin, bleomycin, daunorubicin, and dactinomycin); antiestrogens (e.g., tamoxifen); antimetabolites (e.g., fluorouracil, 5-FU, methotrexate, floxuridine, interferon alpha-2b, glutamic acid, plicamycin,
30 mercaptopurine, and 6-thioguanine); cytotoxic agents (e.g., carmustine, BCNU, lomustine, CCNU, cytosine arabinoside, cyclophosphamide, estramustine,

hydroxyurea, procarbazine, mitomycin, busulfan, cis-platin, and vincristine sulfate); hormones (e.g., medroxyprogesterone, estramustine phosphate sodium, ethinyl estradiol, estradiol, megestrol acetate, methyltestosterone, diethylstilbestrol diphosphate, chlorotrianisene, and testolactone); nitrogen mustard derivatives (e.g.,
5 mephallen, chorambucil, mechlorethamine (nitrogen mustard) and thiotepa); steroids and combinations (e.g., bethamethasone sodium phosphate); and others (e.g., dicarbazine, asparaginase, mitotane, vincristine sulfate, vinblastine sulfate, and etoposide).

In a specific embodiment, Therapeutics of the invention are administered in
10 combination with CHOP (cyclophosphamide, doxorubicin, vincristine, and prednisone) or any combination of the components of CHOP. In another embodiment, Therapeutics of the invention are administered in combination with Rituximab. In a further embodiment, Therapeutics of the invention are administered with Rituxmab and CHOP, or Rituxmab and any combination of the components of
15 CHOP.

In an additional embodiment, the Therapeutics of the invention are administered in combination with cytokines. Cytokines that may be administered with the Therapeutics of the invention include, but are not limited to, IL2, IL3, IL4, IL5, IL6, IL7, IL10, IL12, IL13, IL15, anti-CD40, CD40L, IFN-gamma and TNF-
20 alpha. In another embodiment, Therapeutics of the invention may be administered with any interleukin, including, but not limited to, IL-1alpha, IL-1beta, IL-2, IL-3, IL-4, IL-5, IL-6, IL-7, IL-8, IL-9, IL-10, IL-11, IL-12, IL-13, IL-14, IL-15, IL-16, IL-17, IL-18, IL-19, IL-20, and IL-21.

In an additional embodiment, the Therapeutics of the invention are
25 administered in combination with angiogenic proteins. Angiogenic proteins that may be administered with the Therapeutics of the invention include, but are not limited to, Glioma Derived Growth Factor (GDGF), as disclosed in European Patent Number EP-399816; Platelet Derived Growth Factor-A (PDGF-A), as disclosed in European Patent Number EP-682110; Platelet Derived Growth Factor-B (PDGF-B), as
30 disclosed in European Patent Number EP-282317; Placental Growth Factor (PlGF), as disclosed in International Publication Number WO 92/06194; Placental Growth

Factor-2 (PlGF-2), as disclosed in Hauser et al., Growth Factors, 4:259-268 (1993); Vascular Endothelial Growth Factor (VEGF), as disclosed in International Publication Number WO 90/13649; Vascular Endothelial Growth Factor-A (VEGF-A), as disclosed in European Patent Number EP-506477; Vascular Endothelial Growth
5 Factor-2 (VEGF-2), as disclosed in International Publication Number WO 96/39515; Vascular Endothelial Growth Factor B (VEGF-3); Vascular Endothelial Growth Factor B-186 (VEGF-B186), as disclosed in International Publication Number WO 96/26736; Vascular Endothelial Growth Factor-D (VEGF-D), as disclosed in International Publication Number WO 98/02543; Vascular Endothelial Growth
10 Factor-D (VEGF-D), as disclosed in International Publication Number WO 98/07832; and Vascular Endothelial Growth Factor-E (VEGF-E), as disclosed in German Patent Number DE19639601. The above mentioned references are incorporated herein by reference herein.

In an additional embodiment, the Therapeutics of the invention are
15 administered in combination with hematopoietic growth factors. Hematopoietic growth factors that may be administered with the Therapeutics of the invention include, but are not limited to, LEUKINE™ (SARGRAMOSTIM™) and NEUPOGEN™ (FILGRASTIM™).

In an additional embodiment, the Therapeutics of the invention are
20 administered in combination with Fibroblast Growth Factors. Fibroblast Growth Factors that may be administered with the Therapeutics of the invention include, but are not limited to, FGF-1, FGF-2, FGF-3, FGF-4, FGF-5, FGF-6, FGF-7, FGF-8, FGF-9, FGF-10, FGF-11, FGF-12, FGF-13, FGF-14, and FGF-15.

In additional embodiments, the Therapeutics of the invention are administered in
25 combination with other therapeutic or prophylactic regimens, such as, for example, radiation therapy.

Example 24: Method of Treating Decreased Levels of the Polypeptide

The present invention relates to a method for treating an individual in need of
30 an increased level of a polypeptide of the invention in the body comprising administering to such an individual a composition comprising a therapeutically

effective amount of an agonist of the invention (including polypeptides of the invention). Moreover, it will be appreciated that conditions caused by a decrease in the standard or normal expression level of a secreted protein in an individual can be treated by administering the polypeptide of the present invention, preferably in the secreted form. Thus, the invention also provides a method of treatment of an individual in need of an increased level of the polypeptide comprising administering to such an individual a Therapeutic comprising an amount of the polypeptide to increase the activity level of the polypeptide in such an individual.

For example, a patient with decreased levels of a polypeptide receives a daily dose 0.1-100 ug/kg of the polypeptide for six consecutive days. Preferably, the polypeptide is in the secreted form. The exact details of the dosing scheme, based on administration and formulation, are provided in Example 23.

Example 25: Method of Treating Increased Levels of the Polypeptide

The present invention also relates to a method of treating an individual in need of a decreased level of a polypeptide of the invention in the body comprising administering to such an individual a composition comprising a therapeutically effective amount of an antagonist of the invention (including polypeptides and antibodies of the invention).

In one example, antisense technology is used to inhibit production of a polypeptide of the present invention. This technology is one example of a method of decreasing levels of a polypeptide, preferably a secreted form, due to a variety of etiologies, such as cancer. For example, a patient diagnosed with abnormally increased levels of a polypeptide is administered intravenously antisense polynucleotides at 0.5, 1.0, 1.5, 2.0 and 3.0 mg/kg day for 21 days. This treatment is repeated after a 7-day rest period if the treatment was well tolerated. The formulation of the antisense polynucleotide is provided in Example 23.

Example 26: Method of Treatment Using Gene Therapy-Ex Vivo

One method of gene therapy transplants fibroblasts, which are capable of expressing a polypeptide, onto a patient. Generally, fibroblasts are obtained from a

subject by skin biopsy. The resulting tissue is placed in tissue-culture medium and separated into small pieces. Small chunks of the tissue are placed on a wet surface of a tissue culture flask, approximately ten pieces are placed in each flask. The flask is turned upside down, closed tight and left at room temperature over night. After 24
5 hours at room temperature, the flask is inverted and the chunks of tissue remain fixed to the bottom of the flask and fresh media (e.g., Ham's F12 media, with 10% FBS, penicillin and streptomycin) is added. The flasks are then incubated at 37 degree C for approximately one week.

At this time, fresh media is added and subsequently changed every several
10 days. After an additional two weeks in culture, a monolayer of fibroblasts emerge. The monolayer is trypsinized and scaled into larger flasks.

pMV-7 (Kirschmeier, P.T. et al., DNA, 7:219-25 (1988)), flanked by the long terminal repeats of the Moloney murine sarcoma virus, is digested with EcoRI and HindIII and subsequently treated with calf intestinal phosphatase. The linear vector is
15 fractionated on agarose gel and purified, using glass beads.

The cDNA encoding a polypeptide of the present invention can be amplified using PCR primers which correspond to the 5' and 3' end sequences respectively as set forth in Example 1 using primers and having appropriate restriction sites and initiation/stop codons, if necessary. Preferably, the 5' primer contains an EcoRI site
20 and the 3' primer includes a HindIII site. Equal quantities of the Moloney murine sarcoma virus linear backbone and the amplified EcoRI and HindIII fragment are added together, in the presence of T4 DNA ligase. The resulting mixture is maintained under conditions appropriate for ligation of the two fragments. The ligation mixture is then used to transform bacteria HB101, which are then plated onto
25 agar containing kanamycin for the purpose of confirming that the vector has the gene of interest properly inserted.

The amphotropic pA317 or GP+am12 packaging cells are grown in tissue culture to confluent density in Dulbecco's Modified Eagles Medium (DMEM) with 10% calf serum (CS), penicillin and streptomycin. The MSV vector containing the
30 gene is then added to the media and the packaging cells transduced with the vector.

The packaging cells now produce infectious viral particles containing the gene (the packaging cells are now referred to as producer cells).

Fresh media is added to the transduced producer cells, and subsequently, the media is harvested from a 10 cm plate of confluent producer cells. The spent media,
5 containing the infectious viral particles, is filtered through a millipore filter to remove detached producer cells and this media is then used to infect fibroblast cells. Media is removed from a sub-confluent plate of fibroblasts and quickly replaced with the media from the producer cells. This media is removed and replaced with fresh media. If the titer of virus is high, then virtually all fibroblasts will be infected and no
10 selection is required. If the titer is very low, then it is necessary to use a retroviral vector that has a selectable marker, such as neo or his. Once the fibroblasts have been efficiently infected, the fibroblasts are analyzed to determine whether protein is produced.

The engineered fibroblasts are then transplanted onto the host, either alone or
15 after having been grown to confluence on cytodex 3 microcarrier beads.

Example 27: Gene Therapy Using Endogenous Genes Corresponding To Polynucleotides of the Invention

Another method of gene therapy according to the present invention involves
20 operably associating the endogenous polynucleotide sequence of the invention with a promoter via homologous recombination as described, for example, in U.S. Patent NO: 5,641,670, issued June 24, 1997; International Publication NO: WO 96/29411, published September 26, 1996; International Publication NO: WO 94/12650, published August 4, 1994; Koller et al., *Proc. Natl. Acad. Sci. USA*, 86:8932-8935
25 (1989); and Zijlstra et al., *Nature*, 342:435-438 (1989). This method involves the activation of a gene which is present in the target cells, but which is not expressed in the cells, or is expressed at a lower level than desired.

Polynucleotide constructs are made which contain a promoter and targeting sequences, which are homologous to the 5' non-coding sequence of endogenous
30 polynucleotide sequence, flanking the promoter. The targeting sequence will be sufficiently near the 5' end of the polynucleotide sequence so the promoter will be

operably linked to the endogenous sequence upon homologous recombination. The promoter and the targeting sequences can be amplified using PCR. Preferably, the amplified promoter contains distinct restriction enzyme sites on the 5' and 3' ends. Preferably, the 3' end of the first targeting sequence contains the same restriction enzyme site as the 5' end of the amplified promoter and the 5' end of the second targeting sequence contains the same restriction site as the 3' end of the amplified promoter.

The amplified promoter and the amplified targeting sequences are digested with the appropriate restriction enzymes and subsequently treated with calf intestinal phosphatase. The digested promoter and digested targeting sequences are added together in the presence of T4 DNA ligase. The resulting mixture is maintained under conditions appropriate for ligation of the two fragments. The construct is size fractionated on an agarose gel then purified by phenol extraction and ethanol precipitation.

In this Example, the polynucleotide constructs are administered as naked polynucleotides via electroporation. However, the polynucleotide constructs may also be administered with transfection-facilitating agents, such as liposomes, viral sequences, viral particles, precipitating agents, etc. Such methods of delivery are known in the art.

Once the cells are transfected, homologous recombination will take place which results in the promoter being operably linked to the endogenous polynucleotide sequence. This results in the expression of polynucleotide corresponding to the polynucleotide in the cell. Expression may be detected by immunological staining, or any other method known in the art.

Fibroblasts are obtained from a subject by skin biopsy. The resulting tissue is placed in DMEM + 10% fetal calf serum. Exponentially growing or early stationary phase fibroblasts are trypsinized and rinsed from the plastic surface with nutrient medium. An aliquot of the cell suspension is removed for counting, and the remaining cells are subjected to centrifugation. The supernatant is aspirated and the pellet is resuspended in 5 ml of electroporation buffer (20 mM HEPES pH 7.3, 137 mM NaCl, 5 mM KCl, 0.7 mM Na_2HPO_4 , 6 mM dextrose). The cells are recentrifuged, the

supernatant aspirated, and the cells resuspended in electroporation buffer containing 1 mg/ml acetylated bovine serum albumin. The final cell suspension contains approximately 3×10^6 cells/ml. Electroporation should be performed immediately following resuspension.

5 Plasmid DNA is prepared according to standard techniques. For example, to construct a plasmid for targeting to the locus corresponding to the polynucleotide of the invention, plasmid pUC18 (MBI Fermentas, Amherst, NY) is digested with HindIII. The CMV promoter is amplified by PCR with an XbaI site on the 5' end and a BamHI site on the 3' end. Two non-coding sequences are amplified via PCR: one
10 non-coding sequence (fragment 1) is amplified with a HindIII site at the 5' end and an Xba site at the 3' end; the other non-coding sequence (fragment 2) is amplified with a BamHI site at the 5' end and a HindIII site at the 3' end. The CMV promoter and the fragments (1 and 2) are digested with the appropriate enzymes (CMV promoter - XbaI and BamHI; fragment 1 - XbaI; fragment 2 - BamHI) and ligated together. The
15 resulting ligation product is digested with HindIII, and ligated with the HindIII-digested pUC18 plasmid.

Plasmid DNA is added to a sterile cuvette with a 0.4 cm electrode gap (Bio-Rad). The final DNA concentration is generally at least 120 $\mu\text{g/ml}$. 0.5 ml of the cell suspension (containing approximately 1.5×10^6 cells) is then added to the cuvette,
20 and the cell suspension and DNA solutions are gently mixed. Electroporation is performed with a Gene-Pulser apparatus (Bio-Rad). Capacitance and voltage are set at 960 μF and 250-300 V, respectively. As voltage increases, cell survival decreases, but the percentage of surviving cells that stably incorporate the introduced DNA into their genome increases dramatically. Given these parameters, a pulse time of
25 approximately 14-20 mSec should be observed.

Electroporated cells are maintained at room temperature for approximately 5 min, and the contents of the cuvette are then gently removed with a sterile transfer pipette. The cells are added directly to 10 ml of prewarmed nutrient media (DMEM with 15% calf serum) in a 10 cm dish and incubated at 37 degree C. The following
30 day, the media is aspirated and replaced with 10 ml of fresh media and incubated for a further 16-24 hours.

The engineered fibroblasts are then injected into the host, either alone or after having been grown to confluence on cytodex 3 microcarrier beads. The fibroblasts now produce the protein product. The fibroblasts can then be introduced into a patient as described above.

5

Example 28: Method of Treatment Using Gene Therapy - In Vivo

Another aspect of the present invention is using *in vivo* gene therapy methods to treat disorders, diseases and conditions. The gene therapy method relates to the introduction of naked nucleic acid (DNA, RNA, and antisense DNA or RNA) sequences into an animal to increase or decrease the expression of the polypeptide. The polynucleotide of the present invention may be operatively linked to a promoter or any other genetic elements necessary for the expression of the polypeptide by the target tissue. Such gene therapy and delivery techniques and methods are known in the art, see, for example, WO90/11092, WO98/11779; U.S. Patent NO. 5693622, 10 5705151, 5580859; Tabata et al., Cardiovasc. Res. 35(3):470-479 (1997); Chao et al., Pharmacol. Res. 35(6):517-522 (1997); Wolff, Neuromuscul. Disord. 7(5):314-318 (1997); Schwartz et al., Gene Ther. 3(5):405-411 (1996); Tsurumi et al., Circulation 15 94(12):3281-3290 (1996) (incorporated herein by reference).

The polynucleotide constructs may be delivered by any method that delivers injectable materials to the cells of an animal, such as, injection into the interstitial 20 space of tissues (heart, muscle, skin, lung, liver, intestine and the like). The polynucleotide constructs can be delivered in a pharmaceutically acceptable liquid or aqueous carrier.

The term "naked" polynucleotide, DNA or RNA, refers to sequences that are free from any delivery vehicle that acts to assist, promote, or facilitate entry into the 25 cell, including viral sequences, viral particles, liposome formulations, lipofectin or precipitating agents and the like. However, the polynucleotides of the present invention may also be delivered in liposome formulations (such as those taught in Felgner P.L. et al. (1995) Ann. NY Acad. Sci. 772:126-139 and Abdallah B. et al. 30 (1995) Biol. Cell 85(1):1-7) which can be prepared by methods well known to those skilled in the art.

The polynucleotide vector constructs used in the gene therapy method are preferably constructs that will not integrate into the host genome nor will they contain sequences that allow for replication. Any strong promoter known to those skilled in the art can be used for driving the expression of DNA. Unlike other gene therapies
5 techniques, one major advantage of introducing naked nucleic acid sequences into target cells is the transitory nature of the polynucleotide synthesis in the cells. Studies have shown that non-replicating DNA sequences can be introduced into cells to provide production of the desired polypeptide for periods of up to six months.

The polynucleotide construct can be delivered to the interstitial space of
10 tissues within the an animal, including of muscle, skin, brain, lung, liver, spleen, bone marrow, thymus, heart, lymph, blood, bone, cartilage, pancreas, kidney, gall bladder, stomach, intestine, testis, ovary, uterus, rectum, nervous system, eye, gland, and connective tissue. Interstitial space of the tissues comprises the intercellular fluid, mucopolysaccharide matrix among the reticular fibers of organ tissues, elastic fibers
15 in the walls of vessels or chambers, collagen fibers of fibrous tissues, or that same matrix within connective tissue ensheathing muscle cells or in the lacunae of bone. It is similarly the space occupied by the plasma of the circulation and the lymph fluid of the lymphatic channels. Delivery to the interstitial space of muscle tissue is preferred for the reasons discussed below. They may be conveniently delivered by injection
20 into the tissues comprising these cells. They are preferably delivered to and expressed in persistent, non-dividing cells which are differentiated, although delivery and expression may be achieved in non-differentiated or less completely differentiated cells, such as, for example, stem cells of blood or skin fibroblasts. *In vivo* muscle cells are particularly competent in their ability to take up and express
25 polynucleotides.

For the naked polynucleotide injection, an effective dosage amount of DNA or RNA will be in the range of from about 0.05 g/kg body weight to about 50 mg/kg body weight. Preferably the dosage will be from about 0.005 mg/kg to about 20 mg/kg and more preferably from about 0.05 mg/kg to about 5 mg/kg. Of course, as
30 the artisan of ordinary skill will appreciate, this dosage will vary according to the tissue site of injection. The appropriate and effective dosage of nucleic acid sequence

can readily be determined by those of ordinary skill in the art and may depend on the condition being treated and the route of administration. The preferred route of administration is by the parenteral route of injection into the interstitial space of tissues. However, other parenteral routes may also be used, such as, inhalation of an aerosol formulation particularly for delivery to lungs or bronchial tissues, throat or
5 mucous membranes of the nose. In addition, naked polynucleotide constructs can be delivered to arteries during angioplasty by the catheter used in the procedure.

The dose response effects of injected polynucleotide in muscle *in vivo* is determined as follows. Suitable template DNA for production of mRNA coding for
10 polypeptide of the present invention is prepared in accordance with a standard recombinant DNA methodology. The template DNA, which may be either circular or linear, is either used as naked DNA or complexed with liposomes. The quadriceps muscles of mice are then injected with various amounts of the template DNA.

Five to six week old female and male Balb/C mice are anesthetized by
15 intraperitoneal injection with 0.3 ml of 2.5% Avertin. A 1.5 cm incision is made on the anterior thigh, and the quadriceps muscle is directly visualized. The template DNA is injected in 0.1 ml of carrier in a 1 cc syringe through a 27 gauge needle over one minute, approximately 0.5 cm from the distal insertion site of the muscle into the knee and about 0.2 cm deep. A suture is placed over the injection site for future
20 localization, and the skin is closed with stainless steel clips.

After an appropriate incubation time (e.g., 7 days) muscle extracts are prepared by excising the entire quadriceps. Every fifth 15 um cross-section of the individual quadriceps muscles is histochemically stained for protein expression. A time course for protein expression may be done in a similar fashion except that
25 quadriceps from different mice are harvested at different times. Persistence of DNA in muscle following injection may be determined by Southern blot analysis after preparing total cellular DNA and HIRT supernatants from injected and control mice. The results of the above experimentation in mice can be use to extrapolate proper dosages and other treatment parameters in humans and other animals using naked
30 DNA.

Example 29: Transgenic Animals.

The polypeptides of the invention can also be expressed in transgenic animals. Animals of any species, including, but not limited to, mice, rats, rabbits, hamsters, guinea pigs, pigs, micro-pigs, goats, sheep, cows and non-human primates, *e.g.*,
5 baboons, monkeys, and chimpanzees may be used to generate transgenic animals. In a specific embodiment, techniques described herein or otherwise known in the art, are used to express polypeptides of the invention in humans, as part of a gene therapy protocol.

Any technique known in the art may be used to introduce the transgene (*i.e.*,
10 polynucleotides of the invention) into animals to produce the founder lines of transgenic animals. Such techniques include, but are not limited to, pronuclear microinjection (Paterson et al., Appl. Microbiol. Biotechnol. 40:691-698 (1994); Carver et al., Biotechnology (NY) 11:1263-1270 (1993); Wright et al., Biotechnology (NY) 9:830-834 (1991); and Hoppe et al., U.S. Pat. No. 4,873,191 (1989)); retrovirus
15 mediated gene transfer into germ lines (Van der Putten et al., Proc. Natl. Acad. Sci., USA 82:6148-6152 (1985)), blastocysts or embryos; gene targeting in embryonic stem cells (Thompson et al., Cell 56:313-321 (1989)); electroporation of cells or embryos (Lo, 1983, Mol Cell. Biol. 3:1803-1814 (1983)); introduction of the polynucleotides of the invention using a gene gun (see, *e.g.*, Ulmer et al., Science
20 259:1745 (1993); introducing nucleic acid constructs into embryonic pluripotent stem cells and transferring the stem cells back into the blastocyst; and sperm-mediated gene transfer (Lavitrano et al., Cell 57:717-723 (1989); etc. For a review of such techniques, see Gordon, "Transgenic Animals," Intl. Rev. Cytol. 115:171-229 (1989), which is incorporated by reference herein in its entirety.

25 Any technique known in the art may be used to produce transgenic clones containing polynucleotides of the invention, for example, nuclear transfer into enucleated oocytes of nuclei from cultured embryonic, fetal, or adult cells induced to quiescence (Campell et al., Nature 380:64-66 (1996); Wilmut et al., Nature 385:810-813 (1997)).

30 The present invention provides for transgenic animals that carry the transgene in all their cells, as well as animals which carry the transgene in some, but not all their

cells, *i.e.*, mosaic animals or chimeric. The transgene may be integrated as a single transgene or as multiple copies such as in concatamers, *e.g.*, head-to-head tandems or head-to-tail tandems. The transgene may also be selectively introduced into and activated in a particular cell type by following, for example, the teaching of Lasko et al. (Lasko et al., Proc. Natl. Acad. Sci. USA 89:6232-6236 (1992)). The regulatory sequences required for such a cell-type specific activation will depend upon the particular cell type of interest, and will be apparent to those of skill in the art. When it is desired that the polynucleotide transgene be integrated into the chromosomal site of the endogenous gene, gene targeting is preferred. Briefly, when such a technique is to be utilized, vectors containing some nucleotide sequences homologous to the endogenous gene are designed for the purpose of integrating, via homologous recombination with chromosomal sequences, into and disrupting the function of the nucleotide sequence of the endogenous gene. The transgene may also be selectively introduced into a particular cell type, thus inactivating the endogenous gene in only that cell type, by following, for example, the teaching of Gu et al. (Gu et al., Science 265:103-106 (1994)). The regulatory sequences required for such a cell-type specific inactivation will depend upon the particular cell type of interest, and will be apparent to those of skill in the art.

Once transgenic animals have been generated, the expression of the recombinant gene may be assayed utilizing standard techniques. Initial screening may be accomplished by Southern blot analysis or PCR techniques to analyze animal tissues to verify that integration of the transgene has taken place. The level of mRNA expression of the transgene in the tissues of the transgenic animals may also be assessed using techniques which include, but are not limited to, Northern blot analysis of tissue samples obtained from the animal, *in situ* hybridization analysis, and reverse transcriptase-PCR (rt-PCR). Samples of transgenic gene-expressing tissue may also be evaluated immunocytochemically or immunohistochemically using antibodies specific for the transgene product.

Once the founder animals are produced, they may be bred, inbred, outbred, or crossbred to produce colonies of the particular animal. Examples of such breeding strategies include, but are not limited to: outbreeding of founder animals with more

than one integration site in order to establish separate lines; inbreeding of separate lines in order to produce compound transgenics that express the transgene at higher levels because of the effects of additive expression of each transgene; crossing of heterozygous transgenic animals to produce animals homozygous for a given
5 integration site in order to both augment expression and eliminate the need for screening of animals by DNA analysis; crossing of separate homozygous lines to produce compound heterozygous or homozygous lines; and breeding to place the transgene on a distinct background that is appropriate for an experimental model of interest.

10 Transgenic animals of the invention have uses which include, but are not limited to, animal model systems useful in elaborating the biological function of polypeptides of the present invention, studying conditions and/or disorders associated with aberrant expression, and in screening for compounds effective in ameliorating such conditions and/or disorders.

15

Example 30: Knock-Out Animals.

Endogenous gene expression can also be reduced by inactivating or "knocking out" the gene and/or its promoter using targeted homologous recombination. (*E.g.*, see Smithies et al., Nature 317:230-234 (1985); Thomas & Capecchi, Cell 51:503-
20 512 (1987); Thompson et al., Cell 5:313-321 (1989); each of which is incorporated by reference herein in its entirety). For example, a mutant, non-functional polynucleotide of the invention (or a completely unrelated DNA sequence) flanked by DNA homologous to the endogenous polynucleotide sequence (either the coding regions or regulatory regions of the gene) can be used, with or without a selectable
25 marker and/or a negative selectable marker, to transfect cells that express polypeptides of the invention *in vivo*. In another embodiment, techniques known in the art are used to generate knockouts in cells that contain, but do not express the gene of interest. Insertion of the DNA construct, via targeted homologous recombination, results in inactivation of the targeted gene. Such approaches are particularly suited in
30 research and agricultural fields where modifications to embryonic stem cells can be used to generate animal offspring with an inactive targeted gene (*e.g.*, see Thomas &

Capecchi 1987 and Thompson 1989, *supra*). However this approach can be routinely adapted for use in humans provided the recombinant DNA constructs are directly administered or targeted to the required site *in vivo* using appropriate viral vectors that will be apparent to those of skill in the art.

5 In further embodiments of the invention, cells that are genetically engineered to express the polypeptides of the invention, or alternatively, that are genetically engineered not to express the polypeptides of the invention (e.g., knockouts) are administered to a patient *in vivo*. Such cells may be obtained from the patient (i.e., animal, including human) or an MHC compatible donor and can include, but are not
10 limited to fibroblasts, bone marrow cells, blood cells (e.g., lymphocytes), adipocytes, muscle cells, endothelial cells etc. The cells are genetically engineered *in vitro* using recombinant DNA techniques to introduce the coding sequence of polypeptides of the invention into the cells, or alternatively, to disrupt the coding sequence and/or endogenous regulatory sequence associated with the polypeptides of the invention,
15 e.g., by transduction (using viral vectors, and preferably vectors that integrate the transgene into the cell genome) or transfection procedures, including, but not limited to, the use of plasmids, cosmids, YACs, naked DNA, electroporation, liposomes, etc. The coding sequence of the polypeptides of the invention can be placed under the control of a strong constitutive or inducible promoter or promoter/enhancer to achieve
20 expression, and preferably secretion, of the polypeptides of the invention. The engineered cells which express and preferably secrete the polypeptides of the invention can be introduced into the patient systemically, e.g., in the circulation, or intraperitoneally.

 Alternatively, the cells can be incorporated into a matrix and implanted in the
25 body, e.g., genetically engineered fibroblasts can be implanted as part of a skin graft; genetically engineered endothelial cells can be implanted as part of a lymphatic or vascular graft. (See, for example, Anderson et al. U.S. Patent No. 5,399,349; and Mulligan & Wilson, U.S. Patent No. 5,460,959 each of which is incorporated by reference herein in its entirety).

30 When the cells to be administered are non-autologous or non-MHC compatible cells, they can be administered using well known techniques which

prevent the development of a host immune response against the introduced cells. For example, the cells may be introduced in an encapsulated form which, while allowing for an exchange of components with the immediate extracellular environment, does not allow the introduced cells to be recognized by the host immune system.

5 Transgenic and "knock-out" animals of the invention have uses which include, but are not limited to, animal model systems useful in elaborating the biological function of polypeptides of the present invention, studying conditions and/or disorders associated with aberrant expression, and in screening for compounds effective in ameliorating such conditions and/or disorders.

10

Example 31: Production of an Antibody

a) Hybridoma Technology

The antibodies of the present invention can be prepared by a variety of methods. (See, Current Protocols, Chapter 2.) As one example of such methods, cells
15 expressing polypeptide(s) of the invention are administered to an animal to induce the production of sera containing polyclonal antibodies. In a preferred method, a preparation of polypeptide(s) of the invention is prepared and purified to render it substantially free of natural contaminants. Such a preparation is then introduced into an animal in order to produce polyclonal antisera of greater specific activity.

20 Monoclonal antibodies specific for polypeptide(s) of the invention are prepared using hybridoma technology. (Kohler et al., Nature 256:495 (1975); Kohler et al., Eur. J. Immunol. 6:511 (1976); Kohler et al., Eur. J. Immunol. 6:292 (1976); Hammerling et al., in: Monoclonal Antibodies and T-Cell Hybridomas, Elsevier, N.Y., pp. 563-681 (1981)). In general, an animal (preferably a mouse) is immunized
25 with polypeptide(s) of the invention or, more preferably, with a secreted polypeptide-expressing cell. Such polypeptide-expressing cells are cultured in any suitable tissue culture medium, preferably in Earle's modified Eagle's medium supplemented with 10% fetal bovine serum (inactivated at about 56°C), and supplemented with about 10 g/l of nonessential amino acids, about 1,000 U/ml of penicillin, and about 100 µg/ml
30 of streptomycin.

The splenocytes of such mice are extracted and fused with a suitable myeloma cell line. Any suitable myeloma cell line may be employed in accordance with the present invention; however, it is preferable to employ the parent myeloma cell line (SP2O), available from the ATCC. After fusion, the resulting hybridoma cells are
5 selectively maintained in HAT medium, and then cloned by limiting dilution as described by Wands et al. (Gastroenterology 80:225-232 (1981)). The hybridoma cells obtained through such a selection are then assayed to identify clones which secrete antibodies capable of binding the polypeptide(s) of the invention.

Alternatively, additional antibodies capable of binding to polypeptide(s) of the
10 invention can be produced in a two-step procedure using anti-idiotypic antibodies. Such a method makes use of the fact that antibodies are themselves antigens, and therefore, it is possible to obtain an antibody which binds to a second antibody. In accordance with this method, protein specific antibodies are used to immunize an animal, preferably a mouse. The splenocytes of such an animal are then used to
15 produce hybridoma cells, and the hybridoma cells are screened to identify clones which produce an antibody whose ability to bind to the protein-specific antibody can be blocked by polypeptide(s) of the invention. Such antibodies comprise anti-idiotypic antibodies to the protein-specific antibody and are used to immunize an animal to induce formation of further protein-specific antibodies.

20 For in vivo use of antibodies in humans, an antibody is "humanized". Such antibodies can be produced using genetic constructs derived from hybridoma cells producing the monoclonal antibodies described above. Methods for producing chimeric and humanized antibodies are known in the art and are discussed herein. (See, for review, Morrison, Science 229:1202 (1985); Oi et al., BioTechniques 4:214
25 (1986); Cabilly et al., U.S. Patent No. 4,816,567; Taniguchi et al., EP 171496; Morrison et al., EP 173494; Neuberger et al., WO 8601533; Robinson et al., WO 8702671; Boulianne et al., Nature 312:643 (1984); Neuberger et al., Nature 314:268 (1985).)

30 b) Isolation Of Antibody Fragments Directed Against
Polypeptide(s) From A Library Of scFvs

Naturally occurring V-genes isolated from human PBLs are constructed into a library of antibody fragments which contain reactivities against polypeptide(s) of the invention to which the donor may or may not have been exposed (see e.g., U.S. Patent 5,885,793 incorporated herein by reference in its entirety).

5 Rescue of the Library.

A library of scFvs is constructed from the RNA of human PBLs as described in PCT publication WO 92/01047. To rescue phage displaying antibody fragments, approximately 109 E. coli harboring the phagemid are used to inoculate 50 ml of 2xTY containing 1% glucose and 100 µg/ml of ampicillin (2xTY-AMP-GLU) and
10 grown to an O.D. of 0.8 with shaking. Five ml of this culture is used to inoculate 50 ml of 2xTY-AMP-GLU, 2 x 10⁸ TU of delta gene 3 helper (M13 delta gene III, see PCT publication WO 92/01047) are added and the culture incubated at 37°C for 45 minutes without shaking and then at 37°C for 45 minutes with shaking. The culture is centrifuged at 4000 r.p.m. for 10 min. and the pellet resuspended in 2 liters of 2xTY
15 containing 100 µg/ml ampicillin and 50 µg/ml kanamycin and grown overnight. Phage are prepared as described in PCT publication WO 92/01047.

M13 delta gene III is prepared as follows: M13 delta gene III helper phage does not encode gene III protein, hence the phage(mid) displaying antibody fragments have a greater avidity of binding to antigen. Infectious M13 delta gene III
20 particles are made by growing the helper phage in cells harboring a pUC19 derivative supplying the wild type gene III protein during phage morphogenesis. The culture is incubated for 1 hour at 37° C without shaking and then for a further hour at 37°C with shaking. Cells are spun down (IEC-Centra 8,400 r.p.m. for 10 min), resuspended in 300 ml 2xTY broth containing 100 µg ampicillin/ml and 25 µg kanamycin/ml (2xTY-
25 AMP-KAN) and grown overnight, shaking at 37°C. Phage particles are purified and concentrated from the culture medium by two PEG-precipitations (Sambrook et al., 1990), resuspended in 2 ml PBS and passed through a 0.45 µm filter (Minisart NML; Sartorius) to give a final concentration of approximately 10¹³ transducing units/ml (ampicillin-resistant clones).

30 Panning of the Library.

Immunotubes (Nunc) are coated overnight in PBS with 4 ml of either 100 µg/ml or 10 µg/ml of a polypeptide of the present invention. Tubes are blocked with 2% Marvel-PBS for 2 hours at 37°C and then washed 3 times in PBS. Approximately 1013 TU of phage is applied to the tube and incubated for 30 minutes at room temperature tumbling on an over and under turntable and then left to stand for another 1.5 hours. Tubes are washed 10 times with PBS 0.1% Tween-20 and 10 times with PBS. Phage are eluted by adding 1 ml of 100 mM triethylamine and rotating 15 minutes on an under and over turntable after which the solution is immediately neutralized with 0.5 ml of 1.0M Tris-HCl, pH 7.4. Phage are then used to infect 10 ml of mid-log E. coli TG1 by incubating eluted phage with bacteria for 30 minutes at 37°C. The E. coli are then plated on TYE plates containing 1% glucose and 100 µg/ml ampicillin. The resulting bacterial library is then rescued with delta gene 3 helper phage as described above to prepare phage for a subsequent round of selection. This process is then repeated for a total of 4 rounds of affinity purification with tube-washing increased to 20 times with PBS, 0.1% Tween-20 and 20 times with PBS for rounds 3 and 4.

Characterization of Binders.

Eluted phage from the 3rd and 4th rounds of selection are used to infect E. coli HB 2151 and soluble scFv is produced (Marks, et al., 1991) from single colonies for assay. ELISAs are performed with microtitre plates coated with either 10 pg/ml of the polypeptide of the present invention in 50 mM bicarbonate pH 9.6. Clones positive in ELISA are further characterized by PCR fingerprinting (see, e.g., PCT publication WO 92/01047) and then by sequencing. These ELISA positive clones may also be further characterized by techniques known in the art, such as, for example, epitope mapping, binding affinity, receptor signal transduction, ability to block or competitively inhibit antibody/antigen binding, and competitive agonistic or antagonistic activity.

Example 32: Assays Detecting Stimulation or Inhibition of B cell Proliferation and Differentiation

Generation of functional humoral immune responses requires both soluble and cognate signaling between B-lineage cells and their microenvironment. Signals may impart a positive stimulus that allows a B-lineage cell to continue its programmed development, or a negative stimulus that instructs the cell to arrest its current developmental pathway. To date, numerous stimulatory and inhibitory signals have been found to influence B cell responsiveness including IL-2, IL-4, IL-5, IL-6, IL-7, IL-10, IL-13, IL-14 and IL-15. Interestingly, these signals are by themselves weak effectors but can, in combination with various co-stimulatory proteins, induce activation, proliferation, differentiation, homing, tolerance and death among B cell populations.

One of the best studied classes of B-cell co-stimulatory proteins is the TNF-superfamily. Within this family CD40, CD27, and CD30 along with their respective ligands CD154, CD70, and CD153 have been found to regulate a variety of immune responses. Assays which allow for the detection and/or observation of the proliferation and differentiation of these B-cell populations and their precursors are valuable tools in determining the effects various proteins may have on these B-cell populations in terms of proliferation and differentiation. Listed below are two assays designed to allow for the detection of the differentiation, proliferation, or inhibition of B-cell populations and their precursors.

In Vitro Assay- Purified polypeptides of the invention, or truncated forms thereof, is assessed for its ability to induce activation, proliferation, differentiation or inhibition and/or death in B-cell populations and their precursors. The activity of the polypeptides of the invention on purified human tonsillar B cells, measured qualitatively over the dose range from 0.1 to 10,000 ng/mL, is assessed in a standard B-lymphocyte co-stimulation assay in which purified tonsillar B cells are cultured in the presence of either formalin-fixed *Staphylococcus aureus* Cowan I (SAC) or immobilized anti-human IgM antibody as the priming agent. Second signals such as IL-2 and IL-15 synergize with SAC and IgM crosslinking to elicit B cell proliferation as measured by tritiated-thymidine incorporation. Novel synergizing agents can be readily identified using this assay. The assay involves isolating human tonsillar B cells by magnetic bead (MACS) depletion of CD3-positive

cells. The resulting cell population is greater than 95% B cells as assessed by expression of CD45R(B220).

Various dilutions of each sample are placed into individual wells of a 96-well plate to which are added 10^5 B-cells suspended in culture medium (RPMI 1640 containing 10%
5 FBS, 5×10^{-5} M 2ME, 100U/ml penicillin, 10ug/ml streptomycin, and 10^{-5} dilution of SAC) in a total volume of 150ul. Proliferation or inhibition is quantitated by a 20h pulse (1uCi/well) with 3H-thymidine (6.7 Ci/mM) beginning 72h post factor addition. The positive and negative controls are IL2 and medium respectively.

In Vivo Assay- BALB/c mice are injected (i.p.) twice per day with buffer only, or
10 2 mg/Kg of a polypeptide of the invention, or truncated forms thereof. Mice receive this treatment for 4 consecutive days, at which time they are sacrificed and various tissues and serum collected for analyses. Comparison of H&E sections from normal spleens and spleens treated with polypeptides of the invention identify the results of the activity of the polypeptides on spleen cells, such as the diffusion of peri-arterial lymphatic sheaths,
15 and/or significant increases in the nucleated cellularity of the red pulp regions, which may indicate the activation of the differentiation and proliferation of B-cell populations. Immunohistochemical studies using a B cell marker, anti-CD45R(B220), are used to determine whether any physiological changes to splenic cells, such as splenic disorganization, are due to increased B-cell representation within loosely defined B-cell
20 zones that infiltrate established T-cell regions.

Flow cytometric analyses of the spleens from mice treated with polypeptide is used to indicate whether the polypeptide specifically increases the proportion of ThB+, CD45R(B220)dull B cells over that which is observed in control mice.

Likewise, a predicted consequence of increased mature B-cell representation in
25 vivo is a relative increase in serum Ig titers. Accordingly, serum IgM and IgA levels are compared between buffer and polypeptide-treated mice.

The studies described in this example tested activity of a polypeptide of the invention. However, one skilled in the art could easily modify the exemplified studies to test the activity of polynucleotides of the invention (e.g., gene therapy), agonists, and/or
30 antagonists of polynucleotides or polypeptides of the invention.

Example 33: T Cell Proliferation Assay

A CD3-induced proliferation assay is performed on PBMCs and is measured by the uptake of ^3H -thymidine. The assay is performed as follows. Ninety-six well plates are coated with 100 μl /well of mAb to CD3 (HIT3a, Pharmingen) or isotype-matched control mAb (B33.1) overnight at 4 degrees C (1 $\mu\text{g}/\text{ml}$ in .05M bicarbonate buffer, pH 9.5), then washed three times with PBS. PBMC are isolated by F/H gradient centrifugation from human peripheral blood and added to quadruplicate wells (5 x 10⁴/well) of mAb coated plates in RPMI containing 10% FCS and P/S in the presence of varying concentrations of polypeptides of the invention (total volume 200 μl). Relevant protein buffer and medium alone are controls. After 48 hr. culture at 37 degrees C, plates are spun for 2 min. at 1000 rpm and 100 μl of supernatant is removed and stored -20 degrees C for measurement of IL-2 (or other cytokines) if effect on proliferation is observed. Wells are supplemented with 100 μl of medium containing 0.5 μCi of ^3H -thymidine and cultured at 37 degrees C for 18-24 hr. Wells are harvested and incorporation of ^3H -thymidine used as a measure of proliferation. Anti-CD3 alone is the positive control for proliferation. IL-2 (100 U/ml) is also used as a control which enhances proliferation. Control antibody which does not induce proliferation of T cells is used as the negative controls for the effects of polypeptides of the invention.

The studies described in this example tested activity of polypeptides of the invention. However, one skilled in the art could easily modify the exemplified studies to test the activity of polynucleotides of the invention (e.g., gene therapy), agonists, and/or antagonists of polynucleotides or polypeptides of the invention.

Example 34: Effect of Polypeptides of the Invention on the Expression of MHC Class II, Costimulatory and Adhesion Molecules and Cell Differentiation of Monocytes and Monocyte-Derived Human Dendritic Cells

Dendritic cells are generated by the expansion of proliferating precursors found in the peripheral blood: adherent PBMC or elutriated monocytic fractions are cultured for 7-10 days with GM-CSF (50 ng/ml) and IL-4 (20 ng/ml). These dendritic cells have the characteristic phenotype of immature cells (expression of CD1, CD80, CD86, CD40 and

MHC class II antigens). Treatment with activating factors, such as TNF- α , causes a rapid change in surface phenotype (increased expression of MHC class I and II, costimulatory and adhesion molecules, downregulation of FC γ RII, upregulation of CD83). These changes correlate with increased antigen-presenting capacity and with functional
5 maturation of the dendritic cells.

FACS analysis of surface antigens is performed as follows. Cells are treated 1-3 days with increasing concentrations of polypeptides of the invention or LPS (positive control), washed with PBS containing 1% BSA and 0.02 mM sodium azide, and then incubated with 1:20 dilution of appropriate FITC- or PE-labeled monoclonal antibodies
10 for 30 minutes at 4 degrees C. After an additional wash, the labeled cells are analyzed by flow cytometry on a FACScan (Becton Dickinson).

Effect on the production of cytokines. Cytokines generated by dendritic cells, in particular IL-12, are important in the initiation of T-cell dependent immune responses. IL-
15 12 strongly influences the development of Th1 helper T-cell immune response, and induces cytotoxic T and NK cell function. An ELISA is used to measure the IL-12 release as follows. Dendritic cells (10^6 /ml) are treated with increasing concentrations of polypeptides of the invention for 24 hours. LPS (100 ng/ml) is added to the cell culture as positive control. Supernatants from the cell cultures are then collected and analyzed for
20 IL-12 content using commercial ELISA kit (e.g., R & D Systems (Minneapolis, MN)). The standard protocols provided with the kits are used.

Effect on the expression of MHC Class II, costimulatory and adhesion molecules. Three major families of cell surface antigens can be identified on monocytes: adhesion
25 molecules, molecules involved in antigen presentation, and Fc receptor. Modulation of the expression of MHC class II antigens and other costimulatory molecules, such as B7 and ICAM-1, may result in changes in the antigen presenting capacity of monocytes and ability to induce T cell activation. Increase expression of Fc receptors may correlate with improved monocyte cytotoxic activity, cytokine release and phagocytosis.

30 FACS analysis is used to examine the surface antigens as follows. Monocytes are treated 1-5 days with increasing concentrations of polypeptides of the invention or LPS

(positive control), washed with PBS containing 1% BSA and 0.02 mM sodium azide, and then incubated with 1:20 dilution of appropriate FITC- or PE-labeled monoclonal antibodies for 30 minutes at 4 degreesC. After an additional wash, the labeled cells are analyzed by flow cytometry on a FACScan (Becton Dickinson).

5

Monocyte activation and/or increased survival. Assays for molecules that activate (or alternatively, inactivate) monocytes and/or increase monocyte survival (or alternatively, decrease monocyte survival) are known in the art and may routinely be applied to determine whether a molecule of the invention functions as an inhibitor or
10 activator of monocytes. Polypeptides, agonists, or antagonists of the invention can be screened using the three assays described below. For each of these assays, Peripheral blood mononuclear cells (PBMC) are purified from single donor leukopacks (American Red Cross, Baltimore, MD) by centrifugation through a Histopaque gradient (Sigma). Monocytes are isolated from PBMC by counterflow centrifugal elutriation.

15

Monocyte Survival Assay. Human peripheral blood monocytes progressively lose viability when cultured in absence of serum or other stimuli. Their death results from internally regulated process (apoptosis). Addition to the culture of activating factors, such as TNF-alpha dramatically improves cell survival and prevents DNA fragmentation.
20 Propidium iodide (PI) staining is used to measure apoptosis as follows. Monocytes are cultured for 48 hours in polypropylene tubes in serum-free medium (positive control), in the presence of 100 ng/ml TNF-alpha (negative control), and in the presence of varying concentrations of the compound to be tested. Cells are suspended at a concentration of 2×10^6 /ml in PBS containing PI at a final concentration of 5 μ g/ml, and then incubated at room
25 temperature for 5 minutes before FACScan analysis. PI uptake has been demonstrated to correlate with DNA fragmentation in this experimental paradigm.

Effect on cytokine release. An important function of monocytes/macrophages is their regulatory activity on other cellular populations of the immune system through the
30 release of cytokines after stimulation. An ELISA to measure cytokine release is performed as follows. Human monocytes are incubated at a density of 5×10^5 cells/ml with

increasing concentrations of the a polypeptide of the invention and under the same conditions, but in the absence of the polypeptide. For IL-12 production, the cells are primed overnight with IFN (100 U/ml) in presence of a polypeptide of the invention. LPS (10 ng/ml) is then added. Conditioned media are collected after 24h and kept frozen until
5 use. Measurement of TNF-alpha, IL-10, MCP-1 and IL-8 is then performed using a commercially available ELISA kit (e.g., R & D Systems (Minneapolis, MN)) and applying the standard protocols provided with the kit.

Oxidative burst. Purified monocytes are plated in 96-w plate at 2×10^5 cell/well.
10 Increasing concentrations of polypeptides of the invention are added to the wells in a total volume of 0.2 ml culture medium (RPMI 1640 + 10% FCS, glutamine and antibiotics). After 3 days incubation, the plates are centrifuged and the medium is removed from the wells. To the macrophage monolayers, 0.2 ml per well of phenol red solution (140 mM NaCl, 10 mM potassium phosphate buffer pH 7.0, 5.5 mM dextrose, 0.56 mM phenol red
15 and 19 U/ml of HRPO) is added, together with the stimulant (200 nM PMA). The plates are incubated at 37°C for 2 hours and the reaction is stopped by adding 20 µl 1N NaOH per well. The absorbance is read at 610 nm. To calculate the amount of H_2O_2 produced by the macrophages, a standard curve of a H_2O_2 solution of known molarity is performed for each experiment.

20 The studies described in this example tested activity of a polypeptide of the invention. However, one skilled in the art could easily modify the exemplified studies to test the activity of polypeptides, polynucleotides (e.g., gene therapy), agonists, and/or antagonists of the invention.

25 **Example 35: Biological Effects of Polypeptides of the Invention**

Astrocyte and Neuronal Assays

Recombinant polypeptides of the invention, expressed in *Escherichia coli* and purified as described above, can be tested for activity in promoting the survival, neurite outgrowth, or phenotypic differentiation of cortical neuronal cells and for inducing the
30 proliferation of glial fibrillary acidic protein immunopositive cells, astrocytes. The selection of cortical cells for the bioassay is based on the prevalent expression of FGF-1

and FGF-2 in cortical structures and on the previously reported enhancement of cortical neuronal survival resulting from FGF-2 treatment. A thymidine incorporation assay, for example, can be used to elucidate a polypeptide of the invention's activity on these cells.

Moreover, previous reports describing the biological effects of FGF-2 (basic FGF) on cortical or hippocampal neurons *in vitro* have demonstrated increases in both neuron survival and neurite outgrowth (Walicke et al., "Fibroblast growth factor promotes survival of dissociated hippocampal neurons and enhances neurite extension." *Proc. Natl. Acad. Sci. USA* 83:3012-3016. (1986), assay herein incorporated by reference in its entirety). However, reports from experiments done on PC-12 cells suggest that these two responses are not necessarily synonymous and may depend on not only which FGF is being tested but also on which receptor(s) are expressed on the target cells. Using the primary cortical neuronal culture paradigm, the ability of a polypeptide of the invention to induce neurite outgrowth can be compared to the response achieved with FGF-2 using, for example, a thymidine incorporation assay.

Fibroblast and endothelial cell assays

Human lung fibroblasts are obtained from Clonetics (San Diego, CA) and maintained in growth media from Clonetics. Dermal microvascular endothelial cells are obtained from Cell Applications (San Diego, CA). For proliferation assays, the human lung fibroblasts and dermal microvascular endothelial cells can be cultured at 5,000 cells/well in a 96-well plate for one day in growth medium. The cells are then incubated for one day in 0.1% BSA basal medium. After replacing the medium with fresh 0.1% BSA medium, the cells are incubated with the test proteins for 3 days. Alamar Blue (Alamar Biosciences, Sacramento, CA) is added to each well to a final concentration of 10%. The cells are incubated for 4 hr. Cell viability is measured by reading in a CytoFluor fluorescence reader. For the PGE₂ assays, the human lung fibroblasts are cultured at 5,000 cells/well in a 96-well plate for one day. After a medium change to 0.1% BSA basal medium, the cells are incubated with FGF-2 or polypeptides of the invention with or without IL-1 α for 24 hours. The supernatants are collected and assayed for PGE₂ by EIA

kit (Cayman, Ann Arbor, MI). For the IL-6 assays, the human lung fibroblasts are cultured at 5,000 cells/well in a 96-well plate for one day. After a medium change to 0.1% BSA basal medium, the cells are incubated with FGF-2 or with or without polypeptides of the invention IL-1 α for 24 hours. The supernatants are collected and
5 assayed for IL-6 by ELISA kit (Endogen, Cambridge, MA).

Human lung fibroblasts are cultured with FGF-2 or polypeptides of the invention for 3 days in basal medium before the addition of Alamar Blue to assess effects on growth of the fibroblasts. FGF-2 should show a stimulation at 10 - 2500 ng/ml which can be used to compare stimulation with polypeptides of the invention.

10

Parkinson Models.

The loss of motor function in Parkinson's disease is attributed to a deficiency of striatal dopamine resulting from the degeneration of the nigrostriatal dopaminergic projection neurons. An animal model for Parkinson's that has been extensively
15 characterized involves the systemic administration of 1-methyl-4 phenyl 1,2,3,6-tetrahydropyridine (MPTP). In the CNS, MPTP is taken-up by astrocytes and catabolized by monoamine oxidase B to 1-methyl-4-phenyl pyridine (MPP⁺) and released. Subsequently, MPP⁺ is actively accumulated in dopaminergic neurons by the high-affinity reuptake transporter for dopamine. MPP⁺ is then concentrated in mitochondria by the
20 electrochemical gradient and selectively inhibits nicotinamide adenine disphosphate: ubiquinone oxidoreductionase (complex I), thereby interfering with electron transport and eventually generating oxygen radicals.

It has been demonstrated in tissue culture paradigms that FGF-2 (basic FGF) has trophic activity towards nigral dopaminergic neurons (Ferrari et al., Dev. Biol. 1989).
25 Recently, Dr. Unsicker's group has demonstrated that administering FGF-2 in gel foam implants in the striatum results in the near complete protection of nigral dopaminergic neurons from the toxicity associated with MPTP exposure (Otto and Unsicker, J. Neuroscience, 1990).

Based on the data with FGF-2, polypeptides of the invention can be evaluated to
30 determine whether it has an action similar to that of FGF-2 in enhancing dopaminergic

neuronal survival *in vitro* and it can also be tested *in vivo* for protection of dopaminergic neurons in the striatum from the damage associated with MPTP treatment. The potential effect of a polypeptide of the invention is first examined *in vitro* in a dopaminergic neuronal cell culture paradigm. The cultures are prepared by dissecting the midbrain floor plate from gestation day 14 Wistar rat embryos. The tissue is dissociated with trypsin and seeded at a density of 200,000 cells/cm² on polyorthinine-laminin coated glass coverslips. The cells are maintained in Dulbecco's Modified Eagle's medium and F12 medium containing hormonal supplements (N1). The cultures are fixed with paraformaldehyde after 8 days *in vitro* and are processed for tyrosine hydroxylase, a specific marker for dopaminergic neurons, immunohistochemical staining. Dissociated cell cultures are prepared from embryonic rats. The culture medium is changed every third day and the factors are also added at that time.

Since the dopaminergic neurons are isolated from animals at gestation day 14, a developmental time which is past the stage when the dopaminergic precursor cells are proliferating, an increase in the number of tyrosine hydroxylase immunopositive neurons would represent an increase in the number of dopaminergic neurons surviving *in vitro*. Therefore, if a polypeptide of the invention acts to prolong the survival of dopaminergic neurons, it would suggest that the polypeptide may be involved in Parkinson's Disease.

The studies described in this example tested activity of a polypeptide of the invention. However, one skilled in the art could easily modify the exemplified studies to test the activity of polynucleotides (e.g., gene therapy), agonists, and/or antagonists of the invention.

Example 36: The Effect of Polypeptides of the Invention on the Growth of

Vascular Endothelial Cells

On day 1, human umbilical vein endothelial cells (HUVEC) are seeded at 2-5x10⁴ cells/35 mm dish density in M199 medium containing 4% fetal bovine serum (FBS), 16 units/ml heparin, and 50 units/ml endothelial cell growth supplements (ECGS, Biotechnology, Inc.). On day 2, the medium is replaced with M199 containing 10% FBS, 8 units/ml heparin. A polypeptide having the amino acid sequence of SEQ ID NO:Y, and positive controls, such as VEGF and basic FGF (bFGF) are added, at varying

concentrations. On days 4 and 6, the medium is replaced. On day 8, cell number is determined with a Coulter Counter.

An increase in the number of HUVEC cells indicates that the polypeptide of the invention may proliferate vascular endothelial cells.

- 5 The studies described in this example tested activity of a polypeptide of the invention. However, one skilled in the art could easily modify the exemplified studies to test the activity of polynucleotides (e.g., gene therapy), agonists, and/or antagonists of the invention.

10 **Example 37: Stimulatory Effect of Polypeptides of the Invention on the Proliferation of Vascular Endothelial Cells**

- For evaluation of mitogenic activity of growth factors, the colorimetric MTS (3-(4,5-dimethylthiazol-2-yl)-5-(3-carboxymethoxyphenyl)-2-(4-sulfophenyl)2H-tetrazolium) assay with the electron coupling reagent PMS (phenazine methosulfate) was performed (CellTiter 96 AQ, Promega). Cells are seeded in a 96-well plate (5,000 cells/well) in 0.1 mL serum-supplemented medium and are allowed to attach overnight. After serum-starvation for 12 hours in 0.5% FBS, conditions (bFGF, VEGF₁₆₅ or a polypeptide of the invention in 0.5% FBS) with or without Heparin (8 U/ml) are added to wells for 48 hours. 20 mg of MTS/PMS mixture (1:0.05) are added per well and allowed to incubate for 1 hour at 37°C before measuring the absorbance at 490 nm in an ELISA plate reader. Background absorbance from control wells (some media, no cells) is subtracted, and seven wells are performed in parallel for each condition. See, Leak *et al. In Vitro Cell. Dev. Biol.* 30A:512-518 (1994).
- 15
20

- The studies described in this example tested activity of a polypeptide of the invention. However, one skilled in the art could easily modify the exemplified studies to test the activity of polynucleotides (e.g., gene therapy), agonists, and/or antagonists of the invention.
- 25

Example 38: Inhibition of PDGF-induced Vascular Smooth Muscle Cell Proliferation Stimulatory Effect

HAoSMC proliferation can be measured, for example, by BrdUrd incorporation.

5 Briefly, subconfluent, quiescent cells grown on the 4-chamber slides are transfected with CRP or FITC-labeled AT2-3LP. Then, the cells are pulsed with 10% calf serum and 6 mg/ml BrdUrd. After 24 h, immunocytochemistry is performed by using BrdUrd Staining Kit (Zymed Laboratories). In brief, the cells are incubated with the biotinylated mouse anti-BrdUrd antibody at 4 degrees C for 2 h after being exposed to denaturing solution and

10 then incubated with the streptavidin-peroxidase and diaminobenzidine. After counterstaining with hematoxylin, the cells are mounted for microscopic examination, and the BrdUrd-positive cells are counted. The BrdUrd index is calculated as a percent of the BrdUrd-positive cells to the total cell number. In addition, the simultaneous detection of the BrdUrd staining (nucleus) and the FITC uptake (cytoplasm) is performed for

15 individual cells by the concomitant use of bright field illumination and dark field-UV fluorescent illumination. See, Hayashida et al., J. Biol. Chem. 6:271(36):21985-21992 (1996).

The studies described in this example tested activity of a polypeptide of the invention. However, one skilled in the art could easily modify the exemplified studies to

20 test the activity of polynucleotides (e.g., gene therapy), agonists, and/or antagonists of the invention.

Example 39: Stimulation of Endothelial Migration

This example will be used to explore the possibility that a polypeptide of the

25 invention may stimulate lymphatic endothelial cell migration.

Endothelial cell migration assays are performed using a 48 well microchemotaxis chamber (Neuroprobe Inc., Cabin John, MD; Falk, W., et al., J. Immunological Methods 1980;33:239-247). Polyvinylpyrrolidone-free polycarbonate filters with a pore size of 8 um (Nucleopore Corp. Cambridge, MA) are coated with 0.1% gelatin for at least 6 hours

30 at room temperature and dried under sterile air. Test substances are diluted to appropriate concentrations in M199 supplemented with 0.25% bovine serum albumin (BSA), and 25

ul of the final dilution is placed in the lower chamber of the modified Boyden apparatus. Subconfluent, early passage (2-6) HUVEC or BMEC cultures are washed and trypsinized for the minimum time required to achieve cell detachment. After placing the filter between lower and upper chamber, 2.5×10^5 cells suspended in 50 ul M199 containing 1% FBS are seeded in the upper compartment. The apparatus is then incubated for 5 hours at 37°C in a humidified chamber with 5% CO₂ to allow cell migration. After the incubation period, the filter is removed and the upper side of the filter with the non-migrated cells is scraped with a rubber policeman. The filters are fixed with methanol and stained with a Giemsa solution (Diff-Quick, Baxter, McGraw Park, IL). Migration is quantified by counting cells of three random high-power fields (40x) in each well, and all groups are performed in quadruplicate.


The studies described in this example tested activity of a polypeptide of the invention. However, one skilled in the art could easily modify the exemplified studies to test the activity of polynucleotides (e.g., gene therapy), agonists, and/or antagonists of the invention.

Example 40: Stimulation of Nitric Oxide Production by Endothelial Cells

Nitric oxide released by the vascular endothelium is believed to be a mediator of vascular endothelium relaxation. Thus, activity of a polypeptide of the invention can be assayed by determining nitric oxide production by endothelial cells in response to the polypeptide.

Nitric oxide is measured in 96-well plates of confluent microvascular endothelial cells after 24 hours starvation and a subsequent 4 hr exposure to various levels of a positive control (such as VEGF-1) and the polypeptide of the invention. Nitric oxide in the medium is determined by use of the Griess reagent to measure total nitrite after reduction of nitric oxide-derived nitrate by nitrate reductase. The effect of the polypeptide of the invention on nitric oxide release is examined on HUVEC.

Briefly, NO release from cultured HUVEC monolayer is measured with a NO-specific polarographic electrode connected to a NO meter (Iso-NO, World Precision Instruments Inc.) (1049). Calibration of the NO elements is performed according to the following equation:



The standard calibration curve is obtained by adding graded concentrations of KNO_2 (0, 5, 10, 25, 50, 100, 250, and 500 nmol/L) into the calibration solution containing KI and H_2SO_4 . The specificity of the Iso-NO electrode to NO is previously determined by measurement of NO from authentic NO gas (1050). The culture medium is removed and HUVECs are washed twice with Dulbecco's phosphate buffered saline. The cells are then bathed in 5 ml of filtered Krebs-Henseleit solution in 6-well plates, and the cell plates are kept on a slide warmer (Lab Line Instruments Inc.) To maintain the temperature at 37°C. The NO sensor probe is inserted vertically into the wells, keeping the tip of the electrode 2 mm under the surface of the solution, before addition of the different conditions.

S-nitroso acetyl penicillamin (SNAP) is used as a positive control. The amount of released NO is expressed as picomoles per 1×10^6 endothelial cells. All values reported are means of four to six measurements in each group (number of cell culture wells). See, Leak *et al. Biochem. and Biophys. Res. Comm.* 217:96-105 (1995).

The studies described in this example tested activity of polypeptides of the invention. However, one skilled in the art could easily modify the exemplified studies to test the activity of polynucleotides (e.g., gene therapy), agonists, and/or antagonists of the invention.

Example 41: Effect of Polypeptides of the Invention on Cord Formation in Angiogenesis

Another step in angiogenesis is cord formation, marked by differentiation of endothelial cells. This bioassay measures the ability of microvascular endothelial cells to form capillary-like structures (hollow structures) when cultured *in vitro*.

CADMEC (microvascular endothelial cells) are purchased from Cell Applications, Inc. as proliferating (passage 2) cells and are cultured in Cell Applications' CADMEC Growth Medium and used at passage 5. For the *in vitro* angiogenesis assay, the wells of a 48-well cell culture plate are coated with Cell Applications' Attachment Factor Medium (200 ml/well) for 30 min. at 37°C. CADMEC are seeded onto the coated wells at 7,500 cells/well and cultured overnight in Growth Medium. The Growth Medium is then replaced with 300 mg Cell Applications' Chord Formation Medium containing control

buffer or a polypeptide of the invention (0.1 to 100 ng/ml) and the cells are cultured for an additional 48 hr. The numbers and lengths of the capillary-like chords are quantitated through use of the Boeckeler VIA-170 video image analyzer. All assays are done in triplicate.

- 5 Commercial (R&D) VEGF (50 ng/ml) is used as a positive control. b-esteradiol (1 ng/ml) is used as a negative control. The appropriate buffer (without protein) is also utilized as a control.

 The studies described in this example tested activity of a polypeptide of the invention. However, one skilled in the art could easily modify the exemplified studies to
10 test the activity of polynucleotides (e.g., gene therapy), agonists, and/or antagonists of the invention.

Example 42: Angiogenic Effect on Chick Chorioallantoic Membrane

 Chick chorioallantoic membrane (CAM) is a well-established system to examine
15 angiogenesis. Blood vessel formation on CAM is easily visible and quantifiable. The ability of polypeptides of the invention to stimulate angiogenesis in CAM can be examined.

 Fertilized eggs of the White Leghorn chick (*Gallus gallus*) and the Japanese quail (*Coturnix coturnix*) are incubated at 37.8°C and 80% humidity. Differentiated CAM of
20 16-day-old chick and 13-day-old quail embryos is studied with the following methods.

 On Day 4 of development, a window is made into the egg shell of chick eggs. The embryos are checked for normal development and the eggs sealed with cellotape. They are further incubated until Day 13. Thermanox coverslips (Nunc, Naperville, IL) are cut into disks of about 5 mm in diameter. Sterile and salt-free growth factors are dissolved in
25 distilled water and about 3.3 mg/ 5 ml are pipetted on the disks. After air-drying, the inverted disks are applied on CAM. After 3 days, the specimens are fixed in 3% glutaraldehyde and 2% formaldehyde and rinsed in 0.12 M sodium cacodylate buffer. They are photographed with a stereo microscope [Wild M8] and embedded for semi- and ultrathin sectioning as described above. Controls are performed with carrier disks alone.
30 The studies described in this example tested activity of a polypeptide of the invention. However, one skilled in the art could easily modify the exemplified studies to

test the activity of polynucleotides (e.g., gene therapy), agonists, and/or antagonists of the invention.

Example 43: Angiogenesis Assay Using a Matrigel Implant in Mouse

5 *In vivo* angiogenesis assay of a polypeptide of the invention measures the ability of an existing capillary network to form new vessels in an implanted capsule of murine extracellular matrix material (Matrigel). The protein is mixed with the liquid Matrigel at 4 degree C and the mixture is then injected subcutaneously in mice where it solidifies. After 7 days, the solid “plug” of Matrigel is removed and examined for the presence of new
10 blood vessels. Matrigel is purchased from Becton Dickinson Labware/Collaborative Biomedical Products.

When thawed at 4 degree C the Matrigel material is a liquid. The Matrigel is mixed with a polypeptide of the invention at 150 ng/ml at 4 degrees C and drawn into cold 3 ml syringes. Female C57Bl/6 mice approximately 8 weeks old are injected with the
15 mixture of Matrigel and experimental protein at 2 sites at the midventral aspect of the abdomen (0.5 ml/site). After 7 days, the mice are sacrificed by cervical dislocation, the Matrigel plugs are removed and cleaned (i.e., all clinging membranes and fibrous tissue is removed). Replicate whole plugs are fixed in neutral buffered 10% formaldehyde, embedded in paraffin and used to produce sections for histological examination after
20 staining with Masson’s Trichrome. Cross sections from 3 different regions of each plug are processed. Selected sections are stained for the presence of vWF. The positive control for this assay is bovine basic FGF (150 ng/ml). Matrigel alone is used to determine basal levels of angiogenesis.

The studies described in this example tested activity of a polypeptide of the
25 invention. However, one skilled in the art could easily modify the exemplified studies to test the activity of polynucleotides (e.g., gene therapy), agonists, and/or antagonists of the invention.

Example 44: Rescue of Ischemia in Rabbit Lower Limb Model

To study the in vivo effects of polynucleotides and polypeptides of the invention on ischemia, a rabbit hindlimb ischemia model is created by surgical removal of one femoral arteries as described previously (Takeshita *et al.*, *Am J. Pathol* 147:1649-1660 (1995)). The excision of the femoral artery results in retrograde propagation of thrombus and occlusion of the external iliac artery. Consequently, blood flow to the ischemic limb is dependent upon collateral vessels originating from the internal iliac artery (Takeshita *et al.* *Am J. Pathol* 147:1649-1660 (1995)). An interval of 10 days is allowed for post-operative recovery of rabbits and development of endogenous collateral vessels. At 10 day post-operatively (day 0), after performing a baseline angiogram, the internal iliac artery of the ischemic limb is transfected with 500 mg naked expression plasmid containing a polynucleotide of the invention by arterial gene transfer technology using a hydrogel-coated balloon catheter as described (Riessen *et al.* *Hum Gene Ther.* 4:749-758 (1993); Leclerc *et al.* *J. Clin. Invest.* 90: 936-944 (1992)). When a polypeptide of the invention is used in the treatment, a single bolus of 500 mg polypeptide of the invention or control is delivered into the internal iliac artery of the ischemic limb over a period of 1 min. through an infusion catheter. On day 30, various parameters are measured in these rabbits: (a) BP ratio - The blood pressure ratio of systolic pressure of the ischemic limb to that of normal limb; (b) Blood Flow and Flow Reserve - Resting FL: the blood flow during undilated condition and Max FL: the blood flow during fully dilated condition (also an indirect measure of the blood vessel amount) and Flow Reserve is reflected by the ratio of max FL: resting FL; (c) Angiographic Score - This is measured by the angiogram of collateral vessels. A score is determined by the percentage of circles in an overlaying grid that with crossing opacified arteries divided by the total number in the rabbit thigh; (d) Capillary density - The number of collateral capillaries determined in light microscopic sections taken from hindlimbs.

The studies described in this example tested activity of polynucleotides and polypeptides of the invention. However, one skilled in the art could easily modify the exemplified studies to test the agonists, and/or antagonists of the invention.

Example 45: Effect of Polypeptides of the Invention on Vasodilation

Since dilation of vascular endothelium is important in reducing blood pressure, the ability of polypeptides of the invention to affect the blood pressure in spontaneously hypertensive rats (SHR) is examined. Increasing doses (0, 10, 30, 100, 300, and 900
5 mg/kg) of the polypeptides of the invention are administered to 13-14 week old spontaneously hypertensive rats (SHR). Data are expressed as the mean +/- SEM. Statistical analysis are performed with a paired t-test and statistical significance is defined as $p < 0.05$ vs. the response to buffer alone.

The studies described in this example tested activity of a polypeptide of the
10 invention. However, one skilled in the art could easily modify the exemplified studies to test the activity of polynucleotides (e.g., gene therapy), agonists, and/or antagonists of the invention.

Example 46: Rat Ischemic Skin Flap Model

15 The evaluation parameters include skin blood flow, skin temperature, and factor VIII immunohistochemistry or endothelial alkaline phosphatase reaction. Expression of polypeptides of the invention, during the skin ischemia, is studied using in situ hybridization.

The study in this model is divided into three parts as follows:

- 20
- a) Ischemic skin
 - b) Ischemic skin wounds
 - c) Normal wounds

The experimental protocol includes:

- 25 a) Raising a 3x4 cm, single pedicle full-thickness random skin flap (myocutaneous flap over the lower back of the animal).
- b) An excisional wounding (4-6 mm in diameter) in the ischemic skin (skin-flap).
- c) Topical treatment with a polypeptide of the invention of the excisional wounds (day 0, 1, 2, 3, 4 post-wounding) at the following various dosage ranges: 1mg to 100 mg.
- 30 d) Harvesting the wound tissues at day 3, 5, 7, 10, 14 and 21 post-wounding for histological, immunohistochemical, and in situ studies.

The studies described in this example tested activity of a polypeptide of the invention. However, one skilled in the art could easily modify the exemplified studies to test the activity of polynucleotides (e.g., gene therapy), agonists, and/or antagonists of the invention.

5

Example 47: Peripheral Arterial Disease Model

Angiogenic therapy using a polypeptide of the invention is a novel therapeutic strategy to obtain restoration of blood flow around the ischemia in case of peripheral
10 arterial diseases. The experimental protocol includes:

a) One side of the femoral artery is ligated to create ischemic muscle of the hindlimb, the other side of hindlimb serves as a control.

b) a polypeptide of the invention, in a dosage range of 20 mg - 500 mg, is delivered intravenously and/or intramuscularly 3 times (perhaps more) per week for 2-3
15 weeks.

c) The ischemic muscle tissue is collected after ligation of the femoral artery at 1, 2, and 3 weeks for the analysis of expression of a polypeptide of the invention and histology. Biopsy is also performed on the other side of normal muscle of the contralateral hindlimb.

20 The studies described in this example tested activity of a polypeptide of the invention. However, one skilled in the art could easily modify the exemplified studies to test the activity of polynucleotides (e.g., gene therapy), agonists, and/or antagonists of the invention.

25 **Example 48: Ischemic Myocardial Disease Model**

A polypeptide of the invention is evaluated as a potent mitogen capable of stimulating the development of collateral vessels, and restructuring new vessels after coronary artery occlusion. Alteration of expression of the polypeptide is investigated in
30 situ. The experimental protocol includes:

a) The heart is exposed through a left-side thoracotomy in the rat. Immediately, the left coronary artery is occluded with a thin suture (6-0) and the thorax is closed.

b) a polypeptide of the invention, in a dosage range of 20 mg - 500 mg, is delivered intravenously and/or intramuscularly 3 times (perhaps more) per week for 2-4
5 weeks.

c) Thirty days after the surgery, the heart is removed and cross-sectioned for morphometric and in situ analyzes.

The studies described in this example tested activity of a polypeptide of the invention. However, one skilled in the art could easily modify the exemplified studies to
10 test the activity of polynucleotides (e.g., gene therapy), agonists, and/or antagonists of the invention.

Example 49: Rat Corneal Wound Healing Model

15 This animal model shows the effect of a polypeptide of the invention on neovascularization. The experimental protocol includes:

a) Making a 1-1.5 mm long incision from the center of cornea into the stromal layer.

b) Inserting a spatula below the lip of the incision facing the outer corner of
20 the eye.

c) Making a pocket (its base is 1-1.5 mm from the edge of the eye).

d) Positioning a pellet, containing 50ng- 5ug of a polypeptide of the invention, within the pocket.

e) Treatment with a polypeptide of the invention can also be applied topically
25 to the corneal wounds in a dosage range of 20mg - 500mg (daily treatment for five days).

The studies described in this example tested activity of a polypeptide of the invention. However, one skilled in the art could easily modify the exemplified studies to test the activity of polynucleotides (e.g., gene therapy), agonists, and/or antagonists of the invention.

Example 50: Diabetic Mouse and Glucocorticoid-Impaired Wound Healing Models

A. Diabetic db+/db+ Mouse Model.

5 To demonstrate that a polypeptide of the invention accelerates the healing process, the genetically diabetic mouse model of wound healing is used. The full thickness wound healing model in the db+/db+ mouse is a well characterized, clinically relevant and reproducible model of impaired wound healing. Healing of the diabetic wound is dependent on formation of granulation tissue and re-epithelialization rather than
10 contraction (Gartner, M.H. *et al.*, *J. Surg. Res.* 52:389 (1992); Greenhalgh, D.G. *et al.*, *Am. J. Pathol.* 136:1235 (1990)).

The diabetic animals have many of the characteristic features observed in Type II diabetes mellitus. Homozygous (db+/db+) mice are obese in comparison to their normal heterozygous (db+/+m) littermates. Mutant diabetic (db+/db+) mice have a single
15 autosomal recessive mutation on chromosome 4 (db+) (Coleman *et al.* *Proc. Natl. Acad. Sci. USA* 77:283-293 (1982)). Animals show polyphagia, polydipsia and polyuria. Mutant diabetic mice (db+/db+) have elevated blood glucose, increased or normal insulin levels, and suppressed cell-mediated immunity (Mandel *et al.*, *J. Immunol.* 120:1375 (1978); Debray-Sachs, M. *et al.*, *Clin. Exp. Immunol.* 51(1):1-7 (1983); Leiter *et al.*, *Am.*
20 *J. of Pathol.* 114:46-55 (1985)). Peripheral neuropathy, myocardial complications, and microvascular lesions, basement membrane thickening and glomerular filtration abnormalities have been described in these animals (Norido, F. *et al.*, *Exp. Neurol.* 83(2):221-232 (1984); Robertson *et al.*, *Diabetes* 29(1):60-67 (1980); Giacomelli *et al.*, *Lab Invest.* 40(4):460-473 (1979); Coleman, D.L., *Diabetes* 31 (Suppl):1-6 (1982)). These
25 homozygous diabetic mice develop hyperglycemia that is resistant to insulin analogous to human type II diabetes (Mandel *et al.*, *J. Immunol.* 120:1375-1377 (1978)).

The characteristics observed in these animals suggests that healing in this model may be similar to the healing observed in human diabetes (Greenhalgh, *et al.*, *Am. J. of Pathol.* 136:1235-1246 (1990)).

30 Genetically diabetic female C57BL/KsJ (db+/db+) mice and their non-diabetic (db+/+m) heterozygous littermates are used in this study (Jackson Laboratories). The

animals are purchased at 6 weeks of age and are 8 weeks old at the beginning of the study. Animals are individually housed and received food and water ad libitum. All manipulations are performed using aseptic techniques. The experiments are conducted according to the rules and guidelines of Human Genome Sciences, Inc. Institutional
5 Animal Care and Use Committee and the Guidelines for the Care and Use of Laboratory Animals.

Wounding protocol is performed according to previously reported methods (Tsuboi, R. and Rifkin, D.B., *J. Exp. Med.* 172:245-251 (1990)). Briefly, on the day of wounding, animals are anesthetized with an intraperitoneal injection of Avertin (0.01
10 mg/mL), 2,2,2-tribromoethanol and 2-methyl-2-butanol dissolved in deionized water. The dorsal region of the animal is shaved and the skin washed with 70% ethanol solution and iodine. The surgical area is dried with sterile gauze prior to wounding. An 8 mm full-thickness wound is then created using a Keyes tissue punch. Immediately following wounding, the surrounding skin is gently stretched to eliminate wound expansion. The
15 wounds are left open for the duration of the experiment. Application of the treatment is given topically for 5 consecutive days commencing on the day of wounding. Prior to treatment, wounds are gently cleansed with sterile saline and gauze sponges.

Wounds are visually examined and photographed at a fixed distance at the day of surgery and at two day intervals thereafter. Wound closure is determined by daily
20 measurement on days 1-5 and on day 8. Wounds are measured horizontally and vertically using a calibrated Jameson caliper. Wounds are considered healed if granulation tissue is no longer visible and the wound is covered by a continuous epithelium.

A polypeptide of the invention is administered using at a range different doses, from 4mg to 500mg per wound per day for 8 days in vehicle. Vehicle control groups
25 received 50mL of vehicle solution.

Animals are euthanized on day 8 with an intraperitoneal injection of sodium pentobarbital (300mg/kg). The wounds and surrounding skin are then harvested for histology and immunohistochemistry. Tissue specimens are placed in 10% neutral buffered formalin in tissue cassettes between biopsy sponges for further processing.
30 Three groups of 10 animals each (5 diabetic and 5 non-diabetic controls) are evaluated: 1) Vehicle placebo control, 2) untreated group, and 3) treated group.

Wound closure is analyzed by measuring the area in the vertical and horizontal axis and obtaining the total square area of the wound. Contraction is then estimated by establishing the differences between the initial wound area (day 0) and that of post treatment (day 8). The wound area on day 1 is 64mm², the corresponding size of the dermal punch. Calculations are made using the following formula:

$$[\text{Open area on day 8}] - [\text{Open area on day 1}] / [\text{Open area on day 1}]$$

Specimens are fixed in 10% buffered formalin and paraffin embedded blocks are sectioned perpendicular to the wound surface (5mm) and cut using a Reichert-Jung microtome. Routine hematoxylin-eosin (H&E) staining is performed on cross-sections of bisected wounds. Histologic examination of the wounds are used to assess whether the healing process and the morphologic appearance of the repaired skin is altered by treatment with a polypeptide of the invention. This assessment included verification of the presence of cell accumulation, inflammatory cells, capillaries, fibroblasts, re-epithelialization and epidermal maturity (Greenhalgh, D.G. *et al.*, *Am. J. Pathol.* 136:1235 (1990)). A calibrated lens micrometer is used by a blinded observer.

Tissue sections are also stained immunohistochemically with a polyclonal rabbit anti-human keratin antibody using ABC Elite detection system. Human skin is used as a positive tissue control while non-immune IgG is used as a negative control. Keratinocyte growth is determined by evaluating the extent of reepithelialization of the wound using a calibrated lens micrometer.

Proliferating cell nuclear antigen/cyclin (PCNA) in skin specimens is demonstrated by using anti-PCNA antibody (1:50) with an ABC Elite detection system. Human colon cancer can serve as a positive tissue control and human brain tissue can be used as a negative tissue control. Each specimen includes a section with omission of the primary antibody and substitution with non-immune mouse IgG. Ranking of these sections is based on the extent of proliferation on a scale of 0-8, the lower side of the scale reflecting slight proliferation to the higher side reflecting intense proliferation.

Experimental data are analyzed using an unpaired t test. A p value of < 0.05 is considered significant.

B. Steroid Impaired Rat Model

The inhibition of wound healing by steroids has been well documented in various *in vitro* and *in vivo* systems (Wahl, Glucocorticoids and Wound healing. In: Anti-Inflammatory Steroid Action: Basic and Clinical Aspects. 280-302 (1989); Wahlet *et al.*, *J. Immunol.* 115: 476-481 (1975); Werb *et al.*, *J. Exp. Med.* 147:1684-1694 (1978)). Glucocorticoids retard wound healing by inhibiting angiogenesis, decreasing vascular permeability (Ebert *et al.*, *An. Intern. Med.* 37:701-705 (1952)), fibroblast proliferation, and collagen synthesis (Beck *et al.*, *Growth Factors.* 5: 295-304 (1991); Haynes *et al.*, *J. Clin. Invest.* 61: 703-797 (1978)) and producing a transient reduction of circulating monocytes (Haynes *et al.*, *J. Clin. Invest.* 61: 703-797 (1978); Wahl, "Glucocorticoids and wound healing", In: Antiinflammatory Steroid Action: Basic and Clinical Aspects, Academic Press, New York, pp. 280-302 (1989)). The systemic administration of steroids to impaired wound healing is a well establish phenomenon in rats (Beck *et al.*, *Growth Factors.* 5: 295-304 (1991); Haynes *et al.*, *J. Clin. Invest.* 61: 703-797 (1978); Wahl, "Glucocorticoids and wound healing", In: Antiinflammatory Steroid Action: Basic and Clinical Aspects, Academic Press, New York, pp. 280-302 (1989); Pierce *et al.*, *Proc. Natl. Acad. Sci. USA* 86: 2229-2233 (1989)).

To demonstrate that a polypeptide of the invention can accelerate the healing process, the effects of multiple topical applications of the polypeptide on full thickness excisional skin wounds in rats in which healing has been impaired by the systemic administration of methylprednisolone is assessed.

Young adult male Sprague Dawley rats weighing 250-300 g (Charles River Laboratories) are used in this example. The animals are purchased at 8 weeks of age and are 9 weeks old at the beginning of the study. The healing response of rats is impaired by the systemic administration of methylprednisolone (17mg/kg/rat intramuscularly) at the time of wounding. Animals are individually housed and received food and water *ad libitum*. All manipulations are performed using aseptic techniques. This study is conducted according to the rules and guidelines of Human Genome Sciences, Inc. Institutional Animal Care and Use Committee and the Guidelines for the Care and Use of Laboratory Animals.

The wounding protocol is followed according to section A, above. On the day of wounding, animals are anesthetized with an intramuscular injection of ketamine (50 mg/kg) and xylazine (5 mg/kg). The dorsal region of the animal is shaved and the skin washed with 70% ethanol and iodine solutions. The surgical area is dried with sterile gauze prior to wounding. An 8 mm full-thickness wound is created using a Keyes tissue punch. The wounds are left open for the duration of the experiment. Applications of the testing materials are given topically once a day for 7 consecutive days commencing on the day of wounding and subsequent to methylprednisolone administration. Prior to treatment, wounds are gently cleansed with sterile saline and gauze sponges.

Wounds are visually examined and photographed at a fixed distance at the day of wounding and at the end of treatment. Wound closure is determined by daily measurement on days 1-5 and on day 8. Wounds are measured horizontally and vertically using a calibrated Jameson caliper. Wounds are considered healed if granulation tissue is no longer visible and the wound is covered by a continuous epithelium.

The polypeptide of the invention is administered using at a range different doses, from 4mg to 500mg per wound per day for 8 days in vehicle. Vehicle control groups received 50mL of vehicle solution.

Animals are euthanized on day 8 with an intraperitoneal injection of sodium pentobarbital (300mg/kg). The wounds and surrounding skin are then harvested for histology. Tissue specimens are placed in 10% neutral buffered formalin in tissue cassettes between biopsy sponges for further processing.

Four groups of 10 animals each (5 with methylprednisolone and 5 without glucocorticoid) are evaluated: 1) Untreated group 2) Vehicle placebo control 3) treated groups.

Wound closure is analyzed by measuring the area in the vertical and horizontal axis and obtaining the total area of the wound. Closure is then estimated by establishing the differences between the initial wound area (day 0) and that of post treatment (day 8). The wound area on day 1 is 64mm², the corresponding size of the dermal punch. Calculations are made using the following formula:

$$[\text{Open area on day 8}] - [\text{Open area on day 1}] / [\text{Open area on day 1}]$$

Specimens are fixed in 10% buffered formalin and paraffin embedded blocks are sectioned perpendicular to the wound surface (5mm) and cut using an Olympus microtome. Routine hematoxylin-eosin (H&E) staining is performed on cross-sections of bisected wounds. Histologic examination of the wounds allows assessment of whether the healing process and the morphologic appearance of the repaired skin is improved by treatment with a polypeptide of the invention. A calibrated lens micrometer is used by a blinded observer to determine the distance of the wound gap.

Experimental data are analyzed using an unpaired t test. A p value of < 0.05 is considered significant.

The studies described in this example tested activity of a polypeptide of the invention. However, one skilled in the art could easily modify the exemplified studies to test the activity of polynucleotides (e.g., gene therapy), agonists, and/or antagonists of the invention.

Example 51: Lymphadema Animal Model

The purpose of this experimental approach is to create an appropriate and consistent lymphedema model for testing the therapeutic effects of a polypeptide of the invention in lymphangiogenesis and re-establishment of the lymphatic circulatory system in the rat hind limb. Effectiveness is measured by swelling volume of the affected limb, quantification of the amount of lymphatic vasculature, total blood plasma protein, and histopathology. Acute lymphedema is observed for 7-10 days. Perhaps more importantly, the chronic progress of the edema is followed for up to 3-4 weeks.

Prior to beginning surgery, blood sample is drawn for protein concentration analysis. Male rats weighing approximately ~350g are dosed with Pentobarbital. Subsequently, the right legs are shaved from knee to hip. The shaved area is swabbed with gauze soaked in 70% EtOH. Blood is drawn for serum total protein testing. Circumference and volumetric measurements are made prior to injecting dye into paws after marking 2 measurement levels (0.5 cm above heel, at mid-pt of dorsal paw). The intradermal dorsum of both right and left paws are injected with 0.05 ml of 1% Evan's

Blue. Circumference and volumetric measurements are then made following injection of dye into paws.

Using the knee joint as a landmark, a mid-leg inguinal incision is made circumferentially allowing the femoral vessels to be located. Forceps and hemostats are
5 used to dissect and separate the skin flaps. After locating the femoral vessels, the lymphatic vessel that runs along side and underneath the vessel(s) is located. The main lymphatic vessels in this area are then electrically coagulated suture ligated.

Using a microscope, muscles in back of the leg (near the semitendinosus and adductors) are bluntly dissected. The popliteal lymph node is then located. The 2
10 proximal and 2 distal lymphatic vessels and distal blood supply of the popliteal node are then and ligated by suturing. The popliteal lymph node, and any accompanying adipose tissue, is then removed by cutting connective tissues.

Care is taken to control any mild bleeding resulting from this procedure. After lymphatics are occluded, the skin flaps are sealed by using liquid skin (Vetbond) (AJ
15 Buck). The separated skin edges are sealed to the underlying muscle tissue while leaving a gap of ~0.5 cm around the leg. Skin also may be anchored by suturing to underlying muscle when necessary.

To avoid infection, animals are housed individually with mesh (no bedding). Recovering animals are checked daily through the optimal edematous peak, which
20 typically occurred by day 5-7. The plateau edematous peak are then observed. To evaluate the intensity of the lymphedema, the circumference and volumes of 2 designated places on each paw before operation and daily for 7 days are measured. The effect plasma proteins on lymphedema is determined and whether protein analysis is a useful testing perimeter is also investigated. The weights of both control and edematous limbs are
25 evaluated at 2 places. Analysis is performed in a blind manner.

Circumference Measurements: Under brief gas anesthetic to prevent limb movement, a cloth tape is used to measure limb circumference. Measurements are done at the ankle bone and dorsal paw by 2 different people then those 2 readings are averaged. Readings are taken from both control and edematous limbs.

30 Volumetric Measurements: On the day of surgery, animals are anesthetized with Pentobarbital and are tested prior to surgery. For daily volumetrics animals are under

brief halothane anesthetic (rapid immobilization and quick recovery), both legs are shaved and equally marked using waterproof marker on legs. Legs are first dipped in water, then dipped into instrument to each marked level then measured by Buxco edema software(Chen/Victor). Data is recorded by one person, while the other is dipping the
5 limb to marked area.

Blood-plasma protein measurements: Blood is drawn, spun, and serum separated prior to surgery and then at conclusion for total protein and Ca²⁺ comparison.

Limb Weight Comparison: After drawing blood, the animal is prepared for tissue collection. The limbs are amputated using a quillitine, then both experimental and control
10 legs are cut at the ligature and weighed. A second weighing is done as the tibio-cacaneal joint is disarticulated and the foot is weighed.

Histological Preparations: The transverse muscle located behind the knee (popliteal) area is dissected and arranged in a metal mold, filled with freezeGel, dipped into cold methylbutane, placed into labeled sample bags at - 80EC until sectioning. Upon
15 sectioning, the muscle is observed under fluorescent microscopy for lymphatics..

The studies described in this example tested activity of a polypeptide of the invention. However, one skilled in the art could easily modify the exemplified studies to test the activity of polynucleotides (e.g., gene therapy), agonists, and/or antagonists of the invention.
20

Example 52: Suppression of TNF alpha-induced adhesion molecule expression by a Polypeptide of the Invention

The recruitment of lymphocytes to areas of inflammation and angiogenesis involves specific receptor-ligand interactions between cell surface adhesion molecules
25 (CAMs) on lymphocytes and the vascular endothelium. The adhesion process, in both normal and pathological settings, follows a multi-step cascade that involves intercellular adhesion molecule-1 (ICAM-1), vascular cell adhesion molecule-1 (VCAM-1), and endothelial leukocyte adhesion molecule-1 (E-selectin) expression on endothelial cells (EC). The expression of these molecules and others on the vascular endothelium
30 determines the efficiency with which leukocytes may adhere to the local vasculature and extravasate into the local tissue during the development of an inflammatory response. The

local concentration of cytokines and growth factor participate in the modulation of the expression of these CAMs.

Tumor necrosis factor alpha (TNF- α), a potent proinflammatory cytokine, is a stimulator of all three CAMs on endothelial cells and may be involved in a wide variety of inflammatory responses, often resulting in a pathological outcome.

The potential of a polypeptide of the invention to mediate a suppression of TNF- α induced CAM expression can be examined. A modified ELISA assay which uses ECs as a solid phase absorbent is employed to measure the amount of CAM expression on TNF- α treated ECs when co-stimulated with a member of the FGF family of proteins.

To perform the experiment, human umbilical vein endothelial cell (HUVEC) cultures are obtained from pooled cord harvests and maintained in growth medium (EGM-2; Clonetics, San Diego, CA) supplemented with 10% FCS and 1% penicillin/streptomycin in a 37 degree C humidified incubator containing 5% CO₂. HUVECs are seeded in 96-well plates at concentrations of 1×10^4 cells/well in EGM medium at 37 degree C for 18-24 hrs or until confluent. The monolayers are subsequently washed 3 times with a serum-free solution of RPMI-1640 supplemented with 100 U/ml penicillin and 100 mg/ml streptomycin, and treated with a given cytokine and/or growth factor(s) for 24 h at 37 degree C. Following incubation, the cells are then evaluated for CAM expression.

Human Umbilical Vein Endothelial cells (HUVECs) are grown in a standard 96 well plate to confluence. Growth medium is removed from the cells and replaced with 90 μ l of 199 Medium (10% FBS). Samples for testing and positive or negative controls are added to the plate in triplicate (in 10 μ l volumes). Plates are incubated at 37 degree C for either 5 h (selectin and integrin expression) or 24 h (integrin expression only). Plates are aspirated to remove medium and 100 μ l of 0.1% paraformaldehyde-PBS(with Ca⁺⁺ and Mg⁺⁺) is added to each well. Plates are held at 4°C for 30 min.

Fixative is then removed from the wells and wells are washed 1X with PBS(+Ca,Mg)+0.5% BSA and drained. Do not allow the wells to dry. Add 10 μ l of diluted primary antibody to the test and control wells. Anti-ICAM-1-Biotin, Anti-VCAM-1-Biotin and Anti-E-selectin-Biotin are used at a concentration of 10 μ g/ml (1:10 dilution

of 0.1 mg/ml stock antibody). Cells are incubated at 37°C for 30 min. in a humidified environment. Wells are washed X3 with PBS(+Ca,Mg)+0.5% BSA.

Then add 20 µl of diluted ExtrAvidin-Alkaline Phosphotase (1:5,000 dilution) to each well and incubated at 37°C for 30 min. Wells are washed X3 with

5 PBS(+Ca,Mg)+0.5% BSA. 1 tablet of p-Nitrophenol Phosphate pNPP is dissolved in 5 ml of glycine buffer (pH 10.4). 100 µl of pNPP substrate in glycine buffer is added to each test well. Standard wells in triplicate are prepared from the working dilution of the ExtrAvidin-Alkaline Phosphotase in glycine buffer: 1:5,000 (10^0) > $10^{-0.5}$ > 10^{-1} > $10^{-1.5}$. 5 µl of each dilution is added to triplicate wells and the resulting AP content in each well is

10 5.50 ng, 1.74 ng, 0.55 ng, 0.18 ng. 100 µl of pNPP reagent must then be added to each of the standard wells. The plate must be incubated at 37°C for 4h. A volume of 50 µl of 3M NaOH is added to all wells. The results are quantified on a plate reader at 405 nm. The background subtraction option is used on blank wells filled with glycine buffer only. The template is set up to indicate the concentration of AP-conjugate in each standard well [

15 5.50 ng; 1.74 ng; 0.55 ng; 0.18 ng]. Results are indicated as amount of bound AP-conjugate in each sample.

The studies described in this example tested activity of a polypeptide of the invention. However, one skilled in the art could easily modify the exemplified studies to test the activity of polynucleotides (e.g., gene therapy), agonists, and/or antagonists of the

20 invention.

It will be clear that the invention may be practiced otherwise than as particularly described in the foregoing description and examples. Numerous modifications and variations of the present invention are possible in light of the above teachings and, therefore, are within the scope of the appended claims.


25 The entire disclosure of each document cited (including patents, patent applications, journal articles, abstracts, laboratory manuals, books, or other disclosures) in the Background of the Invention, Detailed Description, and Examples is hereby incorporated herein by reference. Further, the hard copy of the sequence listing submitted herewith and the corresponding computer readable form are both

30 incorporated herein by reference in their entireties.

INDICATIONS RELATING TO A DEPOSITED MICROORGANISM

(PCT Rule 13bis)

A. The indications made below relate to the microorganism referred to in the description on page <u>71</u> , line <u>N/A</u>	
B. IDENTIFICATION OF DEPOSIT Further deposits are identified on an additional sheet <input type="checkbox"/>	
Name of depositary institution <u>American Type Culture Collection</u>	
Address of depositary institution <i>(including postal code and country)</i> <u>10801 University Boulevard</u> <u>Manassas, Virginia 20110-2209</u> <u>United States of America</u>	
Date of deposit <u>08 April 1999</u>	Accession Number <u>203917</u>
C. ADDITIONAL INDICATIONS <i>(leave blank if not applicable)</i> This information is continued on an additional sheet <input type="checkbox"/>	
D. DESIGNATED STATES FOR WHICH INDICATIONS ARE MADE <i>(if the indications are not for all designated States)</i>	
<u>Europe</u> In respect to those designations in which a European Patent is sought a sample of the deposited microorganism will be made available until the publication of the mention of the grant of the European patent or until the date on which application has been refused or withdrawn or is deemed to be withdrawn, only by the issue of such a sample to an expert nominated by the person requesting the sample (Rule 28 (4) EPC).	
E. SEPARATE FURNISHING OF INDICATIONS <i>(leave blank if not applicable)</i>	
The indications listed below will be submitted to the International Bureau later <i>(specify the general nature of the indications e.g., "Accession Number of Deposit")</i>	

For receiving Office use only <input checked="" type="checkbox"/> This sheet was received with the international application Authorized officer 	For International Bureau use only <input type="checkbox"/> This sheet was received by the International Bureau on: Authorized officer
---	--

ATCC Deposit No.: 203917

CANADA

The applicant requests that, until either a Canadian patent has been issued on the basis of an application or the application has been refused, or is abandoned and no longer subject to reinstatement, or is withdrawn, the Commissioner of Patents only authorizes the furnishing of a sample of the deposited biological material referred to in the application to an independent expert nominated by the Commissioner, the applicant must, by a written statement, inform the International Bureau accordingly before completion of technical preparations for publication of the international application.

NORWAY

The applicant hereby requests that the application has been laid open to public inspection (by the Norwegian Patent Office), or has been finally decided upon by the Norwegian Patent Office without having been laid open inspection, the furnishing of a sample shall only be effected to an expert in the art. The request to this effect shall be filed by the applicant with the Norwegian Patent Office not later than at the time when the application is made available to the public under Sections 22 and 33(3) of the Norwegian Patents Act. If such a request has been filed by the applicant, any request made by a third party for the furnishing of a sample shall indicate the expert to be used. That expert may be any person entered on the list of recognized experts drawn up by the Norwegian Patent Office or any person approved by the applicant in the individual case.

AUSTRALIA

The applicant hereby gives notice that the furnishing of a sample of a microorganism shall only be effected prior to the grant of a patent, or prior to the lapsing, refusal or withdrawal of the application, to a person who is a skilled addressee without an interest in the invention (Regulation 3.25(3) of the Australian Patents Regulations).

FINLAND

The applicant hereby requests that, until the application has been laid open to public inspection (by the National Board of Patents and Regulations), or has been finally decided upon by the National Board of Patents and Registration without having been laid open to public inspection, the furnishing of a sample shall only be effected to an expert in the art.

UNITED KINGDOM

The applicant hereby requests that the furnishing of a sample of a microorganism shall only be made available to an expert. The request to this effect must be filed by the applicant with the International Bureau before the completion of the technical preparations for the international publication of the application.

ATCC Deposit No.: 203917

DENMARK

The applicant hereby requests that, until the application has been laid open to public inspection (by the Danish Patent Office), or has been finally decided upon by the Danish Patent office without having been laid open to public inspection, the furnishing of a sample shall only be effected to an expert in the art. The request to this effect shall be filed by the applicant with the Danish Patent Office not later than at the time when the application is made available to the public under Sections 22 and 33(3) of the Danish Patents Act. If such a request has been filed by the applicant, any request made by a third party for the furnishing of a sample shall indicate the expert to be used. That expert may be any person entered on a list of recognized experts drawn up by the Danish Patent Office or any person by the applicant in the individual case.

SWEDEN

The applicant hereby requests that, until the application has been laid open to public inspection (by the Swedish Patent Office), or has been finally decided upon by the Swedish Patent Office without having been laid open to public inspection, the furnishing of a sample shall only be effected to an expert in the art. The request to this effect shall be filed by the applicant with the International Bureau before the expiration of 16 months from the priority date (preferably on the Form PCT/RO/134 reproduced in annex Z of Volume I of the PCT Applicant's Guide). If such a request has been filed by the applicant any request made by a third party for the furnishing of a sample shall indicate the expert to be used. That expert may be any person entered on a list of recognized experts drawn up by the Swedish Patent Office or any person approved by a applicant in the individual case.

NETHERLANDS

The applicant hereby requests that until the date of a grant of a Netherlands patent or until the date on which the application is refused or withdrawn or lapsed, the microorganism shall be made available as provided in the 31F(1) of the Patent Rules only by the issue of a sample to an expert. The request to this effect must be furnished by the applicant with the Netherlands Industrial Property Office before the date on which the application is made available to the public under Section 22C or Section 25 of the Patents Act of the Kingdom of the Netherlands, whichever of the two dates occurs earlier.

What Is Claimed Is:

1. An isolated nucleic acid molecule comprising a polynucleotide having a nucleotide sequence at least 95% identical to a sequence selected from the group
5 consisting of:
 - (a) a polynucleotide fragment of SEQ ID NO:X or a polynucleotide fragment of the cDNA sequence included in ATCC Deposit No:Z, which is hybridizable to SEQ ID NO:X;
 - (b) a polynucleotide encoding a polypeptide fragment of SEQ ID NO:Y or a
10 polypeptide fragment encoded by the cDNA sequence included in ATCC Deposit No:Z, which is hybridizable to SEQ ID NO:X;
 - (c) a polynucleotide encoding a polypeptide domain of SEQ ID NO:Y or a polypeptide domain encoded by the cDNA sequence included in ATCC Deposit No:Z, which is hybridizable to SEQ ID NO:X;
 - 15 (d) a polynucleotide encoding a polypeptide epitope of SEQ ID NO:Y or a polypeptide epitope encoded by the cDNA sequence included in ATCC Deposit No:Z, which is hybridizable to SEQ ID NO:X;
 - (e) a polynucleotide encoding a polypeptide of SEQ ID NO:Y or the cDNA sequence included in ATCC Deposit No:Z, which is hybridizable to SEQ ID NO:X,
20 having biological activity;
 - (f) a polynucleotide which is a variant of SEQ ID NO:X;
 - (g) a polynucleotide which is an allelic variant of SEQ ID NO:X;
 - (h) a polynucleotide which encodes a species homologue of the SEQ ID NO:Y;
 - 25 (i) a polynucleotide capable of hybridizing under stringent conditions to any one of the polynucleotides specified in (a)-(h), wherein said polynucleotide does not hybridize under stringent conditions to a nucleic acid molecule having a nucleotide sequence of only A residues or of only T residues.

2. The isolated nucleic acid molecule of claim 1, wherein the polynucleotide fragment comprises a nucleotide sequence encoding a secreted protein.

5 3. The isolated nucleic acid molecule of claim 1, wherein the polynucleotide fragment comprises a nucleotide sequence encoding the sequence identified as SEQ ID NO:Y or the polypeptide encoded by the cDNA sequence included in ATCC Deposit No:Z, which is hybridizable to SEQ ID NO:X.

10 4. The isolated nucleic acid molecule of claim 1, wherein the polynucleotide fragment comprises the entire nucleotide sequence of SEQ ID NO:X or the cDNA sequence included in ATCC Deposit No:Z, which is hybridizable to SEQ ID NO:X.

15 5. The isolated nucleic acid molecule of claim 2, wherein the nucleotide sequence comprises sequential nucleotide deletions from either the C-terminus or the N-terminus.

20 6. The isolated nucleic acid molecule of claim 3, wherein the nucleotide sequence comprises sequential nucleotide deletions from either the C-terminus or the N-terminus.

7. A recombinant vector comprising the isolated nucleic acid molecule of claim 1.

25

8. A method of making a recombinant host cell comprising the isolated nucleic acid molecule of claim 1.

9. A recombinant host cell produced by the method of claim 8.

30

10. The recombinant host cell of claim 9 comprising vector sequences.

11. An isolated polypeptide comprising an amino acid sequence at least 95% identical to a sequence selected from the group consisting of:

- 5 (a) a polypeptide fragment of SEQ ID NO:Y or the encoded sequence included in ATCC Deposit No:Z;
- (b) a polypeptide fragment of SEQ ID NO:Y or the encoded sequence included in ATCC Deposit No:Z, having biological activity;
- (c) a polypeptide domain of SEQ ID NO:Y or the encoded sequence included in ATCC Deposit No:Z;
- 10 (d) a polypeptide epitope of SEQ ID NO:Y or the encoded sequence included in ATCC Deposit No:Z;
- (e) a secreted form of SEQ ID NO:Y or the encoded sequence included in ATCC Deposit No:Z;
- (f) a full length protein of SEQ ID NO:Y or the encoded sequence included in
15 ATCC Deposit No:Z;
- (g) a variant of SEQ ID NO:Y;
- (h) an allelic variant of SEQ ID NO:Y; or
- (i) a species homologue of the SEQ ID NO:Y.

20 12. The isolated polypeptide of claim 11, wherein the secreted form or the full length protein comprises sequential amino acid deletions from either the C-terminus or the N-terminus.

25 13. An isolated antibody that binds specifically to the isolated polypeptide of claim 11.

14. A recombinant host cell that expresses the isolated polypeptide of claim 11.

30 15. A method of making an isolated polypeptide comprising:
(a) culturing the recombinant host cell of claim 14 under conditions such that said polypeptide is expressed; and

(b) recovering said polypeptide.

16. The polypeptide produced by claim 15.

5 17. A method for preventing, treating, or ameliorating a medical condition, comprising administering to a mammalian subject a therapeutically effective amount of the polypeptide of claim 11 or the polynucleotide of claim 1.

10 18. A method of diagnosing a pathological condition or a susceptibility to a pathological condition in a subject comprising:

(a) determining the presence or absence of a mutation in the polynucleotide of claim 1; and

(b) diagnosing a pathological condition or a susceptibility to a pathological condition based on the presence or absence of said mutation.

15

19. A method of diagnosing a pathological condition or a susceptibility to a pathological condition in a subject comprising:

(a) determining the presence or amount of expression of the polypeptide of claim 11 in a biological sample; and

20 (b) diagnosing a pathological condition or a susceptibility to a pathological condition based on the presence or amount of expression of the polypeptide.

20. A method for identifying a binding partner to the polypeptide of claim 11 comprising:

25 (a) contacting the polypeptide of claim 11 with a binding partner; and

(b) determining whether the binding partner effects an activity of the polypeptide.

30 21. The gene corresponding to the cDNA sequence of SEQ ID NO:Y.

22. A method of identifying an activity in a biological assay, wherein the method comprises:

- (a) expressing SEQ ID NO:X in a cell;
- (b) isolating the supernatant;
- 5 (c) detecting an activity in a biological assay; and
- (d) identifying the protein in the supernatant having the activity.

23. The product produced by the method of claim 20.

<110> Human Genome Sciences, Inc.

<120> 47 Human Secreted Proteins

<130> PS507PCT

<140> Unassigned

<141> 2000-03-09

<150> 60/124,095

<151> 1999-03-12

<150> 60/138,598

<151> 0999-06-11

<150> 60/168,665

<151> 1999-12-03

<160> 148

<170> PatentIn Ver. 2.0

<210> 1

<211> 733

<212> DNA

<213> Homo sapiens

<400> 1

gggatccgga	gccc aaatct	tctgacaaaa	ctcacacatg	cccaccggtg	ccagcacctg	60
aattcgaggg	tgcaccgtca	gtcttctctt	tcccccaaaa	acccaaggac	accctcatga	120
tctcccgga	tcttgaggtc	acatgcgtgg	tgggtggacgt	aagccacga	gaccctgagg	180
tcaagttcaa	ctggtacgtg	gacggcggtg	aggtgcataa	tgccaagaca	aagccgcggg	240
aggagcagta	caacagcacg	taccgtgtgg	tcagcgtcct	caccgtcctg	caccaggact	300
ggctgaatgg	caaggagtac	aagtgcgaag	tctccaacaa	agccctccca	acccccatcg	360
agaaaaccat	ctccaaagcc	aaagggcagc	cccgagaacc	acaggtgtac	accctgcccc	420
catcccggga	tgagctgacc	aagaaccagg	tcagcctgac	ctgcctggtc	aaaggcttct	480
atccaagcga	catcgccgtg	gagtgggaga	gcaatgggca	gccgggagaac	aactacaaga	540
ccacgcctcc	cgtgctggac	tccgacggct	ccttcttctt	ctacagcaag	ctcaccgtgg	600
acaagagcag	gtggcagcag	gggaacgtct	tctcatgtct	cgtgatgcat	gaggctctgc	660
acaaccacta	cacgcagaag	agcctctccc	tgtctccggg	taaatgagtg	cgacggccgc	720
gactctagag	gat					733

<210> 2

<211> 5

<212> PRT

<213> Homo sapiens

<220>

<221> Site

<222> (3)

<223> Xaa equals any of the twenty naturally occurring L-amino acids

<400> 2

Trp Ser Xaa Trp Ser
1 5

<210> 3

<211> 86

<212> DNA

<213> Homo sapiens

<400> 3

gcgcctcgag atttccccga aatctagatt tccccgaaat gatttccccg aaatgatttc 60
cccgaatat ctgccatctc aattag 86

<210> 4

<211> 27

<212> DNA

<213> Homo sapiens

<400> 4

gcggcaagct ttttgcaaag cctaggc 27

<210> 5

<211> 271

<212> DNA

<213> Homo sapiens

<400> 5

ctcgagattt ccccgaaatc tagatttccc cgaaatgatt tccccgaaat gatttccccg 60
aaatatctgc catctcaatt agtcagcaac catagtcccc cccctaactc cgcccatccc 120
gccctaact ccgcccagtt ccgcccattc tccgccccat ggctgactaa ttttttttat 180
ttatgcagag gccgaggccg cctcggcctc tgagctattc cagaagtagt gaggaggctt 240
ttttggaggc ctaggctttt gcaaaaagct t 271

<210> 6

<211> 32

<212> DNA

<213> Homo sapiens

<400> 6

gcgctcgagg gatgacagcg atagaacccc gg 32

<210> 7

<211> 31

<212> DNA

<213> Homo sapiens

<400> 7

gcgaagcttc gcgactcccc ggatccgcct c 31

<210> 8

<211> 12
 <212> DNA
 <213> Homo sapiens

<400> 8
 ggggactttc cc

12

<210> 9
 <211> 73
 <212> DNA
 <213> Homo sapiens

<400> 9
 gcggcctcga ggggactttc ccggggactt tccggggact ttccgggact ttccatcctg
 ccattctcaat tag

60

73

<210> 10
 <211> 256
 <212> DNA
 <213> Homo sapiens

<400> 10
 ctccgagggga ctttcccggg gactttccgg ggactttccg ggactttcca tctgccatct 60
 caattagtca gcaaccatag tcccggccct aactccgccc atcccggccc taactccgcc 120
 cagttccgcc cattctccgc cccatggctg actaattttt tttatttatg cagaggccga 180
 ggccgcctcg gcctctgagc tattccagaa gtagtgagga ggcttttttg gaggcctagg 240
 cttttgcaaa aagctt 256

<210> 11
 <211> 1229
 <212> DNA
 <213> Homo sapiens

<400> 11
 ggcacgagaa aatacaaaaa ttatccaggc gaggtggtgc acgtgtgtag tcccagttac 60
 tcgggaggct gaggtgggag gatggcttga atctgggagg tgaagattgc agtgagctga 120
 gatcacacca ctgcattcag tctggataat agaacaagac tctatctcaa aaaaaaatt 180
 gtatacttta ttgactcatt tatgtctgat gggatatttt gattacaaat tgtttagtca 240
 ctactttaag gcctgcttta tttttttctt aattttattg actcatttat gtcttagtct 300
 tttttattac aaattattta ttgttcagtc actaccttag agcctgtttt attttttctt 360
 aattttatta aaggatgata ttgatgatga aatgtcttac gatgatcatt tagaggttta 420
 ttttgaacaa ctggcaattc cacgaatgat ggaataaaac atatgaagta gaaggactgg 480
 aacctccaga aaaagtactt taagttacct acagggtgat ctagtcagggt atgaattgat 540
 aagaaatgcc tgcaccttcc ctcttctcta tctttccctt gcctacagaa aattaaagg 600
 caaaacaatg gacatctaca tattcttcat tcagatcaac cagtggctag catttgccac 660
 cttttgcagt ttctttctct ttccataagt actttcttct ctgaatcatt tgaaagcaaa 720
 tgaaaacagt agcctaaaagt gtcagtttca accagaaaaa aacagctctg atttctcatg 780
 gctcacactc ttctgaaacg actcgggtag aggctgagga aggccgtgtt gtttgtctac 840
 ctgggactag taagtataga aatagaattc ctttgttctt aaattctacc tttgacttta 900
 cttttaaaat ataatttctt tggtagcatt tagctcatgc ctgtaatcct agcatttttg 960
 gaggccaaag agggagaatt gcttgagccc aggagtttga gaccagcctg gacatcatag 1020
 ggagactctg cgcacgcaca cacacacaca cagcgcacg ctaaccggga atgggtggcat 1080

gcgactgtgg	ccccagctgc	ctgggaggct	gaggttggag	gatcatttgg	accagaggag	1140
tggagactgc	agtgagccat	gattgcacca	ctgcactcca	gcctgggtga	cagagcaaga	1200
ccctgtttca	caaaaaaaaa	aaaaaaaaa				1229

<210> 12
 <211> 1538
 <212> DNA
 <213> Homo sapiens

<220>
 <221> SITE
 <222> (724)
 <223> n equals a,t,g, or c

<220>
 <221> SITE
 <222> (889)
 <223> n equals a,t,g, or c

<220>
 <221> SITE
 <222> (975)
 <223> n equals a,t,g, or c

<220>
 <221> SITE
 <222> (1008)
 <223> n equals a,t,g, or c

<400> 12						
ggcacgagca	aataatcatc	tgttgaaaaa	tatgaagaaa	attgtggatc	aaaacacaaa	60
gcttgctcct	gagaccaagg	ctgtcattca	ttggattatg	gatattcctt	ttgtgctttc	120
tgccaatctc	catggaggag	accttgtggc	caattatcca	tatgatgaga	cgcggagtgg	180
tagtgctcac	gaatacagct	cctccccaga	tgacgccatt	ttccaaagct	tggcccgggc	240
atactcttct	ttcaaccggg	ccatgtctga	ccccaatcgg	ccaccatgtc	gcaagaatga	300
tgatgacagc	agctttgtag	atggaaccac	caacgggtgtg	cttggtacag	cgtacctgga	360
gggatgcaag	acttcaatta	ccttagcagc	aactgttttg	agatcacctg	ggagcttagc	420
tgtgagaagt	tcccacctga	agagactctg	aagacctact	gggaggataa	caaaaactcc	480
ctcattagct	accttgagca	gatacaccga	ggagttaaag	gatttgtccg	agaccttcaa	540
ggtaacccaa	ttgcgatggc	caccaatctc	cgtggaagga	atagaccacg	atgttacatc	600
cgcaaaggat	ggcgattact	ggagattgct	tatacctgga	aactataaac	ttacactcag	660
ctccaggcta	tctggcaata	acaaagaaag	tggcagttcc	ttacagccct	gctgctgggg	720
gttnattttg	aactgggagt	cattttctga	aaggaaagaa	gaggagaagg	aagaattgat	780
ggaatgggtg	aaaatgatgt	cagaaaacttt	aaatttttaa	aaaggcttct	agtttagctgc	840
tttaaatcta	tctatataat	gtagtatgat	gtaatgtggt	cttttttttna	gattttgtgc	900
agttaatact	taacattgat	ttatttttta	atcattttaa	tattaatcaa	ctttccttaa	960
aataaatagc	ctctnaggta	aaaatataag	aacttgatat	atttcattnc	cttatatagt	1020
attcattttc	ctacctatat	ttccccaaaa	agtttaggaa	aggtttaaga	attttttgcca	1080
tccttaggct	taaatgcaat	attcctggta	ttatttacaa	tgcagaattt	tttgagtaat	1140
tctagctttc	aaaaattagt	gaagttcttt	tactgtaatt	ggtgacaatg	tcacataatg	1200
aatgctattg	aaaagggttaa	cagatacagc	tcggagtgtg	gagcactcta	ctgcaagact	1260
taaatagttc	agtataaatt	gtcgtttttt	tcttgtgctg	actaactata	agcatgatct	1320
tgtaaatgca	tttttgatgg	gaagaaaagg	tacatgttta	caaagaggtt	ttatgaaaag	1380
aataaaaaatt	gacttcttgc	ttgtacatat	aggagcaata	ctattatatt	atgtagtccc	1440

gttaacacta cttaaaagtt taggggttttc tcttggttgt agagtggccc agaattgcat 1500
tctgaatgaa taaagggtta aaaaaaaaaa aaaaaaaaaa 1538

<210> 13
<211> 1032
<212> DNA
<213> Homo sapiens

<400> 13
ggcacgaggt taattaactg ctctgtgctg tagtttctcc atctgtaaaa ctgtcaaggt 60
agtagtacat gtcattagga ttaagtaa atagctatgt aaaacacaga taacattgga 120
tgacaccaat ggtaggcag cagatacatt tattagtaaa gtgcaatgga aattttcctc 180
ttgctggata agagtaaact aatgattaga cttgaggatg gcttaactaa gaaagaaagt 240
gatcaagaag gtaagagtag taatgattgg ttttgagaat ttaattttct gtgcttatca 300
tggaataaat tttagctgaat taaaatgcat tataattaca agagttagga tatcttaaaa 360
tatctagtga ggctgtataa cctagcaaga tcaaacaaac tcattgctat tggattttaa 420
gaaattggat tgtcaagaag ataagtaatc aatatggctt ggctgtgtcc ccacccaaat 480
ctcatcttga attatagctc tcattattcc cacatattgt gggagtggacc tgggtgggaga 540
caactgaatc atgggggcag ttttcccat actgttctcg tggtagtgag taactctcac 600
gaaatctgat ggttttataa ggggaactcc cttttgcttg gctttcattc tgtcttgctc 660
gtgccatgt aagatgtgcc tttcgcccta tgccattatt ctgaggcctc ccagccacg 720
tggaattgtg agtcccttaa acctcttttt gtttataaat caccagctct tgggtacatc 780
tttatcagca gtgtgaaaat ggactaatac agtaatcttg caagcctgca caaacttgcg 840
ctttgtataa aaagccacaa taataaaact tgttgagaat atcctaccaa aattactatg 900
aaagttatct ctccctttta ctatgagcag aggtggcca ttagaataat atgattgtct 960
ctggattatg tataataaga aaatgtgtgt atacaaataa aatgcattta ttatgaagta 1020
aaaaaaaaaa aa 1032

<210> 14
<211> 1896
<212> DNA
<213> Homo sapiens

<220>
<221> SITE
<222> (933)
<223> n equals a,t,g, or c

<220>
<221> SITE
<222> (940)
<223> n equals a,t,g, or c

<400> 14
ggcacaggac cagtggagtg agctgttcat ggatgcgcta gggcccttca acttcgtgct 60
ggtgagttcg gtgaggatgc aggtgtcat cctgctgctg ttccccaagt actaccacct 120
gcccttcctg cgagacgtgc agaccgactg cacgcgcact ggctggggcg gctactgggg 180
taacaagggt ggcgtgagcg tgcgcctggc ggcttcggg cacatgctct gcttcctgaa 240
ctgccacttg cctgcgcata tggacaaggc ggagcagcgc aaagacaact tccagaccat 300
cctcagcctc cagcagttcc aaggggccggg cgcacagggc atcctggatc atgacctcgt 360
gttctgggtc ggggacctga acttcgcgat tgagagctat gacctgcact ttgtcaagtt 420
tgccatcgac agtgaccage tccatcagct ctgggagaag gaccagctca acatggccaa 480
gaacacctgg ccattctga agggctttca ggagggggccc ctcaacttcg ctcccacctt 540

caagtttgat	gtgggtacca	acaaatacga	taccagtgcc	aagaaacgga	agccagcttg	600
gacagaccgt	atcctatgga	aggtcaaggc	tccaggtggg	ggtcccagcc	cctcaggacg	660
gaagagccac	cgactccagg	tgacgcagca	cagctaccgc	agccacatgg	aatacacagt	720
cagcgaccac	aagcctgtgg	ytgccagtt	cctcctgcag	tttgcctttc	agggacgaca	780
tgccactggt	gcggctggag	gtgggcagat	gagtgggtgc	ggcccagca	ggcgggtggtg	840
aggttaccgc	wtggaaacak	tkttcgscgg	cagytccctgg	gactggatcg	gcttataccg	900
ggtgggtttc	cgccattgca	aggactatgt	ggnntatgtn	tgggccaac	atgaagatgt	960
ggatgggaat	acataccagg	taacattcag	tgaggaatca	ctgcccagg	gccatggaga	1020
cttcacctcg	ggctacyata	gtcacaacca	cagcaccctc	atcggcacatca	ctgaaccctt	1080
ccagatctcg	ctgccttcct	cggagtgtggc	cagcagcagc	acagacagct	caggcaccag	1140
ctcagaggga	gaggatgaca	gcacactgga	gctccttgca	ccaagtccc	gcagccccag	1200
tccctggcaag	tccaagcgac	accgcagccg	cagcccggga	ctggccagggt	tccctgggct	1260
tgccctacgg	ccctcatccc	gtgaacgccc	tgggtgccagc	cgtagccctt	cacccagag	1320
cgccgcctg	tcccagtggt	ctcctgacag	gagcagtaat	ggcagcagcc	ggggcagtag	1380
tgaagagggg	ccctctgggt	tgccctggccc	ctgggccttc	ccaccagctg	tgccctgaag	1440
cctgggcctg	ttgcccgctt	tgccgcctaga	gactgttagac	cctgggtggtg	gtggctcctg	1500
gggacctgat	cgggaggccc	tgggcgccaa	cagcctgtct	cctagtcccc	agggccatcg	1560
ggggctggag	gaagggggcc	tggggccctg	agggtggggt	aggcagatgg	gccaaggtga	1620
ccaccattct	gcctcaatct	tttgcaagcc	caacctgcctc	tctcctgctg	ctcctccagc	1680
tgtatctgca	cctgcctctc	tgtcctggcc	aggggtggac	aactgggggtc	ccccaaaact	1740
cagtccctggc	acctcaactg	tgacaatcag	caaagcccca	cccaggcccc	catctgggat	1800
gatgggagag	ctctggcaga	tgtcccaatc	ctggagggtca	tccattagga	attaaattct	1860
ccagcctcaa	aaaaaaaaaa	aaaaaaaaaa	ctcgag			1896

<210> 15

<211> 1230

<212> DNA

<213> Homo sapiens

<400> 15

gggcgaatta	accttagcat	atgatagtgc	tttttagtgtt	tcccaccacc	cttttcagag	60
taagcaatta	gacgtctaga	gtttcttttc	ctctatccct	ttgggtaaat	agcaaccagt	120
ggtgcccgc	accacgcctg	gctgattttt	tgtatttttg	gtagagatgg	ggtttcacca	180
tactggccag	gctgggtctca	gactcccaaa	gtgctgggat	tacaggcgtg	agctaccacg	240
cctggccaag	aggcagtttt	tgaattccaa	aattatggag	tgatatctgg	gagtcagcct	300
tcatcagtga	tgctgcttgt	gctttctcca	gcttcttggg	agactgggca	ccaggagact	360
tggagggtct	tccctatctt	acctaactcc	aagtatgcat	gtttttgaaa	aaaatattta	420
agattttctag	agaagatcac	aatcttctcc	agtgtctgtg	taatccattg	tggagcagtt	480
tattacagct	gacctactgt	ctcctgtctg	ttttaggagc	ggtgacttag	gttatcacc	540
cttagtggtt	tcctgtcctc	tgccctctca	tctccacctc	aaatttagtt	tttggtaaaa	600
gatcatgcta	tctaaccctg	tcatcattgc	aaaacaaact	tattttatgt	tatcatgatg	660
ctgacattag	caagctgact	catgggttaa	tgtagtggtt	tcttactact	ggtcactggc	720
tgacagtcgg	aaaagtgatg	ccctattttt	tttttttttt	ttttgagatg	gaatcttgct	780
ctggtgccc	ggctggagtg	cagtgggtgc	atctcggtc	actgcaacct	cttcctccca	840
ggttcaagcg	attctcctgc	ctcagccacc	tgagtagctg	ggattacagg	cacccgctat	900
catgcccagc	taatttttgt	atttttatta	gagaccatgt	tggctagact	ggtcctgaac	960
tccctgacct	atgatccgcc	caccttggcc	tcacaaagtg	ctgggattac	agggtgtgagc	1020
cactgcgtcc	ggctcctgtc	ctgtttttat	ctgaaattat	gttagccttt	gcctagtatt	1080
ggtgaattaa	ccttagcata	tgatagttct	cttaattgtt	cccaccaccc	ttttcagagt	1140
aagcaatttg	acgtctacag	cttcttttcc	tttatctctt	tgggtaaata	gcaaccaatc	1200
caacagcaaa	aaaaaaaaaa	aaaactcgag				1230

<210> 16

<211> 1744

<212> DNA

<213> Homo sapiens

<400> 16

```

ggcagcagca ggcgcggggc gcactgtcct agctgctggt tttccacgct ggttttagct      60
cccggcgtct gcaaaatgaa gattgaggag gtgaagagca ctacgaagac gcagcgcctc      120
gcctcccaca gccacgtgaa agggctgggg ctggacgaga gcggcttggc caagcaggcg      180
gcctcagggc ttgtggggca ggagaacgcg cgagaggcat gtggcgctcat agtagaatta      240
atcaaaagca agaaaatggc tggaaagagct gtcttgttgg caggacctcc tggaaactggc      300
aagacagctc tggctctggc tattgtctcag gagctgggta gtaaggctccc cttctgcccc      360
atgggtgggga gtgaagttta ctcaactgag atcaagaaga cagagggtgct gatggagaac      420
ttccgcaggc cattgggctg cgaataaagg agaccaagga agtttatgaa gtgaagtac      480
agagctaact ccgtgtgaga cagagaatcc catgggagga tatggcaaaa ccattagcca      540
tgtgatcata ggactcaaaa cagccaaagg aaccaaacag ttgaaactgg accccagcat      600
ttttgaaagt ttgcagaaag agcagtaga agctggagat gtgatttaca ttgaagccaa      660
cagtggggcc gtgaagaggc agggcagggt tgatacctat gccacagaat tgcaccttga      720
agctgaagag tatgtccctc tgccaaaagg ggatgtgcac aaaaagaaag aaatcatcca      780
agatgtgacc ttgcatgact tggatgtggc taatgcgcgg ccagggggg acaagatatc      840
ctgtccatga tgggccagct aatgaagcca agaagacaga aatcacgaca aacttcaggg      900
gagattaata aggtgtgtgaa caagtacatc gaccagggca ttgtgtagct ggtcccgggg      960
gtgtctgttt ttgatgaggt ccacatgctg gacattgagt gcttcacctc cctgcaccgc     1020
gccctggagt cttctatcgc tcccatcgtc atctttgcat ccaaccgagg caactgtgtc     1080
atcagaggca ctgaggacat cacatccctc caccgcatcc ctcttgrcct tctggaccga     1140
gtgatgataa tccggaccat gctgtatact ccacaggaaa tgaaacagat cattaaaatc     1200
cgtgcccaga cggaaggatc aacatcagtg aggaggcact gaaccacctg ggggagattg     1260
gcaccaagac cactctgagg tactcagtc agctgtgac ccgggccaac ttgcttgcta     1320
aaatcaacgg gaaggacagc attgagaaag agcatgtcga agagatcagt gaacttttct     1380
atgatgccaa gtctctcgcc aaaatcctgg ctgaccagca ggataagtac atgaagttag     1440
atggctgagg ttttcagcag taagagactc ccagggtgtg cctggcctgg gtccagcctg     1500
tgggcgcttg cccctgggct tggggctgcc gtccctcctc aggcgtggtc tgcagcgtg     1560
tcagttcagt gtggaaagca tttcttttta agttatcgt actgttctct tggttgcttt     1620
gaaagaacct ttcttacct ggtgtgtttt ctataaatct tcataggtta ttttgattct     1680
ctctctctct ctctctaagt tttttaaaaa taaacttttc agaacaaaaa aaaaaaaaaa     1740
aaaa                                              1744

```

<210> 17

<211> 1714

<212> DNA

<213> Homo sapiens

<400> 17

```

ggcagagaa atgtagtaac caattacaga gacttcttta aagatataac tgttgtgctt      60
ttttgaaggc aaagaacctg ccctgaaaaa gaatctgacc tgaacacatg ctgtcctaag     120
cagttagtct agtgtgtctc tggggctctat gcaggtttgc agtgtacaga ctgttatctc     180
ctgtctcagt atttgccctt cttatggaat tacttttctt cctcttacta cttattttcc     240
tcctcttcatt ttgtctcatt agttgtatta gccagatata tatttacttg taggggaggg     300
ggaatgtcaa atttcagttt ctaggaaatta tatatggcat gagtcatcgc tatagcatca     360
cagaaatctt catcaacgtc tatctagaac agtcaggagt tctagttatg agttctgtaa     420
tagatatatta ttgaacatgt gagaacaatg taaagccaat agttggttca ctcaaatatt     480
tactgagcac tgggatacct ccatgaataa aacaaagatc ctaacccttg tgaagcttca     540
ttttaatgca gggggagaca gaaataaacc ataaacttaa caaataggta tattttatag     600
taaatacagta tgtaatagat gcagtggggg ggcagggaat atgaaactgt gtaaagggaa     660
attgatagtg cttggggaca gtggctgtag gttacactac tcagtatggg atcaggaaag     720

```

gcctccttaa	gaagaaatgt	gtgtaggcca	agtgccgtgg	ctcacgcctg	tgggtcccacc	780
gctttggggg	gccaaggcgg	ccggatcacc	tgagttcagg	agttcaagac	tattctgccc	840
aacatggcga	accccatctc	tcctaaaaat	acaaaaatta	cccgggcctg	gtggtgcaact	900
cctgtagtcc	cagctacatg	ggaggctgag	gcaggagaat	cacttgaacc	cgggatgtgg	960
aggttgcagt	gagctgagat	tgagccactg	cactccagcg	tgggcgacag	aaagaagagt	1020
gtgcaaagac	ttaaaggagg	tggggggagt	aagccaaaca	gatagaggga	aaaagcattc	1080
caggcacagc	cacagccatg	tcctaagaat	catgcaatgc	gttggaggag	ctgcaaggag	1140
gctggtgagt	ctgggtggag	tgagcaagag	gagtagaaag	aaatgcagtg	cgagaggcat	1200
caatacctgg	ggctgttcct	ggagcacctt	atatactgtg	aggactgctt	ttcactagga	1260
gttagagagc	acttttgggt	tgggggtttt	tttaattgtt	ttattgagct	aaaatttaca	1320
taccatacaa	gtcaccatt	tgaagtgtaa	aattcagtgg	tttttagtat	attcacagat	1380
atattcaact	attaccacag	tcaatttttag	aacattttca	ttggctgggc	gtggtggctc	1440
acgcctgtaa	tcccagcact	ttgggaggct	gaggcgggtg	gatcgcctga	ggtcgggagt	1500
ttgagaccgg	cctggccaac	gtggtgaagc	cccatttcta	ctacaaatac	aaaaattggc	1560
caggcgtggg	ggcacgcacc	tgtaatccca	gtactcggg	gggctgaggc	gggagaattg	1620
cttgagcccc	ggaggcggag	gttgcatcac	gccattacac	tccagcctgg	gcaacaagag	1680
cgaaactcct	tctcaaaaaa	aaaaaaaaaa	aaaa			1714

<210> 18

<211> 1419

<212> DNA

<213> Homo sapiens

<400> 18

aattcggcag	agatttacct	tagcttactg	taattttttt	aacttaataa	gctttttaac	60
ttttttttta	actttctgtc	tttttaaata	acacgtagct	taaaacagcc	tcataatatg	120
gatccactgt	agtatatgtg	gtctgttgtg	ttgacagaaa	tgtcattagg	tggtgcatga	180
ctaattactt	ttgggcctca	gcctcatctt	gtactgtttt	ttcccttgct	ttcctttgct	240
ccagttgtgt	ggggcctcct	tcattctcct	gtgtccacca	tattctccac	caggaatctt	300
ggcacttggt	cttttctcct	gcctggaatg	cctttctttg	ttaacagtta	ttcctggctt	360
ttagagtttg	gctcaagcat	cacttttaat	gaaattttct	gtgtgcgtgc	ttgcacattt	420
tcataagata	aagagattgg	actgtactga	tattcatcac	cgactccaac	aagagcgaaa	480
ctccatctga	aaaaaaaaaa	aaaacaaaaa	agttacaact	ttggcaccaa	ctcaggccta	540
agtgtcatct	tccctgaatt	actcctgtgc	ttctgcatgc	atctatctct	tctctattat	600
ctctttggat	gtctatgtcc	ccttccaaac	tctactgtat	taaatgtcct	gttatgatgg	660
tactcacac	cctctcaggt	gtttaggaag	ccaataaca	tcaattgagg	acgtaagtaa	720
gagatccagt	cctcagcctg	taatcacacc	tttctgcccc	ctgaaaagat	ttaccaccat	780
aatctgaagc	atgtcctcct	aaatggaccc	ataactaatc	atagttcact	cattagcttt	840
catatatttg	atgtctccag	tatcaacacc	ctcccagata	ccccatccag	tatgtctagt	900
tatcgatttg	atgccaccac	ttattgggtc	ctgattcttc	atatggccaa	aacaaccctc	960
tggatatcca	cagtgaatat	tatgtcaata	ccaacgtcat	ctcagctgaa	gcagtgtgca	1020
tggttacagc	tgctaagatc	aaaaaccttg	ggctcatgcc	cgattcctct	ttcatgctcc	1080
tatcatccac	attagaaaat	ctagctgtct	gcaaggcatg	gtggctcgtg	cctgtaatcc	1140
cagcacttgg	ggaggcagag	gccggcgagt	ctcctgaggt	cgggagtttg	agactagcct	1200
ggccgacatg	gtgaaacccc	gactttacta	aaaatagaaa	aattagccga	gtgtggtggg	1260
acatgcctgt	gatcccagct	actcgggagg	ctgaggcagg	agaatcactt	gagcccgagg	1320
ggtgaagggt	gcagtgggct	gggattgtgc	cactagactc	caggctggac	aagagagtga	1380
aactcctgtc	caaaaaaaaa	aaaaaaaaaa	aaaaaaaa			1419

<210> 19

<211> 1165

<212> DNA

<213> Homo sapiens

<400> 19

gggataagag	cttcagagtt	tattctacag	aaacagggaa	attgactcag	attgtatttg	60
gccattggga	tgtggtcact	tgcttggcca	ggcccgagtc	atacattggt	ggggactgct	120
acatcgtgtc	cggatctcga	gatgccaccc	tgctgtctctg	gtactggagt	gggcggcacc	180
atatcatagg	agacaaccct	aacagcagtg	actatccggc	accaagagcc	gtcctcacag	240
gccatgacca	tgaagttgtc	tgtgtttctg	tctgtgcaga	acttgggctt	gttatcagtg	300
gtgctaaaga	gggcccttgc	cttgtccaca	ccatcaactgg	agatttgctg	agagcccttg	360
aaggaccaga	aaactgctta	ttcccacgct	tgatatctgt	ctccagcgaa	ggccactgta	420
tcatatacta	tgaacgaggg	cgattcagta	atttcagcat	taatgggaaa	cttttggtctc	480
aaatggagat	caatgattca	acacggggcca	ttctcctgag	cagtgcaggc	cagaacctgg	540
tcaccggagg	ggacaatggg	gtagtagagg	tctggcaggc	ctgtgacttc	aagcaactgt	600
acatttacct	tggatgtgat	gctggcatta	gagcaatgga	cttgtcccat	gaccagagga	660
ctctgatcac	tggcatggct	tctggtagca	ttgtagcttt	taatatagat	tttaatcggt	720
ggcattatga	gcatacagaac	agatactgaa	gataaaggaa	gaacccaaaag	ccaagttaaa	780
gctgagagca	caagtgtctg	atggaaaaggc	aatatctctg	gtggaaaaaaa	ctcgtctaca	840
tcgacctccg	tttgtacatt	ccatcacacc	cagcaatagc	tgtacattgt	agtcagcaac	900
cattttactt	tgtgtgtttt	ttcacgactg	aacaccagct	gctatcaagc	aagcttatat	960
catgtaaatt	atatgaatta	ggagatgttt	tggttaattat	ttcatatatt	gttgtttatt	1020
gagaaaaggt	tgtaggatgt	gtcacaagag	acttttgaca	attctgagga	accttgtgtc	1080
cagttgttac	aaagtttaag	ctttgaacct	aacctgcac	ccatttccag	cctcttttca	1140
agctgagaaa	aaaaaaaaaa	aaaaa				1165

<210> 20

<211> 2280

<212> DNA

<213> Homo sapiens

<400> 20

ggcacgagga	attatgaagt	tgtgtgcccc	atgatgggga	gtggactttg	agatgagaca	60
ctggcctgca	ttataaggga	cctgtcattt	ccactggatt	ttagttttgtg	tggcttaaaag	120
agaatgcctt	tccagttagg	tatgctcgtg	gctttgaata	taaatgggaa	cagtctaacg	180
ctagctattt	tgctcctttc	cttctgatcc	ccagctggtg	gaaatcggtg	ctgcgtactc	240
agcaacctta	catagggcag	gtatggtctc	atgagctggg	agcaagacag	gttttctttc	300
ttgaaatgag	gtccccccac	ttccctttgt	ctttgtctgt	ttgtgtcttt	ctttccattg	360
tcttgctgct	tctctcctt	ggatctgtag	acactcagag	aactgcctgg	ccagacccta	420
tagaggtaca	attaggtcac	tctagggcac	ctgaccatgg	gtggcagttt	tccacaaaat	480
tcacttggtc	tgtgggttga	aggaaaatac	aaaccagctg	gcaagttcta	ggcctcttgt	540
cttcttggtg	tcttctagtc	aaatttacag	tatatccctt	ctgatttaat	atttgcagga	600
aactgatgca	tggagatgct	gacatccccc	gtgttaattc	tggggcaccc	tgatcctatt	660
ctctaggatg	gcattgcccc	tttctgagcc	atcaatccaa	gtcctttccc	cgtattttcta	720
tgtgggttgg	atataaattca	gtgagacact	ctttcttgct	cctgtacttg	aatgtaggag	780
tgtattgcag	tggccctgaa	gatcccagta	gagctctggg	agggttgatt	aggacagagt	840
gatttccacc	ctgcccctcc	tcaagcagag	gagctttctt	caaccttttt	ctctttgaag	900
aaattgagac	tcagggaaga	agtattgtcg	cagatgtgat	gcggtgagtc	actgggtttt	960
aaacctctgt	tttggcgatg	ggctgggtat	ggatgagtta	aggtttatcc	ccctaaatct	1020
gtattacact	ctaagtcaac	atttgtgctt	taggctgggc	gcagtggctc	acgcctgtgg	1080
ttccagccct	ttgggaggct	gaggcgggtg	gatcacaagg	tcaggagatc	aaggccatcc	1140
tggctaaccac	gggtgaaacc	catctctgct	aaaaacacac	acacacacaa	aattagccgg	1200
gtgtgggtgg	agggtgcctgt	gtcccagct	gctcgggggg	ctgaggcagg	agaatggcgt	1260
gggcccggga	ggcagagctt	gcagtgggct	caccgcactc	cagcctgggc		1320
aacagggcga	gactccgtct	caaaaaaac	aaaacaaaaa	aacaaaaaaa	accattgtgt	1380
ccttaaattg	aatatccttg	atagaaaagg	gtagggcctt	agcgtactgg	ctctatggaa	1440
tacgtactct	ctgtctttca	gtagaattca	gcagtcattg	ttcacctgct	ttttgccaga	1500

tgttgagcta	attgctattg	gagtaggatg	gtaaagattg	gtcttggtat	gcttggttga	1560
gaaggcacia	cttttttacac	atgtgggcct	gcaaatatct	atgcagattt	ttttgagact	1620
ttaaaaagca	tgtttaaaag	attattttctt	ctaatagaga	ctccaaaagt	actcaattac	1680
tgtcttatcc	tctgggatcc	tagttcctat	tacttgtgct	tgattttccct	ctgctgatta	1740
gataagttac	cttgaaaagag	gcaaagaaaag	ccaaatggga	attgacttgt	ttttcttcct	1800
gtcattaaaa	aaaaatggaa	ataattttcaa	atttatagaa	gttatacaaa	taaaattata	1860
caaagaatat	ttgtaatect	ttttatccag	aacctcctat	tgttcgtgtt	cattttacc	1920
catttgcttt	atcatttgca	cattctctct	ccatgtatca	atacataatt	gttttctgaa	1980
tcattaagag	gagagataat	tggcccaggc	acggaggctc	acacctgtga	cctgagtact	2040
ttgggaggct	gaggcaggag	gattgcttga	gccaggagt	tcaagaccag	cctgggcaac	2100
agagttagat	cctgcctcta	ccaaaaaaaa	aaaaaaaaat	tagtcagaca	tggtggcatg	2160
ccgctagcta	ctttggaggc	tgaagtggga	ggttgagact	gcagtgagct	gtaattgtgc	2220
actgcgctct	agcctgggtt	acagagggag	acccttgctc	aaaaaaaaaa	aaaaaaaaaa	2280

<210> 21

<211> 677

<212> DNA

<213> Homo sapiens

<220>

<221> SITE

<222> (53)

<223> n equals a,t,g, or c

<400> 21

aactggtgga	tcccccgggc	ctggcaggaa	ttcgggcacg	agaactgtat	ggngttgggtg	60
atggatgaac	tgcaaggatc	agtgaacacag	ctgcaggcct	ttatggatga	aagtacccag	120
tgcttcagaa	aggtgtcagt	acagctcgga	aagagaagca	tgcaacaatt	agatccctca	180
ccagctcgaa	aactgttgaa	gcttcagcta	cagaaccac	ctgccataca	tggatctgga	240
tctggatctt	gtcagtgact	ttatgagagt	ttctgccaca	aggtgcccaa	gaggagagga	300
atgggaagag	tgcccagca	cgtggtgact	gcgtgatttc	tgctcgttgc	ctttaaagrt	360
aactggcagg	actgactgta	gaacactttg	acttttttca	aaaagtgatg	gaatttgtac	420
atccaaatga	atattgtata	gacaattttc	ccaggaatgt	gcaaaatgct	tgaaagtcca	480
aacttctttt	ttgaaatgat	cttcagatcc	agtggcccat	tcttttatct	ttatcctgtg	540
aaggtgtttt	tcagggtttg	aaacaatcca	aaaatcattt	aggaccaagt	ctaaggaaac	600
atttttagtg	ccaagttgga	ttccgattgt	aaaggaatga	tactaatttt	ctagcatggc	660
tctgaagggtg	atttttag					677

<210> 22

<211> 858

<212> DNA

<213> Homo sapiens

<400> 22

tttttttttt	tttttttttt	tttttttttt	tttttttttt	tttttttttt	tttttttttt	60
gaaaagaaaa	agagtttact	tgaaaagggt	caaggaggtc	agacttcaga	aaaactggat	120
aagcaagtga	aaaatataac	aggtatttaa	agaaaaatat	aaagaagatt	accatgatta	180
ctaagtttgc	attaatttgt	tcatggagaa	aaaacaaaat	cttatcacat	gatgatcata	240
ttcagaaaga	tcttgcaaca	tgagctaaat	aaaagtcaag	aataaaatgg	tgtttgctgg	300
ggatctggga	ttataagaaa	tactcaatgt	ttcattatct	tgggaccaga	aattgacagg	360
gtaacgctgt	tatttttggg	taatcttagg	gtacaagggt	cttcatcatg	atggaaacaa	420
acagggcatg	atttctgctt	tgcttcagaa	agcataaagc	aaatgtcact	actacaaaat	480
atagatagga	tattttcttat	aactctaate	actgttgtaa	gtaaataaaa	atccaaagtc	540

aaatcataga	ctctcaatta	ccatggtttt	tacacatttc	taccatccct	tcctgggtgca	600
ctgtattgag	ccatactgag	ctatatTTTT	ctacttcaca	acttttcact	tacgggaaga	660
cagctattgt	ccccacacc	accctaattg	gggcaagagg	accacttgac	cctcatatat	720
ttaacactct	tgaccctcat	atatttaaca	ttcataatta	tgtattgcct	aaatactcac	780
tgttgtagtc	actgaaatgc	ctgtgaaata	atttttaaaa	ctgtattttac	aaaagaaagc	840
ctattttctag	ctcgtgcc					858

<210> 23

<211> 1085

<212> DNA

<213> Homo sapiens

<400> 23

ctgcaggaat	tcggcacgag	gtgcattcttg	cccatttgatt	tctaaatgta	ttactacttt	60
aaattaatcc	tgaatctttt	cccaggctta	agtgggataa	tgttttattg	tagatgcata	120
tttcttggt	ctaccagtc	tttctttgaa	gactttatca	tcctattttc	tgaatccagt	180
ggctgacttt	aatcttctct	ggaggaacta	gataatttct	agactaatgc	ttactcat	240
gatccagatt	gtaatttctg	aactccttct	tccaaataga	atcaaaacaa	gaaaggggaa	300
agcctctcaa	agcaactgtg	cgtaataaat	gaaacactct	ttttttctaa	tccaaggagg	360
gtttcatact	ttttcttagt	ttcttgccct	cttcccttct	gatcaataat	tgtaatagga	420
aatttgcaat	tgtgccaata	ctcagattca	atactgaact	actttcttgc	attgtaattc	480
aaattccaag	gttaacaact	agctgtatgt	ttccaaaaca	atcttattgt	atatgtattt	540
tcttaggtga	agtttccaga	aatgattttt	tttttttgca	gagccaaaca	cacatggtaa	600
tttaaaaaaa	taatgcacgt	atgtggtaaa	aacagtaaaa	gcaagggtatc	aagtaaaaag	660
tgaagagtcc	tccctttctc	attcccatc	ctactctcta	attttttata	tatccatttt	720
gtaagctata	atacagagat	tccatatact	cttcacatac	tttccccagt	ggtaacctct	780
tgcataacta	taggacaata	acacactata	ggacaaaaat	caagaaattg	acattgatac	840
aatccatctc	caaaaaaaga	aaaaagaaaa	agccagacat	ggtgatgtgt	gcctgcagtc	900
ccagctaatt	gggaggctac	agtgggagga	tcccttgagc	ctgtgaggcg	gagggtgaag	960
tcagctatga	tcacgccact	gcactccagc	ttgggcaaca	gagccagacc	ctgtctcaaa	1020
aaaacaaaaa	caaacaaaaa	aaaagaaaaa	aaaaaaaaaa	aaaaaaaaaa	aaaaaaaaaa	1080
aaaaa						1085

<210> 24

<211> 1422

<212> DNA

<213> Homo sapiens

<220>

<221> SITE

<222> (44)

<223> n equals a,t,g, or c

<400> 24

gttgtggatg	cgcgagagg	ccggcagcgg	tggcagcggc	acgnaggagg	aagctgagca	60
gggcggcggc	ggcgggtgaa	cctgcggggc	tggggcgcg	gccatgggcc	gcctgcactg	120
cactgaggac	ccggtgccgg	aggccgtggg	cggcgacatg	cagcagctga	accagctggg	180
cgcgagcag	ttctcagccc	tgacagaggt	gcttttccac	ttcctaactg	agccaaaaga	240
ggtggaaaga	tttctggctc	agctctctga	atttgccacc	accaatcaga	tcagtcttgg	300
ctccctcaga	agcatcgtga	aaagcctcct	tctggttoca	aatggtgctt	tgaagaagag	360
tctcacagcc	aagcaggtcc	aggcggattt	cataactctg	ggtcttagtg	aggagaaagc	420
cacttacttt	tctgaaaagt	ggaagcagaa	tgctcccacc	cttgctcgat	gggccatagg	480
tcagactctg	atgattaacc	agctcataga	tatggagtg	aaatttgagg	tgacatctgg	540

gagcagcgaa	ttggagaaaag	tgggaagtat	atttttacaa	ctaaagttgg	tggttaagaa	600
aggaaatcaa	accgaaaatg	tgtatataga	attaaccttg	cctcagttct	acagcttcct	660
gcacgagatg	gagcgagtca	gaaccagcat	ggagtgttct	tgctgatttc	tgtccctgca	720
tctcccctrg	ccccgttccc	tgccctcctc	ccttccctgg	gtgactgctc	tgagaggcac	780
ttcactcaca	ggcctgtggg	atgctccatg	gggccctgct	ggctccatgg	ggcccagggtg	840
caaagggttt	ctgaaaaaca	gcaggattaa	gtactgaaag	agcccaacac	aattaccctg	900
taaaactctct	gttagggcaa	ccaccaccac	ctgtcttcca	ggacacattt	ttagatactc	960
tgacaggcca	ctgcatctca	gattcagggg	agaaaataag	ttgtcacctc	cccttcaaag	1020
ttccagagta	aacaaatggt	gccatcattc	aagataacat	gctgatcacc	ctcctcccaa	1080
aaagcaagag	cttgtttatg	gctgaggaat	cggcggattg	tctgaatgac	acatatacag	1140
agccccacg	gatttctgca	cactctgggt	ctgtgctggt	ggaacattgc	caatcagttc	1200
ttaatgaggg	acctgtgtgt	aaatacatgc	ttggtcttct	ctgcagagaa	ctgaggctaa	1260
actctgtccc	tacttctggt	tttgccctgt	catgtcgtaa	cgagggtgggc	cttttgagggc	1320
cattttagtt	tgagttcgag	ccaaccacct	ctgttggtta	gatgatgaat	aaaaagggtc	1380
tgaagaaaaa	aaaaaaaaaa	aaaaaaaaaa	aaaaaaaaaa	ag		1422

<210> 25

<211> 419

<212> DNA

<213> Homo sapiens

<400> 25

ggcagagcat	gaacttgctg	ctagtttctg	ttccctctca	aattgcccag	gctgtcctgt	60
gagtgcagc	agctatttctg	tgggcctcct	ctggaatcat	gagaagtcac	ccaaacaatc	120
tcagttttct	agctcactcc	gtcttgacat	ttctacactg	tcatecttgg	ttttcttgga	180
aattaatttg	cttttcttca	ttgtctttct	ttggagctgc	tttccttttg	ttggttacta	240
ttttattttt	agcttctcac	accataccga	catatgttgg	ttattctttt	agacatgttt	300
tttggtgttg	ttgtcacctg	gaacttttgt	atcttgaata	aatttgggga	tcaaatacaa	360
aaaaaaaaaa	aaaaaaaaaa	aaaaaaaaaa	aaaaaaaaaa	aaaaaaaaaa	aaaaaaaaaa	419

<210> 26

<211> 1531

<212> DNA

<213> Homo sapiens

<400> 26

aaagaggggg	cctccttccc	tggaggccat	ctgatgaggg	tcgatggggc	agtttgagaa	60
ccatttgtgg	caagcgttta	aatcccaggc	tgctttttgc	cccagtacct	ggtatccagc	120
catcgattga	tgtgctacga	gagaagatga	accaatttta	ttattagcct	gaacaaaaca	180
taattaggag	gagaaatttg	cttctgctgt	tgcatgtgct	gcagcaaaga	agttacataa	240
ggttcattcg	gatgacttgg	gaacaccttc	cctacttccc	tattcctgtt	cccgtttctc	300
tggcttgaa	caagtaaata	gagggtccaga	gcctccactc	tgcttcttag	aaattcccaa	360
gacagaaatg	tgctcaagga	gcagatctaa	gaccctcagc	ctatagcgga	catggaacaa	420
gggcacaaat	cgaataagtc	agtgcattgt	catgatgaac	atgtggacaa	aggcaagaaa	480
agggagaagg	gacaagagag	aggagcagac	tggaggagaa	ggaataaaga	gctgaggcat	540
agctctgctc	ctgtgatgtt	agaggaaggg	aagggtggaaa	aaaaaggcat	tgaattatct	600
tcagacaatt	tcagaaggac	cagatggatt	ctggaaggaa	ttaactctgt	tatgtcctgt	660
actagtatca	ttactacact	ctaagatggg	caaacgaaag	gatttggagc	attttggcag	720
atactctttg	ggtgattgag	agggggaggt	aggaggtcat	ggctgttggg	gtgactatat	780
gggggtcgca	tttgggctga	aagttaagac	tctgtctcaa	aaaaaaaaag	aaagcaaaga	840
aatgtcaac	tactatttta	agtaaacaaa	taatacaaat	tgtttcta	ctttatttta	900
ggagatgttg	caaatatgga	ccagagactg	cagaaagcta	gccagatcaa	aggccagcag	960
ttaaaggcag	taactatagt	taactcccag	ttgacccaat	agaaggaaga	gccttgaccc	1020

gagtaacacc	cagccactga	ttttttttta	atcttttctt	tttaactttg	ggatgcatca	1080
tggtgataat	aatgatgata	aataatcaga	tttgcccttg	ttattagagt	tgacttggtc	1140
tgctatttaa	atgagtttct	aattccctgc	tcacaccaga	ctgacaaccg	aaaccttctg	1200
ggtggcaaa	ggaatcagcc	tgccctgggt	taagattttt	ccatgggtta	tctccaaagt	1260
gaggaacct	cgtgtgagta	attgagagct	gacagcattt	cgttggatca	gttttgccct	1320
gtcagacagg	gagagggtct	agccttggaa	agacaatagg	gtaagacaag	aggaagttga	1380
cagagtgtga	cttctatcca	tttctcagcc	ttgcttgac	aacctgccct	agctcagggt	1440
aagtggccc	tgcttctttt	gtctcttatt	tgctcatttc	caacaaacct	ttcatttggt	1500
ccaagggtta	aaagaaaaaa	aaaaaaaaaa	a			1531

<210> 27

<211> 957

<212> DNA

<213> Homo sapiens

<400> 27

gagattaagg	cctttttccac	acgcattaat	agtcceattt	ttctcttgcc	attttagtct	60
ttgcccattg	tcttattggc	acatgggtgg	acacggatct	gctgggctct	gccttaaaca	120
cacattgcag	cttcaacttt	tctcttttagt	gttctgtttg	aaactaatac	ttaccgagtc	180
agactttgtg	ttcatttcat	ttcaggggtct	tggtgccttg	tggtgctccc	caggtggcct	240
ggaggtgggc	aaagggaagt	aacagacaca	cgatgttgct	aaggatgggt	ttgggactag	300
aggctcagtg	gtgggagaga	tccctgcaga	acccaccaac	cagaacgtgg	tttgccctgaa	360
gctgtaactg	agagaaagat	tctggggctg	tcttatgaaa	atatagacat	tctcacataa	420
gcccagttca	tcaccatttc	ctcctttacc	tttcagtgc	gtttcttttc	acattaggct	480
gttggttcaa	acttttggga	gcacggactg	tcagttctct	gggaagtggg	cagcgcatcc	540
tgcaaggctt	ctcctcctct	gtcttttggg	gaaccagggc	tcttctcagg	ggctctaggg	600
actgccaggc	tgtttcagcc	aggaaggcca	aaatcaagag	tgagatgtag	aaagttgtaa	660
aatagaaaaa	gtggagtgtg	tgaatcggtt	gttctttcct	cacatttgga	tgattgtcat	720
aagggtttta	gcatgttctt	cctttttctt	accctccctt	tttggttctt	attaatcaag	780
agaaacttca	aagttaatgg	gatgggtcga	tctcacaggc	tgagaactcg	ttcacctcca	840
agcatttcat	gaaaaagctg	cttcttatta	atcatataaa	ctctcaccat	gatgtgaaga	900
gtttcacaaa	tctttcaaaa	taaaaagtaa	tgacttagaa	aaaaaaaaaa	aaaaaaa	957

<210> 28

<211> 1390

<212> DNA

<213> Homo sapiens

<400> 28

ggcacgaggg	agtctgatgg	ggagaagaag	tacccatgcc	ctgaatgtgg	gagcttcttc	60
cgctctaagt	cctacttgaa	caaacacatc	cagaagggtg	atgtccgggc	tctcgggggc	120
cccttggggg	acctggggcc	tgcccttggt	tcacctttct	ctcctcagca	gaacatgtct	180
ctcctcgagt	cctttgggtt	tcagattgtt	cagtcggcat	ttgcgtcatc	tttagtagat	240
cctgaggttg	accagcagcc	catggggcct	gaagggaat	gaggcagctg	ctgtgtcccc	300
acggaaacaa	ccatctgggg	actgctggga	aatgctgtga	atgctggagg	aagtgtgttt	360
tggtttctgt	agctgagaga	tttttattca	tttttaactg	ccccccaacc	ccactccaac	420
tccttctcca	ccacccattc	tcccaatggt	ctttagaaat	agattttcat	ctgatattct	480
gcagaaatat	caatgagact	tggtatggga	caggggcaga	aaacactaca	taggcctcca	540
aggcaaaacc	agtcccagtt	tctttaatgg	gaagaagctg	gaattcctgg	tgctcaattc	600
ttagtgaacc	caatcctata	cccaaactta	tgatattctg	ggacctcagt	gattttgggt	660
ccctcccact	tctctagttc	gtcatcctcc	cttcccatat	ccttcaaaaag	aaccacacta	720
gggtctccac	ctacttatac	aatgcggatg	cccaactgtt	tttaagggaag	ccagaagcat	780
cccatggacc	atgggggtgag	tgtcctccaa	gagccccctg	agctcagccc	tctgcctgga	840

gggctccaga cctttctgag ccttgcttgg aggcgagcat tttcactgct aggacaagct	900
cagctgttga ggacaccccc accccaaatt tcagttctta cgtgatttta accattcaac	960
atgctgttgg gttttaattc tctaattatt attattattg ttattatttt ttaggaccag	1020
ttgtagtga ttgctactga aagctatccc aggtgataca gagctctttg taaaccgcag	1080
tcacacatta gggtagtat taaactttgt ttagatgtac cataattaac ttggctagtt	1140
gattgtttga agtctatgga agaaatagtt ttatgcaaaa ttttaaaaaa tgccagtctg	1200
gtcagggaag taggggggtt caatgctgtt gggaaccagg aagggtgggac agccggcagg	1260
tagggacatt gtgtacctca gttgtgtcac atgtgagcaa gcccagggtg accttgtgat	1320
gtgaattgat ctgatcagac tgtattaaaa atgttagtac attaaaaaa aaaaaaaaaa	1380
aaaaaaaaa	1390

<210> 29

<211> 1590

<212> DNA

<213> Homo sapiens

<400> 29

ctgcaggaat tcggcacgag cggcacgagg ttctgtctc acctcactct ggtactcgcc	60
tcttggggcg gctcagccca ttcatgggga tggcaccaag cggccatgct cagtcttcca	120
gccccgctga gggtaaaccg aggcctctgg cagctgtgca cagggtgctg cctctggctc	180
cttcaaggag cactgcctgt cactcgctcc tgggctgtct agccatgtct cccaccccca	240
ctttaccgca gccagctgct gggatcaaag caagtctgtt cttatgttat ttgcctgtat	300
gaaatcattt ctcatattat cacaattcct tcaactcagc ttactcgcgt ggctgcctgt	360
tcataattga aagcagccac cgtgctgttg ctttggtttg gaaaagcata gcacgcactt	420
cccttggttt tcccttccca gagccgaccg cagctggtea gccctctctt cccgctcctg	480
aacctttact tactgacttt gagctctgtg actccgtcgg ttctcgcagg aattaactaa	540
cttaccaatt ggttcaatcc acttgagcgc catagctctg agctcctctg tgtgacatgc	600
cacagatgac tattgcacac ctgggwcctg cccagcagg ccatgcccct cccatgtgcc	660
gtgcctgttg ctgcagctgc cccccgaccc cgccactggc tgcaggaatt cagcctttag	720
aggcagaggc agctgcagcg gccctgagg tcaaacccca gtgtgactgc atagcagtgt	780
tagctggttg gtttcaaact actggattcc aggcaaaggc ctacagattg accttattat	840
ttttgaaaat atgttaaggg ttttttcata gagagagaaa gaatggattt ttttttaact	900
gggaacctcc tgattcttac ggaaaattat ccttctataa gaagatacca gagagattta	960
ttcaaggtaa tttgataacc taaaatcaat tctccatttt ttatcatatg tgggatttgt	1020
tgctaagtcg tgttcaacaa tagcttttat gttcctaaca tatctgaaag cttattttatg	1080
aatggatata ctggattatt gatatactga tttttttttt aatggggaca tttgccattt	1140
tcttccaga aatatgtaat cccctggctg actaggactg ttaaacatag tgtggactgg	1200
atgatgcctt cgacaaacca gagaagccaa gttgggggga gctgggtgct ggagtgggcc	1260
ctgtgcacct cacctggcgg aggtggggg ggctctgtca gcaggaccct agaggagact	1320
ctcattcgat tttaaagaag cacaacgggt cattttcctt tgtatgttcc tagcgcagaa	1380
ctgtttctaa aacaacttga agtatagttt tgttatctaa gcaatttttg ttttaagtaa	1440
gtaagtgtac tagaatgcga agccgttatg gttcagggtt ttaaaaactg gtacagtatt	1500
gtatytgtct catctgttgc actgtatttc aatcatctgt aattaaaatg atcatatgtt	1560
taaaaaaaaa aaaaaaaaaa aaaaaaattt	1590

<210> 30

<211> 1651

<212> DNA

<213> Homo sapiens

<400> 30

cggcacgagc ggaccaccag cccccgaaat gataccttcc teggctctac ggagtgcacc	60
ttggggccaga ttgtgtcaca aaccaaggct actaagccat tattgctgaa gaatgggaag	120

actgcgggca	agtccaccat	cacgatcgtg	gccgaggagg	tatcaggcac	aaacgactat	180
gtgcaactca	ccttcagagc	ctacaagctg	gacaacaagg	atctgttcag	caagtctgac	240
cctttcatgg	aatctataa	gaccaacgag	gaccaaagtg	atcagctggt	ctggagaact	300
gaggtgggtga	agaacaacct	gaaccccagc	tgggagccgt	tccgcctgtc	cctgcattcc	360
ctatgcagct	gtgatgttca	ccgacctctc	aagttcctgg	tgtatgacta	tgactccagt	420
gggaagcatg	acttcacatg	cgagttcacc	agcactttcc	aggagatgca	ggaagggacg	480
gcaaaccctg	ggcaggagat	gcagtgaggc	tgtatcaacc	caagtatcgg	gacaagaaga	540
agaattacaa	gagctcaggg	acggtagtgc	tggcccagtg	cacggtggag	aaggtgcaca	600
ccttcctgga	ttacatcatg	ggtggctgcc	agatcagctt	cacggtggcc	atcgacttca	660
ccgcctccaa	tggggacccg	aggagcagcc	agtccttgca	ctgcctcagt	ccccgacagc	720
ccaaccacta	cctgcaggcc	ctgcgtgcag	tgggaggcat	ctgccaggac	tatgacagtg	780
ataagcgggt	cccagctttt	ggctttgggg	ctcgaatccc	ccccaaacttc	gaggtgtccc	840
atgactttgc	tatcaacttt	gacccggaag	atcctgaatg	tgaagaatct	caggggtcat	900
cgcttcctac	cgctgttgcc	tgccccagat	caactctacg	gccccaccaa	tgtggccccc	960
atcatcaacc	gtgtggctga	gccgggccag	cgggagcaga	gcaccggcca	agccacgaag	1020
tactcgggtg	tgctgggtgt	cactgacggt	gtggtgagcg	acatggctga	gactcgcact	1080
gctatcgtgc	gtgcctcccg	cctgcccagc	tccatcatca	tcgtaggcgt	gggcaatgct	1140
gacttctctg	acatgcggct	gctggatggc	gacgacggcc	ccttgcgctg	ccccgagggg	1200
gtgcctgcag	cccagacat	tgtccagttc	gtgcccttcc	gagacttcaa	ggatgtgccc	1260
ccctctgcac	tcgccaagtg	tgtcctggct	gaggtgccac	ggcaggtggt	ggagtactac	1320
gccagccagg	gcacagccc	tggggctccc	aggccttgca	cactggctac	gactcccagc	1380
cctagcccg	gactgcctcc	ctccggaccg	acactccctc	agcctctcag	tgctgtctct	1440
gaccctcgtg	actccagtga	ccaatgcctc	cacctcttgg	accaggtgtg	ccccctgggt	1500
tctggacgtg	agtgggtggg	cctgctccta	tctctccaaa	ccccataccc	ttcaatgctg	1560
tggccctca	gtgacttcct	tgggtgatcc	tgactttcta	gccattaata	aagagaactg	1620
ctcctagcac	ctcagcctct	aaaaaaaaaa	a			1651

<210> 31

<211> 1720

<212> DNA

<213> Homo sapiens

<400> 31

ggcacgagca	actctccagg	aggttggggag	ggtgcccgct	tctctgctct	gtgcccatta	60
gcctcgccgc	acaccctagg	accattgact	ttggaggagg	ttgggagggt	gccactttgc	120
ccactcctcc	gcagacagcc	ctggcttcac	ctcagcgcca	caggtctaca	ccggaccctc	180
gtgcctaggt	ctgtggcggc	tgcaaacatc	aggacctgct	tctgtcagc	acgtggagtc	240
tctgtgtctc	atggcatctg	ctggcagtga	gcactggcca	tgtggctctg	agccccaggc	300
tttcgggcat	gacaggcggg	cagtaggaca	gcgcagtctg	tccccctgca	cgcgtcaggg	360
tccgagcctt	tcgccattgc	actccagcct	gggtgacagg	gagagggact	ctgtctcaaa	420
aaaaaaactg	aggtcagggg	gggtgagatg	acggtgagag	ctcggacttg	aacgcaggtc	480
ccaccagaaa	cagcagccct	aactctgagc	aaggtctgtg	ctgttcagta	gctctattga	540
gatgtgattt	ccacactgtg	taattcattc	acttacgggt	tacagtccag	tgggtcttag	600
catgctcgg	gttgacagtc	acatcgtctt	cacccccaaa	aggaaacccc	gtgcccata	660
gcagtcgctt	tgtctgcccc	tcgtccccag	ccccaggcaa	ccacaaatcc	atgctctgtc	720
tctgtagatt	tgctgtttcc	agacgtttca	cagcaatggg	ccttttctgc	ctggcttctt	780
taacgttgca	tcacatcttc	aaggctccatc	ccagctgcag	cgtgtcagtg	cctcctggct	840
tttcaactgt	gagtgtgccc	cgttgcatgg	acagaccacg	ttgtgtcac	ctgtttgccc	900
taatgggccc	ctgcttgggt	ctttccacct	ttgggaggct	gtgaatcgtg	ctccagccac	960
actctgaccc	ccgcccggct	ccaagagatg	acaggattgg	cagcttcttc	accaaccagg	1020
gactctgtgg	cctgcgcac	cgccatcctg	ttggctgtgg	taccctgtgc	tcaaccgtgg	1080
cctttgtgct	gcctcgctg	ctgcacagcc	accccttgca	gcacccacct	agagggcact	1140
ggccaggagt	ctgtttccca	cgagaccgtg	aggcagtcct	ttctcatacc	acgatggagg	1200
ccgagtcact	gtgcgctcca	cgatgaggct	gcaagtcact	atgcgttcca	cgcctccgt	1260

gcactctggg	ctctgccacc	ctactctgca	gggtgctggt	catctgggca	tcgccgagct	1320
caggctggca	cgacaggctg	ctcttgggag	taacagacct	tttgggtgct	ttccagagcc	1380
ttaaactgaa	tgcgtataag	aaaggcaggg	aggccaggca	cagtcactca	cacctgtaat	1440
cccagagctt	tgggaggcca	gggcaggagg	atcacctgcg	cctaggagtt	caagaccagc	1500
ctggtcaaca	tagcgagacc	ccatctctac	aaaacatttt	ataaaaaaat	taggcaggca	1560
tgggtggtg	cacctgtaag	tcccagctac	ttgggaggct	gaagtgggag	gatcacttca	1620
gccgggagtt	tgaggctgca	gtgagctgtg	agcacaccac	tgcactccag	cctgggtgac	1680
agagtgaggc	cctgtctctt	aaaaaaaaa	aaaaaaaaa			1720

<210> 32

<211> 1247

<212> DNA

<213> Homo sapiens

<400> 32

gataatgtga	gcaatctcca	gccacataca	gaagtataga	agaatctcac	aagtacaatt	60
tgagttaaaa	acaaaaaccc	agacctaaag	atatacatac	tctgtgactc	caaatttaaat	120
aattgctcat	agaagtcaaa	tagaggttac	cctgtaagaa	tgggttggtg	actagagtga	180
gcacacccag	attctggggg	cctagtcattg	ctctgtgtct	tgacctaatg	atacaggtag	240
atgtaagttt	gtgaaaatgt	aacaagctgt	atgatacgat	ttttctaata	ttgtctcatct	300
gtgtttgaaa	catatacttc	agaaaatgtt	aaggaaaaaa	tagaaaagga	aaagaaaaat	360
agaaaaccta	aataatctag	tacttattaa	agacttgaac	ccgtcataaa	aagcctttcc	420
acaaattaaa	tactaagcca	agcagactta	ctaaattcta	ccaaaagcta	agaagctatg	480
tcaattttat	acaaactttt	caactaattt	atgagcatta	atttgatacc	caaacctgga	540
agggactata	taggaaagga	aaatgataaa	gtcagtccta	attatgacta	tagattgcaa	600
gaacctttac	caaaatagga	gcaaattgaa	tcaaacaatt	tttttaataa	ggcaaaatgt	660
aacaaccaag	ttgggtatat	tcctgggtatg	caacactgct	gcagcattta	aaaactaatc	720
agttatggag	ctagagggaa	gatggcagag	taggaagctc	tgggaatctg	tctccccacc	780
cagacgacaa	ctgcaactgg	agaatctgtc	tgatgttact	atttcggatc	tctgttcacc	840
atttttatca	ggttgggaac	tctgaagtct	attgaaggct	tatgacttcc	aggggaaggt	900
ttgaaaggtt	aatttcagct	cttaacatgg	tagcaactac	ctgttcgtct	accctcagtc	960
ccatagcatg	tagccatgta	cacattttca	agcagctttg	catagggttt	gtgggagcca	1020
gtgtggacaa	taaggattct	gtcctccaaa	tagcaggaat	ctttgtctctg	atagctaatt	1080
actgcttctg	atcatggagg	tgcagccata	cagtgaagca	ttgktgcaaa	ccccactgsc	1140
actgktacaa	gccccctcac	ttgaggctga	agtgaactaa	agattttaaag	ggccattgac	1200
ttcagtgggtg	attagtggag	agaaaaaaa	aaaaaaaaa	ctcgaag		1247

<210> 33

<211> 977

<212> DNA

<213> Homo sapiens

<400> 33

ggcacgagga	cgggccgagc	gggtcggggc	ttgccgtttg	actggaattg	ccagaatggc	60
ggaccgagcc	ccacgacaac	ctacctccct	gggtctctcg	ccgcagcgct	gcggtctgcc	120
tccctctgct	cctcctcttc	cgccggatcg	cggcgagcgg	atcgaggact	gcctagcgcc	180
cctctgccca	ccggtgggtg	gaggccgctg	cggctgcgctg	ttgagtcgtt	tcctgcccgt	240
tgacctgagc	ctacttcgca	gtagcaggac	cgctgctgtg	gagctgggtc	caggcggtgt	300
gtgccggtcg	cctagtcagg	agaactagtc	ctcgactcac	ggtgagggaa	tggacgacac	360
gggtattgta	ccgctgaggg	aaaggagcgg	gactccggac	ctccaggagg	tagggagtga	420
ggccagtaag	accggcgctc	ctccgggggg	attcctcccc	gggcgttgag	ttgcccacc	480
tgggacccga	ggaagatccc	ggcgtgggtg	tgtgcttttt	gttggtgtta	accctctctg	540
gatttctcga	atttcacacc	actgtccata	tgcgatgatg	tttggttgcc	ccttgacgca	600

cttactcatg gatggtactt cttcagcctc gttagacagc ctggtgatgg aggatgaaga	660
aaccatgtgc ttttcattca gttctggact tagtctccct tttcttccct cagcaagtta	720
tttttgtag ttccttatca aaaagtgtac ataaaaatta ggcaactcca aacatgcctc	780
cagggttaat gtgtgaaata ataagataat atatgtaaag tggaattagc tcctaggcat	840
agggaaagtg cagaatattg ccgtgtgtgc atttacagtt ctggtgatgt cgataacgtt	900
tggtgggtgta attggttagtg ttctgtccct ccaaggagtt aataaaacaa agcaaacata	960
aaaaaaaaa aaaaaaa	977

<210> 34

<211> 1893

<212> DNA

<213> Homo sapiens

<220>

<221> SITE

<222> (1842)

<223> n equals a,t,g, or c

<400> 34

gagcagacgg ctatgggttt gtgtttgtct cttccaaaac tcatgttaaa atgtaattgc	60
cattgtgatg atagtgggag atgggacctt taagaggtgt ctaggtcgtg aaggctctta	120
cttcatgtac ggaccaatgc cattattgtg agagtgggtt ttgccctgtc ttgctctgtt	180
tcatgccctt tctttgcact tctgtcatgc gatgccttcc accacattat gacacagcaa	240
gaaggccctc acaagggtgcc agcaccttga tatttgactt ccagcctaca gaactgtgag	300
tcaataaatt tctgttcatt ataaagtagt caatctcagg tattctgtta cagcagtacg	360
aaattgatta agacagctaa ttatttcaat taggcagaat tcatctgctt atatgcaaga	420
gtagccaggg gttcycctct tatacacata ccttaaagga tgggttatcc tgaaaagcag	480
atatgaagaa ttgtgaaaaa gagctatctc aacagtttagc tgtggagtgt ataatttaaat	540
acaaataaat acaattgtaa acgttcatgt atakttcagt taggkttttg ttttgttttg	600
ttttgttttg tttttttgag atggagtcta gctctgtcac ccaggctgta gtgcagtgc	660
cgcatcttg gctcactgca gcctstgcct cctagggttcg ggcatctctc tgcctcagcc	720
tcccagatgg ctggtattgc ggggtgccac caccacgcct ggctcatttt tgtatttttg	780
gtaggacggg gtttcgccat cttggccggg ctggtcttga acttctgacc tcatgatccg	840
ccgccttg gctcccaaag tgcaggggta caggcgtgag ccactgtgcc tggcccatag	900
ttcagttttt aaaatgcatt cagacttcaa ttccaaaccg aatgactcat ttaaattgtg	960
ttcacatttt gttgaataat atttttaaag cagtaacaac cgtatatcca gttggcctac	1020
atttacttct gaattcaaga agaatagctc ataccataaa tgacaaagag cagcacttac	1080
aattggcaac atactgaagt cactcactct caactaaaag atatgatcat ttctagtata	1140
gctttttact tttctttatt ttctttcttc cttttttctt tctttcgttc cttttttttc	1200
tagaaacatt ttttagcata gtctagccaa tcaactaaat ttgctactca tgttactttg	1260
ctaacattct tcaaaacagt aaatcctact taggtctgat aatttaattt ctgggggtatg	1320
catggtccgc atcttacaat aaaagtataa tttkkattwt taataatgca gaaaagctga	1380
taggaggcta ttgctaattc agagagttcc atcagctaca ggaaaatcat ggccaacttg	1440
cttactcacc tatgtagtca cttagaattg tgggtaaaaa ttaatttgta caattgttta	1500
tttggtgagc tgtatcttta atgctgcata gctccacca agaateccagg tcgtttttgt	1560
ttattatttt attatataac tcctagaaaaa ggggaatcat gattataaaa caaaacgaaa	1620
aacaatctgc ctttatggaa taaaatgatt attctgaaaa tattctataa aacataata	1680
tatatttagt tatttttcaa ataaatgatg tagagcaggg ttttcagcac tctctactaa	1740
tcacatctac tgtggaggct gaggcaggag aattgcttga acccggtgg cagaggttgc	1800
agtgaaccga gatcacgtca ttgcactcca gcctgggtga cnacagtga actccgtctc	1860
aaaaaaaaa aaaaaaaaaa aaaaaaaact cga	1893

<210> 35

<211> 847
 <212> DNA
 <213> Homo sapiens

<220>
 <221> SITE
 <222> (765)
 <223> n equals a,t,g, or c

<220>
 <221> SITE
 <222> (812)
 <223> n equals a,t,g, or c

<400> 35
 gaaatggaga agcttaaggc aaaggggtgac tctccggcct gaccctcact atgaagatat 60
 ttatgtatgt tcccataata ctttttcagat cccaatgtct gtgtttatcc ttgacagaaa 120
 taatagaact gctgtcttct tccaagttaa acaaaacttt ggagctaaaa gctagctagg 180
 tcaagagaaa caaaatctaa tcttcaccca attgagagac tgaatataca aagaatggcc 240
 aaacctaacc atgacctcct acccataaacc tcttcagtaa tcggccagaa gggtcaggac 300
 ttaatcttta accggcagct tccctatttc ttgcccctgc ttccagctct gggccaatca 360
 gagaaagcca aatgtgctcc tcacaccaat cgcgtaagaa gccctgcttc tagtatagcc 420
 ctctccaac ttcccgatgc caacatcttc caatcagagc caacctgaag tttctccctt 480
 ttttcagtat atgtaaagct tttccactcc cctgcctgcc tatgagtctg ttaaatcaag 540
 tgatgggtcg cggctgactc ttgccagctc tgagtaaata gcctattctc atttggtggt 600
 cgttatttct acacagccgg tgggaagggtg gtggagaata gtgaagacta ttcttaaggt 660
 taaaaagaag gttcggctta catttttttt ttatttttaa tttacagcat ggcgtctcca 720
 gcccggttt taaaatacct tctaattgtaa ggaaatagcc ctgcncttg tgcccatcct 780
 agagagaaaa tggtttctta ataaaaaatg tntgaggcaa aaaaaaaaaa aaaaaaaaaa 840
 actcgag 847

<210> 36
 <211> 1100
 <212> DNA
 <213> Homo sapiens

<220>
 <221> SITE
 <222> (1088)
 <223> n equals a,t,g, or c

<400> 36
 ggcacagggtg acatcgccgg gggaggttgt gggcaacggt ggaggaggag agacgggagg 60
 ggaccatttg ggaaggagg gctcttcag agtttttaaa ggcgtttgtg ggggtggagtt 120
 gagtgtgctc tgggcttga cacttgccgt ggtgccctg gctggccgag gagactggct 180
 ctggccagg gcccgtcct gagaggtcct cagcgtctga ctctcgcca ggcgccagca 240
 aggaggggccc ggtccccggg gctaccaggc aggcacgtgc acatcgccat cgccacagc 300
 caactccgcc tgggttttac aaagtcgttg ctttaatgca tgtggacagg aactccctga 360
 ggtcgcccca tgccccctgg ctgtgccagt acggacgccc tggacctgc gaacaggtgg 420
 ggcgggagcagg gggcccaagg gacgggtcc agagacagc gcagggcagg aggggtctca 480
 cggaggggtc tcgactgag gcgccagag ctggtggtcc cgctggacgc catccctctg 540
 cccgggatac acacggccca cgtgtgcccg ccacgcccgc gcccacgcc attgcagtct 600
 gccatcctct ggccgtgacg gtggctgcag cttccccatt tgcgccgttg cctctggctg 660
 tctgcacttt tgttcagct ccaaagaaca tttcataatg ctttcagtac cgacgtacac 720

ttctgaccat	tttgtatgtg	tccttgtgcc	gtagtgacca	ggcctttttt	tggtggatgt	780
gttaccocgc	acacttcaat	ctcaactttg	tgcaccgtcc	atcttctagg	gatagacgcc	840
cagggaaatga	actctagttt	tctaacagat	tagctgagat	attaacttac	tcacacggac	900
aggttgatgc	cagagccgta	agaatgcgcc	agtgcgggtt	tgcgggggac	ttcgggtgtg	960
gggtcctgcg	gccgcgatgg	ccgtggaagg	ttctggggat	ccctgctgcc	acggggacga	1020
gttcggacgc	caggtggacc	tgtgcactca	gtaaaacgca	gtgattcaac	ctggaaaaaa	1080
aaaaaanaa	aaaactcgag					1100

<210> 37

<211> 3145

<212> DNA

<213> Homo sapiens

<220>

<221> SITE

<222> (368)

<223> n equals a,t,g, or c

<220>

<221> SITE

<222> (2402)

<223> n equals a,t,g, or c

<400> 37

ggaggaagaa	gaggaggagg	aggatgaacc	cgcccccgag	gccccagacc	ccaccagga	60
gcgcggcag	aagcctgttg	tgcacccctc	ggcacctgcc	cccctcccta	aggactacgc	120
tttacccttc	ttcgatccca	atgacccggc	gtgccaggag	atcctgtttg	accctcagac	180
caccatcccc	gagctgtttg	ccattgtgcg	ccagtgggtg	ccccaagtcc	agcacaagat	240
agacgtcatc	ggcaatgaga	ttctgcgcgc	aggctgccat	gtgaacgata	gtgacggct	300
gaccgacatg	acactgctcc	actatgcgtg	caaagctggg	gccccagggg	tcggggaccc	360
cgcgccancg	tgcgcctctc	gcagcagctg	ctggcgctgg	gcamgaatgt	gacgctgcgy	420
agcgtctggc	caacatgaac	gcgcttcact	acgcggccta	ttttgatgtg	cccacactcg	480
tgcgtgtgct	gctgaagggt	gcgaggccgc	gagtggtgaa	ctccacgtgc	agtgacttca	540
accacggctc	agccctgcac	atcgctgctt	ccagcctgtg	cctggggcgcc	gcaaatgttt	600
gctggagcac	ggcgccaacc	ctgcgctgag	gaatcgaaaa	ggacaggtgc	cggcggagggt	660
ggctcccagat	cctatggaca	tgtccctgga	caaggcagag	gcggcactgg	tggccaagga	720
gctgcggacg	cttctggaag	aggcagtgcc	actatcttgc	gccctcccca	aggtcacgct	780
acccaactat	gacaacgtcc	caggcaatct	catgcttagc	gcactgggct	tgcgctggg	840
agaccgcgtg	ctgctggatg	gccagaagac	gggcacactg	cggttctgtg	ggaccacgga	900
gtttgccagc	ggcagtggtg	gggcgtggag	ctggacgaac	ctgagggcaa	gaacgatggc	960
agcgttgggg	gcgttcggta	cttcatctgc	cctcccaagc	agggctctct	tgctccgtg	1020
tccaagatct	ccaaggcagt	ggacgcaccc	ccctcctctg	tcacctccac	accccggaac	1080
ccccggatgg	acttctcccg	tgtcaccggc	aaaggccgca	gggaacacaa	aggcaagaag	1140
aagaccccat	catcccatc	tytgggcagc	ttgcagcagc	gtgacggggc	caaggctgag	1200
gttgagacc	aggctccttg	cgcgggccag	aagcagggga	tcgtgcgctt	ctacgggaag	1260
acagactttg	ccccaggtta	ctggtatggc	attgagctgg	accagcccac	aggcaagcat	1320
gatggctctg	tcttcggtgt	ccggtacttc	acttgcccc	cgaggcatgg	ggtcttcgca	1380
ccagcatccc	gtattcagag	gattggcgga	tccactgatt	cccccgggga	cagcgttgga	1440
gccccaaaag	tgcatcaagt	gacaatgacg	cagcccaaac	gcaccttcac	cacagtccgg	1500
accccaaagg	acattgcata	agagaactcc	atttccaggt	tgctgttctg	ctgctggttc	1560
ccctggatgc	tgaggcgga	gatgcagtct	tagaggccct	ggacacctga	caaagagaca	1620
gagtccccac	tagcatctcc	tgacacccga	ggagccctga	gtcaccctga	gatagagatt	1680
cccagtaaca	catccagagt	agagacccct	gttagccagc	cctcgatcat	tgaggcccca	1740
ttattaacag	atactcccat	aataaccccc	aaatacagac	cccatgtcac	ccagaaagag	1800

attccctgag	tagcaccttc	aggctagtc	ctatcccca	cccctcagag	cagattccca	1860
gattaacaga	tttccatata	accccaaatg	atggtgaccc	tctccacata	atgcattaca	1920
acagaacatt	cttgaatcac	ccaaccctgg	atcagaaacc	tccccattaa	caaacactgc	1980
cccttaagtc	ctcttgaaat	aaacataggt	cacaccccca	aagcaaaaaga	gtaacagaca	2040
ttcatgtcat	tgttcccat	ttaacatcag	tctctcaag	atgtcgtgac	cccatgggtca	2100
ccctgaagcc	cttagattcc	aacccctcaa	tcagagactt	ccttcattaa	caaagaccct	2160
tgttcttata	cctcaagaag	aaaccaccca	taaccagccc	actgtcacc	ctaatttaca	2220
gacacaaaa	cagtcctgga	agtgttaatt	acaggacccc	ccaagtcttc	ctaccctctg	2280
caccctcaag	aaacccccag	tgccttgat	gaagcccacc	ccacatggcc	cacagctcct	2340
gtgctggcca	gactccaga	aaattctcta	ttttttaagt	aacgacttcc	ccctttgggg	2400
rncccaaaa	tttgagggcc	ccattctagg	actctgggga	tcccaaaccc	tagagtacac	2460
acgtccaaa	ctccctgtg	ccctcaagtc	ctacagcccc	tagaagaccc	caatgccgta	2520
actcctagga	cccccaaatc	atggaatccc	aaatcccag	ggaatcccaa	atttgaaaat	2580
ccaatcccaa	gtcccagga	aacccaatca	tgaggctcct	gtgcctggta	tggaggagac	2640
tgagtcagg	atatgattc	caggctccca	gacacctcaa	gccctattca	caggcaccag	2700
gaaaccccac	acaggaattc	ccatccctg	aaactggaga	atttcaatgc	cccaggtcca	2760
tgggtttcaa	gacacaaaat	tccaagagcc	ccagccctaa	gggaacccca	aatcctaaag	2820
cctccatctc	taataaatgg	aargcccaa	ggccctgagg	ggatctcaaa	tccctgaacc	2880
ccgatttcaa	tctacgttct	agtcactggc	ctcaaaggac	cccacagcac	ctgggcccaga	2940
ccaacagctc	gagggagAAC	ctgaaggccc	aggggggtcca	gggaggacct	ggggcccaga	3000
ccaccaagga	cagctcacga	ctgccccttc	actgcatgtc	cccaaactca	gcatgactcc	3060
tgtcctcttc	aataaagacg	tttctatggc	aaaaaaaaaa	aaaaaaaaaa	aaaaaaaaaa	3120
aaaaaaaaaa	aaaaaaaaaa	aaagg				3145

<210> 38

<211> 840

<212> DNA

<213> Homo sapiens

<400> 38

ggcacgaggc	cgcaaggct	tctctaggg	ccaccaggct	gaggactcgc	ccaggacatg	60
gactgggtctc	tcagaccct	ggccaccat	gtaggccacc	actccaggcc	gtggacttcc	120
cccaacttgg	ggacagcctt	attcccaaat	gtctctatcc	ttttgactgg	agcatcttct	180
gcacaacctt	gggagcccat	ccaagggttg	gtgaggactg	gtctcccggt	gggtgggggtc	240
tgggggggtac	cctctgggg	tatagattcc	ccactgccc	cagctctgac	tggaccccaa	300
gtggctgcta	tggtaaatta	aatctctccc	cgcgtctcct	ttgcctcatg	tctgctgctc	360
cctgggcagt	ggttgcctcc	tactgaaggg	ctgtggactc	tcggattggc	gttttccctat	420
ggcacttgta	tccctcacgt	gtaggaagca	atagcagcac	cagccttgcc	tctagaagag	480
acattgtcaa	gctactgggg	catggagggtc	atctgcctgc	ctgaccttgg	gggtgggtga	540
gccagtggaa	tgaagggcag	tgtattggca	tcattgcggg	gctgttagcc	ctagcctggg	600
ctcagcctca	gctgaagggg	tcttgggttc	tgtcgtcaca	gccccactg	atgggcagtt	660
gaactgattg	tgtaacctgg	cctcacacgt	ggccagctgc	tttctccagt	catatctggc	720
tccggcttgc	cctgcttttc	tgtctcttag	actcagcact	gaagagaaac	catctttgtc	780
ctcaaaaaaa	aaaaaaaaaa	aaaaaaaaaa	aaaaaaaaaa	aaaaaaaaaa	aaaaaaaaaa	840

<210> 39

<211> 1732

<212> DNA

<213> Homo sapiens

<220>

<221> SITE

<222> (6)

<223> n equals a,t,g, or c

<220>

<221> SITE

<222> (56)

<223> n equals a,t,g, or c

<220>

<221> SITE

<222> (83)

<223> n equals a,t,g, or c

<400> 39

ttaaantgggt	ttttgggggtt	ttattcccaa	aggaaaacct	tcacacaaag	gttacntatt	60
tacctagggtt	caaaattata	ttnttttagga	agggtgccctg	gattttttggt	caggagatttt	120
cttacmaacc	atwttaaaaa	mcamactttg	aaaagtmcc	tactaccaga	agcacttaga	180
ctatcacaga	aaaatacatg	aagagttaat	gtctgagata	agaccagcag	cttatagtta	240
taagaaaaac	acactgtaca	atgtttgggg	gggaaatctg	ttgtttatag	aaaagcttta	300
cctttttctt	atttatectc	agataacgtt	gtgaaatctg	ctcatcagta	aggatgatat	360
ggaaagaatt	ctctttttct	atctatgggt	tatttacata	ttaaaatttt	actcagttta	420
ttaagtacac	atatgatacc	acactagatg	caaatatata	tacaggctat	gacagtcaat	480
gttcaattta	atactgggtca	ataaagtga	tttattcaaa	tgtaacaact	tcacttgaat	540
tttttggtac	acacatatcc	tatataaagc	catctgggtt	ctaattattat	gcacaacaca	600
ttccatttta	attataacat	tcacttaca	gaaaaacaag	gaaaacaatt	caaaactatg	660
ttagtaatta	gttcttggtt	ggtccattgt	acgcaatcag	graagtatat	gacttaatta	720
gttgctatca	aattgaagcm	atgaaaaagg	taaaaatcaa	gtacttaaaa	taaaaatatg	780
gaaataaccc	cccctcccca	atactwwagg	gacacwwaac	aacaactact	gtcccatcaa	840
gcaaaagtgg	aaaacaaaca	gagcatgtgt	gtaacctcac	ttaccatctt	tttgttattc	900
tacttcaacg	ggtcaagaaa	ggtgaagaga	gagaaggtag	aagtaagagt	cagaaaaggc	960
ctaaataaaa	tcctcactga	aatgtttaaa	catacaagca	atagagacaa	ttagggttgag	1020
gtcagatgca	gtactactat	atttaatgga	gttaataatt	agggccaaat	taacatggac	1080
agttattcct	gaatgcaaat	taactcatta	aacgatttaa	atttccattt	ttcagtctac	1140
cacatttttag	ctaacaagac	acaaaaagta	taagtcaaaa	tactaagcac	aggaattcca	1200
aaaagtaagc	ttctttaatt	cattactgaa	cttaagaact	ttaattaga	aaaataaatg	1260
ataacagcaa	aggctctagct	gagtaggcag	agtgttagat	agctcagggt	tgtttttcca	1320
agctctagt	ttcaagttaa	atttatttga	cacagggtatc	ttttgctgkt	ttctactega	1380
agaataattt	aactgatag	gcttacaaga	ttgcacagag	tgaacagaat	tgagccaata	1440
gaatacagaa	tgacaaacca	tacmagcaaa	tgctgctagg	gaattccttg	gcaaaaatgk	1500
tttattagca	tactgacaaa	tttgatttct	atcacttggc	cttaaacacg	aagaaagtgg	1560
gcatgaagag	atgtgtagtt	atgtgtgcct	atgttgga	ataaatttat	ggctatacat	1620
gatatttctt	catacctcaa	aaattttatt	gtccagtata	taaaaaagcc	attctttaaa	1680
aacctgacac	tgaataaaaa	gtatagcctt	caacttcatt	aaaccggcac	ga	1732

<210> 40

<211> 2207

<212> DNA

<213> Homo sapiens

<400> 40

ggtcacgaga	aaccaagggtg	ttggccaggc	catgctctct	ctgaaggctc	tgggggcaga	60
ccccttccag	gccactcttc	tagcgtctga	cgccccagc	agtacttgct	tccatggcgg	120
gtaaggcatc	ccttacatca	tctgcctctg	tcattgcgtg	gccctctctg	ctgtgtcttc	180
tccttttatg	gatgccagtc	attggatttt	gcgcctacca	caccccaatg	tgacctcatc	240
ttacctaatt	atacctgcaa	aatctctgtt	tccaaataag	gtcacattct	gaggtaatgg	300

gggttaggat	ttcaacacgt	cttttttggg	agcacgattc	aacccataac	aatatgtaat	360
atacccaagg	ttaccatct	gggaggtaac	tgtcttcagg	gtaaacttgg	ggaagcatcc	420
cctctctgtg	tgaggcaaga	ttygaattct	tttcatagca	gcaagaggac	agttgaggat	480
tccgtgaggt	ggtttgacca	aaggattttg	tctgtttcat	tttctctgtc	caaagcaatg	540
ccatccacat	aattgcaatt	ttttcagaac	atggattgag	aaatgcttag	aggttcgttt	600
tgtttttata	tgtttttaaa	aaatgttact	tttaacaatg	aaaagagatc	attccttata	660
tattgaccag	tgattagatt	ttaaatctac	ttttctataa	aacacaactt	ttattctttc	720
ctatctttgt	ggaaaacatt	cacatcaaag	tgccacatca	tgggtatgca	tacatgtttg	780
tatatcatag	actatctgga	tgttttttaa	tttcttatcg	agatataatt	catataccat	840
aagtgttgcc	atthttgaag	gtacaattca	gtggctttta	gtatatgtag	aaggttgtac	900
aatcataatt	ccagaatgtt	ttcatcacct	ccagaaaaga	aacctgtac	ccatgaacaa	960
tcagtccctc	tctctcacc	ccattcccta	gcaaccacta	atctgctttc	tgthttctatg	1020
gatttgccta	ttctggacat	ttcatataaa	tggaatcatc	caatatgtgg	cctttttatgt	1080
ctggcttcta	cttagcatca	tatttttgag	gatcatccgt	gctatatcat	ggatgggtac	1140
catgttctct	tgatggctga	ataatacccc	cttgtaggga	tggaccacgt	ttagttttatg	1200
cacacatgag	cttgagtggc	tgctctttaa	agggataaaa	agaaaccagt	tctccctgga	1260
ggggaccagg	gaaatggggc	tcatggatgg	caagagggtt	tttcatcatg	gtgtttttgtg	1320
ttgtttaaat	ttttacattg	tttgtatgtt	acctttaaga	aatgaaggaa	ttgatgttta	1380
caattcagtg	ttcttgtaga	atgcagggtg	tggcattttg	gatatgacca	gggcctctct	1440
acacctctgc	agcttgagat	tctcaccaaa	gcacagcacg	tccctcacc	aggagtaggg	1500
acagcagccc	aggtccccaa	gggctggcca	gtttcagctg	ggagatgggt	gactgggtcc	1560
ttgcaccgcc	cctccgcatt	cacaatccat	gtgtccctcg	gcacagttat	gaatcaytcc	1620
atagtggagc	caytttaggg	tgaggagaaa	ttaggttagc	aaatgctaac	actggaaacg	1680
cactgtctcg	ctggktcctg	cgaggctggc	acagtgtcaa	cagctgtgtg	taggagacgt	1740
gattgcctgg	aacccccaa	aatgcaccta	aaaatcatca	aacaccatca	gattacaaac	1800
caatataaag	aattgatagt	ttttctatgg	aagaatcaaa	aatcactttg	gaaaatagaa	1860
tggaatcaa	aatcctgtaa	tagcaataaa	aactaaagac	ttggcctggc	gtgggtgggtc	1920
atgcctgtaa	accagcact	ttgggaggct	caggtagggg	gacttgaggc	taggagtttg	1980
aaaccaaact	gggtgacata	gtgagaccct	gcctctacaa	aaaaataaaa	aaattaacta	2040
ggcatggtgg	tgcatgcctg	tagtcctagc	tacttgagag	gctgaggcag	gaggattgct	2100
tgagcccagg	acttgagggc	tatagtgagc	catgatcgtg	ccactgcact	gcagactgag	2160
accctgtctc	aaaacaacca	aacaaaaaaa	aaaaaaaaaa	actcgag		2207

<210> 41

<211> 1436

<212> DNA

<213> Homo sapiens

<400> 41

ggcacgagaa	acctacaaga	gacatctctc	tatgccttct	taaaccgagt	ttactccatt	60
tcagcctgtt	ctgaattgg	gactctgtca	ccaataacga	ctgcggagaa	ctgtagcgtg	120
cagatgtgtt	gccccctcct	tttaaaattt	tattttctgt	tttctattgg	gtatttgtht	180
tgthttctgt	actthttctc	tctctccttg	ccccctccc	gccccctccg	ccccatacct	240
tttcttcccc	tggattttca	ccctttgggc	tgcttgtctc	atctttatgc	cccagcacta	300
ggtagcgggc	ccaacacgtg	gtaggcactc	catcagtgtt	tgctgaattg	aaaacattgt	360
tgactgtggc	ttctatcaga	gtgtctacct	tttgcactc	ttccccctcc	tcatttaatt	420
tgctgctttt	aatctacgtg	gtctgagaat	ttgtgaaacc	agtgtgttta	gaagtgtata	480
taatctgaat	caataagctc	tgaatgggtg	ccaagggcct	ctcttatggc	acaaagatgc	540
atggacttca	tgacagctct	tttgggtggc	cagaagccat	tttttataga	atcatggaat	600
ctagaatatt	cctgctggaa	agaacctgag	agttggthtt	gaccaattcc	ctggthttcc	660
agcagatgaa	acaggccaaa	gaggthaaat	gactgggtga	aaatcacata	gctgtctggt	720
gccagagcca	gcctatagta	aagtccttga	acccaagccc	ggtgctcatt	ccactacctc	780
tcacacttca	caacaatttc	ctcaacactt	gaggccagaa	agtctgatct	ctccagaatg	840
atcagcccag	aggaatgctg	agaaatcacc	tggaggaggg	agcagaaaga	gaaggthttt	900

aaggagggct	tctgaatact	tgggagatac	ggaacggacc	aaggaccaca	ctccaggggtg	960
cattcggtgc	tccttggggc	accacttctg	gattacagtg	tgccaggtcc	tttgagggcc	1020
ctacccttc	cccattcatt	gccaccagtg	agaaatgggg	gtgcccctgt	gtaaagaaac	1080
ctaccaagg	tttacatttg	caccttagcc	tcaatagcta	cgaaccctag	agaagcagct	1140
agctggagct	catgtgcaac	tcctgattct	caggagaaag	atggatttta	acccaaaatt	1200
atgagtgagc	tgtaaactct	aaaatgtact	tgggagatag	gccaaagcag	aggtcatggg	1260
ccaactaagt	gttatccagt	agaaaagaca	gtacactgct	tttcttttag	tgtttgcttt	1320
tcctttgcta	tatgttttgc	tatttccttg	tggcttagaa	tgtaaaattg	attgttaaaa	1380
gtttgttct	gaataaatat	ttatcttttg	tattgctaaa	aaaaaaaaaa	aaaaaa	1436

<210> 42

<211> 1419

<212> DNA

<213> Homo sapiens

<400> 42

ctgcagggtca	cagattttat	aggctctgtt	catttttctt	cttttttctt	tctgttcttc	60
agactagata	atgtcagttg	gcttatcctc	aagttttctg	attctcttta	tgcttgttca	120
agtatactgt	ttatatcctt	tagtgaattt	ttttatctca	gttattatac	gtctcaaaaa	180
aaaaaaaaaa	aaaaaaaaag	aaagtagctt	acttggcaat	gcctggacac	tctgctagac	240
agattgcatt	tgaaagggga	gttaagtaga	aaaatttgtt	tcaacgttgt	atcaccaaaa	300
atgtctaagt	atggttcaca	ttcttataaa	agactagaaa	aatttaatac	ttattctaag	360
tgattaaact	gtaatgcaga	actgtaaggg	agaaaaatta	ctacttatcc	aggtactaaa	420
actgtaaatg	aagaacaata	aggaaattca	ttaaactgtt	acttctttat	gcacaattca	480
tggttaggcc	tatgatgaga	atatgcaatg	agcagatgtg	cctggtcctt	gttgtgtctt	540
gtgctgctgt	acaacctgtt	tcaattttat	atgcataaac	tctctgtgta	gctcgaattt	600
tttttaagac	taatttagag	ctacagtttt	cttaaacaat	catgcagtta	aactattttc	660
aaatgatggc	agagatcctg	gtggcataaa	gtaatgctgc	tgttctctgc	tgtaaaatat	720
ggtgagacat	ttttaattgt	atttttaaaa	ataattacaa	ttgtatat	ttttaattt	780
tggatccaac	tttttttcaa	gcctcttgaa	gcactaagt	ggatccctgg	ggttaacaat	840
ttataaagca	ttaatatata	gaaggatata	aaagcacatg	aagacatatt	gatatgggaa	900
catatataca	tattacaagt	gtttgaagca	ttgtcatagc	ctttgttaga	gagtagaatt	960
taacataaag	tgggttctcg	aaggagccca	cttgtgagga	tttgaggact	aaaataggat	1020
ctaaaaaggt	tatttccagc	cgggttttat	ggcctataca	cagcaagcct	ccagcatgct	1080
gttccacaac	actagatgag	taatacatga	gatgatatac	agggccacga	gatctctaag	1140
aatgtggctt	ttacattttt	gtatagtggg	aaagaaatat	atgttcaacta	gaaagaacaa	1200
aaagacaaag	aagaaaataa	aacatctcat	ccccaaaaat	aactgcta	aatttggtat	1260
gtgtctttcc	aatgtttctg	agcatatagg	agtggtatat	ttacacacac	acacaaacac	1320
acacaaacac	aaagcaccca	acttacactt	aggagatttc	aacttaactt	gaccgctctg	1380
accaaaaaaa	aaaaaaaaaa	aaaaaaaaaa	aaaaaaaaaa			1419

<210> 43

<211> 2396

<212> DNA

<213> Homo sapiens

<400> 43

gggtaagttg	gcataaattt	tccatgaaac	acttgactgc	catttgcagc	atcatatcta	60
ctttttatgc	acctaattat	atttttctct	tcacttttat	gtwyttttct	aaaactgcc	120
cctaaatacc	tttccactcc	gttaaatgta	tggagtcaag	ataaatttct	cattcttget	180
ctacagttta	agatgtataa	gaagtaaaca	cttccgtgga	atagggctta	atttggggct	240
caaatgtttt	gttttctctc	tgagatgttt	ttgaatggrr	aagactactc	tggttttgga	300
tgcagcaact	tttctcctgg	catggaagaa	ctaaggactt	tgataaacag	tttcccatga	360

tagtataata	taatgagggg	cccagtcctgc	tcttgtgaaa	atggctgagc	ggcagcagcc	420
catcccagct	gtgtgtgagt	agccccgctg	ttaaaagtac	acctgctggg	aattgttggc	480
acatgctcat	agcatatgct	ctgtggggga	ggtgctttta	attccaagag	gagcttttct	540
cccacatctg	cggatgcaga	acaaagatgc	agcaggactt	tagaaaacag	gagtgtttct	600
actgacattt	tattcattgc	tttgttagtc	ccgcaaatgg	agcaacttct	ggtgttttgt	660
aagcagaaaa	acaaaacatt	caaaacccaa	aacctaatta	gtacttttgt	cctttatata	720
cctaggatgc	tctagggagt	agaggtggca	ggcatcttta	tcttcaccca	ctggcacatt	780
agcagcttat	gcaacttagg	aagtcactgg	gagatgagtc	ctaaaagcct	gaatttcagg	840
atggggagga	gataagaagc	agagaacttt	gggagaccat	ggggcattta	gtcctggcat	900
agacacagaa	tggatattca	caggaagatt	cctgaggcca	gatatgcagg	agagcacaca	960
ttgtcacaag	gggtgtggtg	ggaagagagg	tgctggatgc	ctaagacaga	cttctgtgct	1020
tggcagattg	actgagtgtt	gactctgaat	actttgtggt	ccagtcacata	ataggccagt	1080
cactcctgcc	tgatgcatcc	ctgttgcaat	gttcataaaa	cctgtatcat	atgggggaagc	1140
ttcaaggctg	gcaggctttt	catcaagtaa	tcattcattcc	atgttcctac	gtatccttga	1200
aatcaaaaacc	agccaatatg	tgtgtttttg	gttttgtttt	acawaaaggg	catwcaatga	1260
gacgaattct	taagataatg	casatacaga	atcatcaata	aagtttcaaa	gagtttatga	1320
agaaaaggta	ttttcctttc	ttgtaaaaaa	tggtatatat	atataatatt	ctgatctgca	1380
ttcaccttag	ggcttagctc	ttattttatg	aattttcagg	ctgtgatttg	tggtggatgg	1440
ctaagctcta	aaataatgca	aatagcccca	atctttaaat	atagcgggtg	taagttgaac	1500
aagtaacaca	atacagaaaa	tgctgatatg	aatacttagt	aataatacaa	aagttcctgc	1560
taaattatat	atgtgtatca	ctgcctgact	agaaacccca	cttttctttt	ctaattccagc	1620
acaaaatcaa	actctgattc	tacaaccagt	ctttttaaag	ggcaaaaatg	actcaacacc	1680
tttgttttgt	atggaaaact	ttttttttta	cactaactgc	aaaactgctt	taaaaaaagg	1740
cttattcagg	atagataagc	atgtactcct	tttttaactt	gctggaagac	ttgcccctcc	1800
aaactagcac	ccccaaaaga	caacttcctt	cagaaacggg	gtgttttacc	taaacatagt	1860
agcttacatg	ttagccagca	gtaggtcggc	actagtgttt	tccacgggta	tcacctttga	1920
cagggtgatgt	gcatctatag	atagtgggaag	ccaccccatg	aggaggtggt	aatagcagca	1980
tgggttctact	tttggttaatc	aggtaatcat	gtgtatatat	ttagattcgc	attattttta	2040
catttctctg	ctactctgca	cttcagggttc	gttaagctat	tttaataatt	actgggggta	2100
tggcaaacac	caatggaaat	gtatatggca	actgctttcc	tgagcaagtg	tgatttgttt	2160
tatggctggt	caagttataa	aattgttctt	acattgtagg	taaacaaaaat	cttgatgttt	2220
ttaaagggtca	ctgtaactta	aggttcaaat	ttctggcaca	gttttattag	tattcacttc	2280
ggaagctaat	aagataccat	ggtttttctat	gttactccca	ttgtaacat	agtaaaagtga	2340
ctttcaataa	aagattttatg	ttatttttgaa	aaaaaaaaaa	aaaaaaaaaa	actcga	2396

<210> 44

<211> 1984

<212> DNA

<213> Homo sapiens

<400> 44

tttttttttt	tttttttttt	tttttttttt	tttttttttt	tttttttttt	ttttttgcca	60
tattgaatca	atatttat	caggacatgc	catgtcaaaa	taaaacaaaag	agtcaaccct	120
tgcttttaac	aattatattg	tattataaaa	gcactttaca	actccatccc	atctttaagt	180
tactggtatg	tgggctaattg	attatctgta	agcattttctc	tattcagatc	cataatccaa	240
gtgctctctg	aattattacaa	ggtgacaata	agtgggaagt	ggaggaggaa	gaggaaagag	300
aggggactaa	ggttctccca	gtttaagggt	ttgttccaat	gaggggatga	ggaagtatga	360
agatattttt	gttgtctttt	catctttata	ctgtgttaag	taatgcttac	aacataaatt	420
cagcaggctt	ttcctgagct	gtaactcaga	agttttcgtc	ctgcaacaa	tgtatttaca	480
aatgtgttta	ttagcttaca	cagcaatctc	acaataacta	gtaaagatta	aaagggcaca	540
ctcctagtat	atgtgtttgc	agtgttttaa	gggaaataca	tattgccatg	gtgaagctct	600
aaatagattc	aacgaaacat	ctgaagattg	aaagtgtgta	aaaaaaaaaa	aaggcacaaga	660
aataaatatc	accaaagaaa	aaagattaat	tgtagcttaa	atataaaact	aatcaccag	720
ttaattaaac	tatacagatc	taatacaaac	caaaaccagc	ctgaaagacc	caaccttaaa	780

aaatgctaaa	aaataaggca	taaatctgca	taatataggag	ttttatttcc	attctctcct	840
ttccctctac	tatgtatgct	ttacctgac	tgcccttagg	ggcttacaag	aaaacgggtt	900
ccggtttccg	tcttcaat	gacctcaaat	gtccctgagca	aagtttttgt	tattctgctg	960
agggttcttt	tggttgtaag	ctttagacat	cattatttct	actaattcag	tagttttgat	1020
cgtagcgcca	aatttaaatt	cttccactgg	ttcttctgga	aggaattaaa	actttttacag	1080
tgccctgcct	gcacagtatt	atgtttttta	aaaagaaaac	ccaagcaaaa	tctattgctt	1140
aaagagggtt	cttatttttt	aaacaaacag	aataactctt	gacaatttta	aaaccttggg	1200
agaaacagtt	cattagaact	tcatttttct	accgtgaaga	attaaatact	aaaaacctgt	1260
tctgaagcac	ttggttaatt	ttctctccca	gagtctaata	aagcacatgt	gaaagagcca	1320
tttgttttag	tcagaaatac	atcttatgtt	cttctacttc	taagtactca	gtatgttctt	1380
taggactcat	tttgaagatg	caccaggagc	cttttctcat	tcaagcactg	cctaccatga	1440
tggctgaatt	ttgacctcaa	agaagaccgt	aattttttatc	agtgtctcaag	aataattttg	1500
gacatcttgg	tccgtagcct	acagcaagtg	gtatctgtaa	aattaaagga	taattccaat	1560
gggcttggtg	gaactgctgc	tttgccatct	cttgtttgtt	ttgaggaagt	cgggggaggc	1620
taggtaagaa	tagggaataa	gggaatgggg	gtaagtgaga	ggtgagaaaa	gcaaggagag	1680
ataaagtagg	ctgtgaacat	accgctcggt	aaccaagcca	tactcatact	gttgagattt	1740
ccatcatttt	gaagtacatt	atcataacat	taaaaaagaa	aaaaatgtta	agaaaatgta	1800
tctaattttt	aaagttatca	ctggaatatg	ctgaaatatt	ttggcttttt	gtaaaatata	1860
aataatgaag	acactgactt	ttgttgtgtt	tgtgaagcta	atagatcctc	tccacgagac	1920
aggcagcaat	gatgaattgc	aatacgttat	tactgaaggg	aaaagggttca	agccaatatt	1980
caca						1984

<210> 45

<211> 1953

<212> DNA

<213> Homo sapiens

<400> 45

tgaagaatct	caaaaagccc	acccaacttt	cagctgacat	ttccaccagc	cctctcatat	60
ttgttaacaa	ttgggtatctt	tgagtattta	ccaaagagct	gccaagggtta	cagtgaacag	120
agttttgaaa	ggcattgctt	taaaggaaaa	aagtataggt	atgtgtacat	atataatata	180
tacaaacaca	tgtacttctg	tatacat	catattttta	caattcatat	tttaatttct	240
aggctataac	tcagaccaaa	ttatacctaa	aagttccaac	aaagtccctt	tttcaatctc	300
acattaccaa	aaagatggct	gcaaagttaa	tttggacctt	tcattaaatt	tgttttcaaa	360
actagaataa	ttccaccaca	gaatcagaat	tttctaccgt	tccacaccca	accttctcaa	420
atacacacaa	ccttggttact	tttcaactcca	gcaccttcat	acgtttttct	ccaggaggag	480
gttcttgcag	ctggaaacag	cctattttgt	ggtcactgtc	aagtggatgg	atattctagc	540
gctcccaaaa	aagcactatg	gccttatatg	caggggaaggc	acataccacc	aagttcaatg	600
agaaatatta	gagctaaccg	tactctcttc	tctgcgtacg	ttcgagtata	cgttgcccat	660
atccctccca	tattttcttt	ttgctgcttt	tgctctggaa	ctttgctttt	agcagggaaa	720
gcagctgtcc	cctgagtgtt	ttgaattggg	aatataccca	gtgtgtgttc	tccccctct	780
tacgaggcta	cataacacat	ctatgatgtt	gctttaagtt	tttagaggct	atacctcaaa	840
gtagctgcgg	attttgtctc	ctgcactgcc	aatatgcaac	tgatcccgct	tttattaatt	900
ttttgaagaa	gtacacagaa	tttttacaga	atgtagtatt	ttgatatcat	taagtaaacc	960
aatcagaaaa	ctccttgagc	aatagttgtt	tctttgtcag	tttcagttac	aatcatcttt	1020
acccattaag	acttacatta	acattccttt	tatataaaga	gttggtatatg	tccacctaaa	1080
ttcctatgtc	cacacttaac	ctttaaagat	gtacattgag	ggaatatcaa	aaaatagcgt	1140
tcattggtac	gaatattgtg	aattgttaaaa	gcacagcaaa	ctgcactgca	cttataaagc	1200
aaatctatta	gaaaaaaaagc	atttttctaa	atgctcaaat	gttatcaaaa	tactatat	1260
atagaagtaa	tcttctctgt	taacagccaa	gatttctgtg	tagaaataac	tcttgtgagt	1320
tttatattgk	gcttttttggg	gggttctaata	atctcagcag	tagcgtctta	aacagcagta	1380
ttactataag	cagttgcttc	aaaatgtgaa	tttaactgtt	gaaactgtgg	ctttaacatc	1440
catgtgacta	gtgtatatgg	tatttgcctc	ccattagcaa	aataattcat	tgtaggtaa	1500
acttcactag	tgcaaatgtc	agatctgtga	gcaatgtttc	ctattgaata	taaactactg	1560

tctaaaatat	acatatcagc	agcccagcct	ttatcaggaa	aattatactt	ggcaagttgc	1620
tgaaaatgca	caaagttatg	aaagttaaag	gtatgctgca	aataactagc	cattattcta	1680
tgtattatta	aataattact	agttctgtta	aaagcagagc	agaagttaga	cactaaggat	1740
ctctttgtga	actctgtgtt	ctctatatta	gattgctgtt	tatatgtaag	aattttattg	1800
cttatgtggc	atacaatatt	tataactata	aactttatag	aagtacagta	ttaaagtcag	1860
tggtacacag	acattctgta	catatcctgt	gaaacgtgct	gtcatatgaa	ataaatatat	1920
ctgtctttaa	aaaaaaaaaa	aaaaaaactc	gag			1953

<210> 46

<211> 1052

<212> DNA

<213> Homo sapiens

<400> 46

gctgccgtgt	gcacggccgt	ctgggtctctc	tcccacacgt	gtgcgcaacc	tgtcatggag	60
atgtgagggc	cttgtgtgtg	cttctgtgtg	tgactgtgtg	actgcgggtc	cagacccccg	120
cctggcgggtg	atgtgggccc	ttaaatcact	cttctgtctc	acccccctcc	cagtgattcg	180
gttttacttt	gcagcactgt	ggatccgggc	agctgggcgg	cttctcgggg	gtggggggtc	240
ccccaccccc	cccacaagtc	tggtcccagg	gttctcggag	gcaggggggtc	tctgttagtg	300
cgctccctcc	agctgcaggc	acatagcccc	agctcacagc	tcgcctgagt	cgacgcgggc	360
tggggtgaaa	gctccaagtg	ggcctctggc	cttcccgtct	ctctgggtcc	agagtgtctg	420
gagcatgtgg	cacagaccag	ggccccctgt	cctccgagga	gggtgggaca	tcctctctgt	480
ctcacgcccc	tgggtggaga	ttctggctgg	cctcctctcc	ctgtttgcc	aggtcaaagt	540
gggccaagg	tgcaagtgct	tagcctgggt	ccctctcccg	ggccccgagg	ttctgtgggt	600
tgggcagatt	ggagacagga	ctcgtgtaag	ggctctgtct	gggtgaagga	tggagacaga	660
gaaaatcaag	atcctttcac	aagttaattc	tacgtctgct	gagccccagc	ccccgacaca	720
tcaccctgag	gaggtgctag	gcttctctgg	gccccctgtg	ccccatccac	atgttgacaga	780
gtaaactctg	ccccttgagc	ctgggggtccg	agatggacgc	ctggctgccc	ctcctggact	840
gcgggtgaca	gctggcgaga	cactgcgggg	cttgggtgcg	gggagatgga	gtggggctga	900
gctgcatttt	tccagccacc	ccacatccca	cagaagggga	gtcatggtca	gtgccttgag	960
ctggaaagac	gggcaatgct	tccggccccc	accaaccaag	aaaaccacca	ggggctcatt	1020
catcctctca	aaaaaaaaaa	aaaaaaactc	ga			1052

<210> 47

<211> 1381

<212> DNA

<213> Homo sapiens

<400> 47

gaggggtgaag	tgtagctgct	ccctcctctc	cccgttctct	tctgtccccc	catcagggga	60
atacctactc	attcttcaag	ccccggttca	aagtcgcctc	ctcctgatgc	ccacctgctg	120
caggaattga	acctccactt	tgcatctata	gtgcttcaga	ccagccttga	ggagaatgtc	180
actgtgggtg	ggtgtagcca	tctatacttg	tcctcctcct	acaaggctgt	gagccacttg	240
aaggcttctg	atgaagcagg	aacaaaaata	ttaagatgtg	aataattaca	gcctttcttc	300
tcctgccatg	ttcccacatc	tgagctgac	aaggtgatac	aaagtgggaag	atgcttcaact	360
gaagaaagag	cataaagtta	gcagcaaaaag	ctggatcaga	atagaatcga	tccagtcaac	420
tagaggtgga	ggcaagagag	agcagagcta	gagttttagg	agagaaaaag	gaaaaagaaa	480
aaaggggggtg	gggagtgtag	tgacctatgg	cactttgggg	aaaggcaggc	ctgaagagga	540
tttgagagaga	gtttctgtct	gttgtagcac	atagttgctt	ttcccacctt	gccgcttggt	600
tgggagtttt	gcaaaggctt	cattggccaat	caatgctttt	ggagactatc	agggtttctt	660
ctggatgtga	cccaggcttc	aaatggccta	agatttgctt	aggttgcaat	gtacattttg	720
ggcccagctc	agcaggtatg	cagaataata	caaagatgag	acaaaagagc	aggtggaggg	780
acaggaggag	gagtgacac	aggctgtggc	tctctgtccc	tctctgggca	gaccaccatg	840

gtggacaatg	ctcatgcggt	catgacaact	tggtccccag	ggagccccag	atthttgatga	900
accatctccc	cagggtcccgc	ctctcccaaa	ttgtggataa	ataaagccct	aagatthttta	960
ataggcacia	ccattgagga	tgtatcagaa	atactccctc	tctgtgggtt	tggtgcagac	1020
aggaggggat	gggtgttggg	tgtgattctc	tttcagcctt	tctttcttcc	cctctcaaga	1080
aagcttatca	ggaagcacac	aagcacaata	cttctagata	aatattactt	catttgactc	1140
cttctaggaa	tcttgtgagg	tggtattaat	gttcttattt	taatttcaac	ttccaattgt	1200
ttattatcag	catgtagaga	tataatttat	ttttgaatat	tgacctatta	tgaaatthtta	1260
ctgaacttaa	tacttctagt	tttatttttg	attgattctt	agagtthttct	atgaagacaa	1320
tcatgttgte	tgtgaataaa	gatggtttct	atthtttctt	tccaaaaaaa	aaaaaaaaaa	1380
a						1381

<210> 48

<211> 992

<212> DNA

<213> Homo sapiens

<220>

<221> SITE

<222> (405)

<223> n equals a,t,g, or c

<400> 48

ggttccagcc	caccccatcc	cacacccacc	tggtctcatgt	ctctctgacc	tcagttcccg	60
cctgcacccc	cagggatggg	tgggaagggt	ctagaaggta	tggacctgga	gtgggtcact	120
tctgggtggg	aatctggcta	gtcagaagca	tgagtgatcg	aatgaacaag	aatgctctgt	180
aaactctgag	acctttggag	gctgcagagg	gagaggggag	cagaactcca	gagatctgca	240
tcttgtcaga	ggagtccatc	aagctgattt	ggggagaagt	ccccgcactc	ccccagctcc	300
ccggctctcc	cagcagctca	tctacctgcc	cccacctggg	ggcacctgca	aagctgttac	360
catggactcg	gatttggtatt	tcttcgactc	ctacagcatc	actgncctgc	gcctcgactg	420
tgagacgcgc	tactgggtgga	gaactgcaac	tgcgcgatgg	tgcacatgcc	aggtcaggcy	480
kggggytccg	agcatactcc	tggggctcctg	ggsctttgct	gccyttcact	agctcccat	540
ccatatcaat	ctcccttccc	atcttctccc	agcttacacc	ttctaggcct	ttggtactac	600
caccatcacc	agacccttg	aattcccata	ctgcatcatt	gtcttttctc	ctctgtaggg	660
gatgccccat	actgtactcc	agagcagtac	aaggagtgtg	cagatcctgc	tctgggtgag	720
cgcccttggc	ctggggcagt	ctgggggagg	gaaaagggtg	tgccagccac	gtgcgcagga	780
gtttcagggt	acgctcccag	aagcctcaca	taactcctgg	actgggctgc	acctctctt	840
gtgtctgaca	ttaaagctga	gaatgggtcag	tcatgggtgg	ttacacctgt	aatcccagca	900
cttggggagg	ctgaggtggg	caaattgctt	gagcccagga	gttcgagacc	agcctgggtt	960
catggtgaaa	ccctgactct	atgaaaaaaaa	aa			992

<210> 49

<211> 2129

<212> DNA

<213> Homo sapiens

<220>

<221> SITE

<222> (2097)

<223> n equals a,t,g, or c

<220>

<221> SITE

<222> (2120)

<223> n equals a,t,g, or c

<220>

<221> SITE

<222> (2125)

<223> n equals a,t,g, or c

<400> 49

ttggcccatc	ccatcagtg	gcatctggct	gacgccattc	actggacggt	ccctgaacac	60
ctaggaatgc	acacaccgtg	cttctcagac	actggagacg	caaaggcagg	aggatgcagt	120
ccggtgagag	gacacgatct	ttacctgcac	aatcagactg	taagcccagc	agagaacccc	180
aggggcgcct	gggtacttct	cggaggtcat	cttagttgtg	gtggggaaga	caaagaaata	240
agcaaacaag	aaactagagt	tactatacaa	gaaactctcc	tgagtttgta	aaccttaagc	300
ataaggattc	agttgacctt	tttcttggtt	catcaatctg	gaaagaactt	acataaagcg	360
ccattgacac	gttcacctgg	gagctccatg	ggccgtaagt	ctttgacagc	caatttaatt	420
tgaggtcaga	gggccttgag	gtacacagtc	agcactgttt	gaacactttt	cctgaaagca	480
aaactcacag	ctccctgcgc	cctctgacaa	cactagctat	ttctgccaga	gtaagaactt	540
ctattactat	tttattattg	ttcatatgtc	ttttgatgat	ggttgtgtga	cagggggaag	600
caggatctat	ttggtttctt	ccccctcccc	ccaccccttc	ctttttgtct	ctcttttttt	660
ttctctaaga	aaatcaccag	actagttttt	ccatcttgag	taattttctt	tgtgggacag	720
ttttgatcct	cattttgaaa	gcatgcgtgt	gcacatgtgt	gttgccctgtg	gtgccaggtg	780
agacaggttg	cactaaactcc	agctgcttgg	aaggcatccc	aagggcgcat	cttaaagtgtg	840
gagcagacct	ccctttttcca	gcccctgggg	ccattagacc	acgtgctgga	actagcattg	900
taaaattccc	atcccagttc	cactccccctg	aagtgaacc	cttttttttt	tgtgacagta	960
aatcttaaaa	atcattgtct	ctttatgaac	atttcctcag	tttcttctct	gctgaaaatg	1020
taagccatgc	tactttttaa	tgtattttga	attttgtgct	cattggaaat	tgatatgcta	1080
atgcctcccc	cacccccccgc	cagacttttc	ttttataact	ttgtctgtgt	tttactgggg	1140
taggctgggc	atgcgtgcgt	gcctttaggg	cagcatttta	aacctttgcc	aaaattgcaa	1200
atgggacatg	tacattcttc	tgtctccatcc	tacttaaaaca	cctatcagct	atttttatct	1260
ttaacctttt	ctgtatgttt	gaagtgtgtg	gggggtgtgt	gtgtgtgtga	aagagcgaga	1320
gaatgatgtc	atctaaagtt	ttttgaagaa	ttatttggtt	ttcattgcat	taaaattcta	1380
tactcccag	ctttgttttc	atttaaaaaa	atatacaaag	agctttgtaa	atacaacaca	1440
ttttatttct	cccccttctt	ttaatgtaca	gcttttttgc	cacttatata	tacttaaaat	1500
attcccattg	attatgtcca	gttcttcttg	gaaaaaaatt	tggttttgaa	tgaacctgca	1560
aagcatcctg	cagcgtgagc	agctcctcca	cctggagctc	cgaagcatct	tctcaggcca	1620
aagcggcatt	acccgtgaat	ctgtcttctc	cgccacagca	tggtttgagg	cgcagctctgt	1680
taatatagct	gggccatgtc	agtgactgtt	gtgtttgttg	ggtcaggtgg	ggggcatggt	1740
atttgcaaaa	aaaacaaatt	atggctaatt	tattattttg	ttgcagtggg	gttaactgta	1800
aactcatgta	agagtctgtg	atttcctcat	tggttgatct	ctctctctgt	aatcctcatt	1860
gcaaattttc	accaggacag	cgttttttga	ttagagggga	gctctggcac	agtatgcttt	1920
aatttagcag	gaacttccag	atgatttaaa	ttctcgatgc	tgtgatgaca	cacatatgat	1980
ctttcgtgtt	tctgagcgac	tctactttca	ttgtttgcca	gcgtggctcg	ttgctgttgc	2040
ccaataaagc	ttgtgtacgt	tcaaaaaaaaa	aaaaaaaaaac	ccgggggggg	ccccganccc	2100
cattggccct	aaaggggggn	gtttnaaaa				2129

<210> 50

<211> 1474

<212> DNA

<213> Homo sapiens

<400> 50

ggcacgagca	gtgtctgaca	tttcaatgtt	aaatttgtca	ccgttaagga	accaatgttg	60
gtacattact	actaactaaa	attcatttat	ttggattttg	ttagttttta	cctaaggttc	120
ttttttgggt	ccaggatcct	atccagaata	cgacaatata	tttagttgac	atcttccctt	180

agcctatctg	ggctgtgatg	gcatttcaga	ctttccttgt	ttttgatgac	tttgatgggt	240
ttgaggagta	ttgggtcaggg	gattttgtaga	atgtccctca	gtttgggttt	gtctgatgtt	300
ttcttcatgg	ttagactggg	attatgggtt	tgggggagga	gagtcacaga	gataaagtgc	360
cgctctcatt	gcatacatc	aaggggaaat	actatcagta	cggcttatca	ctgatattgt	420
taacctcggt	caccaggctg	agctagtgtt	tgccaggttt	ctttactgta	ttgttttccc	480
gtactcgtct	ttttggaaac	aagtcattaa	ttgtagacca	cactcagtag	gtctgttttc	540
ttaaagggga	gctgtctata	aaattatattg	gaattcttct	gctatggata	gtcttttttt	600
cttctccgtt	tatatattta	ttcaatcatt	tatttatata	tgtatggact	catagatggt	660
tattttatac	ttgggggtcat	aatcctatac	tactttatatt	attttgttgc	tcaaattttt	720
ctaccaacag	gacatactcc	attgttttgt	tttttgagta	cttcccttact	ttctggcact	780
agaagaagtc	cagactcatc	ttgtatatatt	cctgccccag	ccctagaatc	agccattttt	840
ccaaggatcc	ctggttctat	tttttggtga	atcatgttag	aaaccaagat	ctggttactc	900
agtgtgctca	ttgctgttgg	aatatcatga	cttctaggcc	ttttcagtgt	atagacctgg	960
aaaatatatg	tatgtatata	catatataac	tattatctgt	atctatattg	aggtaaacat	1020
gagttcatat	tgatgacttt	gactatcagt	actatatagt	tcattctggc	tttactctgc	1080
tatctattac	ctccctctcc	aaaagtgaga	aatctggctc	acgccatctg	acatctatta	1140
acttatttgt	tcagtcctcat	cgtccatgta	cagcactttt	agaattgtta	acctgtatct	1200
ccatgagaaa	tgcttttacc	aaaaaagaac	agtgcctata	tacagttctt	tttgtgttta	1260
gagtttccag	tttcaagttt	cccaaagtta	cttaaataca	accttttccc	ctgacatctg	1320
tatttttaatt	ttattttttaa	taaattgatc	taaattacat	ggtaattata	tataaagctt	1380
ataaaaaaatt	tcccttgccc	aactaaagat	cttttagcttt	agatgtgttt	tttttttcc	1440
cctgactttt	gtgaccaaaa	aaaaaaaaaa	aaaa			1474

<210> 51

<211> 2100

<212> DNA

<213> Homo sapiens

<400> 51

ggcacgaggt	gaaattgggtg	tgtacatttt	gggcaaagca	aagatacaaa	tgttattcat	60
gcggtgcata	agtcctaaag	gataccagta	tccttctagc	attaactgat	aaaattaact	120
ctccaaagac	caaaagacca	gctctcactt	tctccttcaa	aacatcttac	actgaaatgc	180
agatttatatt	gacagcaact	tcaattttca	ttcactcaca	gcatgtgaaa	ttattttgct	240
gaacctgtgt	tacaacttat	ttactggcat	cttaataatc	tattagcaga	aattgggtgt	300
ctctgtgtgt	agtaactgtt	ctactttaa	gagctggctc	agcctaggct	cctgaccaga	360
tctggctgac	aagctagttt	tcaaaaccag	tgaagtgtgt	tctgcaggga	gaactgtgct	420
tgcttttgat	aattgcctca	gttttatgaa	atcaatcagc	ctgaagggaa	gacctcatta	480
cagatctgac	aagatagact	ccaatttcaa	tactagaaca	actctctgtc	ctcaggctct	540
gaaatctgca	attattttatt	ttatttagac	agagacaacc	ataatagagt	acctctgggc	600
acattctcag	ggaagtactc	aacagataac	ctaattgtca	gtgtgctagg	aacctgccct	660
tttctgttca	ctcagggtgga	aaattgggtca	acaggaatta	cagcatggcc	tagagccagc	720
tcaaagcaga	cactgatgaa	tcaccatctt	tccaatcccc	taccccaaaa	gatgtggaga	780
gaaagggtag	ctgctgtaac	tgttcccagg	gctaaactac	aacaaaaata	ggcttttctct	840
tctcagttcc	tctgaagact	gccagacaga	ctttattcta	aggaaaaatga	tagatcaccc	900
atctaccatc	tgtcactttg	aactgttcta	tcaaatacaac	atttttttga	actttgtttt	960
caatgaggaa	tgtgattttg	ctgtgtttcc	caagccttaa	attggacaag	ggaaaaaagt	1020
cctgaagtgt	tcttcttctt	gaagttggaa	atatgtattt	tgcttttctg	ctcacggaac	1080
acaaacacct	taacagattt	agttcagact	tttctcgaat	aattcacccct	ttaccaggcc	1140
tgggagatgt	cagccccaca	gaaaaaacac	agagaacatt	atgagtgtac	gaaaaatatgg	1200
atttagagag	aaatgatact	ctctctagtc	cagctgtagt	atcactgcta	gcagtgttct	1260
gatagtacag	caccagatg	gaacagagca	ttcctgaaag	atttgggagg	agacatttcc	1320
attgttaggg	gaaaatgtga	ttctttcatg	ttggacagtt	caaggtagtg	atgaatatgg	1380
tctcctaaac	agaatccaag	caaagagcaa	tcactgacta	ctcccgcat	atttattcat	1440
gatttacttt	aatggctttg	tgttattaag	aaaaagtagg	ttagaagagg	aatgcaaagc	1500

taacatcttc	caagctgtaa	atagactgta	gcctggagggc	cgaaggagcc	acactactca	1560
aatgttctta	ttccattttt	gagacaaatt	tatataaacc	ccccattggg	ttagaagggc	1620
attgacattt	gtttagtgtc	caacatgtgc	tacaccttaa	acatacaatc	tcttatttaa	1680
ttttcctcac	tttgctgtaa	ggggggcatt	atcatcccca	ttttacaatg	cagaaacagt	1740
ctgagagaga	tcaagtgacc	tgttttatac	accttgtaag	tggcacactt	agtaagtggc	1800
acacctagta	agttgcagat	gtggctttga	atgcagatct	gactctagag	cagcactgtc	1860
cagcagaagt	ttctgggatg	attgaaatgt	tatataaata	tgctgtccaa	aacagcagcc	1920
actagctaca	tgtaactatt	gagcacttga	aatgtgatta	gtgcaactga	gcaactgaac	1980
ttttcacttt	ttttctttct	ttctgggttt	gttttttgtt	ttttgttttt	ttagatggag	2040
cattgcactc	cagcctgggt	gacagagcaa	aaccctgtt	tcaaaaaaaaa	aaaaaaaaaa	2100

<210> 52

<211> 1607

<212> DNA

<213> Homo sapiens

<400> 52

ggcagcaggt	caccctaaca	ccagattact	ttccccaaga	catgccagga	tcattttcttc	60
tagcccagct	gtgttcgaac	tagcccttca	ccaagtgggt	cagtttcctt	ggtggggcct	120
gtatgattgt	gaggggctgg	gggaagtctg	gaggcgggtg	tcagggaagt	atgaagccca	180
gttcctgcca	caaggttcac	acccaatgaa	tggtcactgt	tattacacag	gatggtaggc	240
ttgtggtttg	ttttttccct	ctattctttg	gtcacttttc	tataagcata	ttatttttag	300
aatcagaaaa	aaaaactcat	gaaacaaaag	aaatgtgatg	agtgatattac	aggagacaaa	360
gggcagagag	agagagacag	aaagagagag	agagagacag	atgctcttta	gaaaccgtgg	420
ctctgggccc	cgaatttaa	tctgcagtta	aggaggctga	gagaggcagg	tgggtgcacc	480
cacaggctctc	ctaactgaga	acttcctgcc	tttgagccca	cctactagtc	cccaggttac	540
agatctcagg	gaaacccccc	ggctcccacc	ccctccacac	acacacccac	tcccagtcag	600
gcggcctgtg	agcctggggt	ggggctaacc	ggaacccagc	cagcctctct	cccaccttct	660
accttctgtt	ctcaatgctg	tgccccctgg	ctggggcagc	ctgagctagg	tgatgtcgag	720
ccaggaacaa	gagccaagga	ccacagaacc	tgcttttggg	tccacatgtc	tctgccccat	780
cttccttttt	tctttttcca	cctcgtttct	ttttttttct	cttttctttc	tctctccctt	840
cctttccctt	cccttgccct	ccccattccc	cttcccttcc	ttttcttttc	tccttccctc	900
ctttcttttt	ctctttcttt	ctttttcttt	ctttctttct	ctttctttct	ttccctctct	960
ctccctccct	tccttccctc	ctcccttccct	tcctctcttt	ctttctttct	ttctctcttt	1020
ctctctttct	ttctctcttt	ctctctttct	ttctttttct	tagagaaagg	atgtccctct	1080
gttgcccagg	ctgcagtga	gtagcacaac	catggctcac	tgtaaccttg	aactcctggg	1140
ctcaagcaat	cctcctgcct	caacctcctg	agttgctggg	actacagggtg	tgagcaaatg	1200
cacccggcac	ttttctcctt	ttttatttta	aatactttta	gagtaattcc	tttacataaa	1260
ttgaacttca	taaaaagtta	gaactttgca	cattaaaaga	cactatcaac	agagtgaaaa	1320
gaccatccaa	gaatgagata	cagtatttgc	aaatcatata	tccgattggg	gattaatcca	1380
ggatatctaa	agaactctaa	aactcagtaa	caacaaaaac	ccaaacaacc	caattcaaga	1440
aatgagaaaa	ggacttgaat	agacattttt	ccaaaaaaga	tatacaaatg	atatactcat	1500
gttcataaca	gtattattca	caatagtaaa	ggtggaaaca	acctaataatg	tcactgacaa	1560
atgaataaat	aaacaaaatg	tgatacaata	aaaaaaaaaa	aaaaaaa		1607

<210> 53

<211> 1432

<212> DNA

<213> Homo sapiens

<400> 53

ggcacgagag	aaagttcatc	ctctgggctt	ggacacagta	gtgatgacag	cagttcacat	60
ggatgtaaag	taaatacagc	tccgcaggaa	tcctgaaaaa	taatttcta	gttactatct	120

taggaatagc	aaattatgtc	cagtcataga	gaagaaagct	tcataataat	acattcttac	180
ctaaagctca	ctgtcatgat	gttaggtatt	taaattctta	aagatgttgg	gttggttatt	240
agtgggtattt	ttatgttgtc	ttatttttagg	taagcttctg	tgtaaagcta	aaaatcctgt	300
gaatacaata	ctatccttta	caggcagaca	ttattggtaa	acaagatctt	gccctccaat	360
gaaatgactt	acatgtttta	aaaaaccgag	ttggttttat	tgaatttaaa	aagataggta	420
actaagtagc	atttaaaatc	aagatagagc	attccttctt	gtatcagtgg	ggcagtgtta	480
ccataaacac	gggttatatg	ttgttaaacc	ctatgaagag	taacagtgtg	gaccagactg	540
cctctctcag	atatgtgcct	gatattttgt	ggatacctcc	cctgcactgg	caaaacacta	600
tgcttttggg	tgtagactg	aaatatttta	agagtattta	acctttccag	tattctgttt	660
cacgcttaga	tggaaatgta	tcttatgaat	agagacatat	taaaataatg	tttacatctt	720
agaaaaaaca	tagatagtgc	tagtaatatt	acttataact	gtaatatata	gattcagaaa	780
tacattttca	ttatccaaaa	tcagcttcaa	caaatggttt	ctggagacaa	ataatttggt	840
ttcattatca	ttgtataatc	aggttaatga	tttatttttt	gactaaatgt	gcaatttctt	900
atcactagat	aactttcagt	atcagtgggtg	gttacttatt	acttaaatca	gaggaaggat	960
tttataagat	tcataaattt	aattttacca	ataaatattc	ccataattta	gaaaaggatg	1020
tcgacttgct	aatttcagaa	ataattattc	atttttataa	gcccccttta	aagcatctac	1080
ttgaagattg	gtataatttt	cataaaatgt	cttttttttt	agtgtcccaa	agatatctta	1140
gataaactat	tttgaagtgc	agatttcaga	tgaggcaaca	ttttcttgag	ataattacc	1200
aagtttcac	catgttgaat	ggtacaaaat	atttctgtga	aactaacagg	aagatatttt	1260
cagataacta	ggataacttg	ttgctttggt	accagccta	attgaagagt	ggcagaggct	1320
actacaaaa	gcaacctttt	cattttcact	aagagttaa	aagctattgt	attattaaaa	1380
agtctttaca	atgcttgttt	caaagaacca	acagaaaaaa	aaaaaaaaaa	aa	1432

<210> 54

<211> 2261

<212> DNA

<213> Homo sapiens

<400> 54

gtggccagt	ccctgggcta	ggagagggat	aaagtcagct	gtggccaagc	agaagcagta	60
ttgcagggga	aggggtggga	gagactgtgc	tatgagctct	gagcaggagg	caggacatgg	120
agagaagggt	gggagacgga	cagagccagc	tgtgccagcc	gagggaccag	agtgggcagt	180
ggggacggag	cacagaccac	cgcccacaag	ggtctctcct	gtgacttctg	gcttcccag	240
ggcagaggct	gggctgggca	tgtggcggt	ggcaccagc	aggctccgc	aggccacgc	300
caagccagcc	tggctcagct	ctggcttct	gctcacagc	tggatgcctg	tccccaggcc	360
tcctgacagg	gctctgcagc	attggagagg	cttgtggtgg	ggccctcgct	gtagaacagg	420
cactgccagt	gctcactgac	tcctcattgc	agccagggca	ggggcgctcc	caccacctcg	480
gtttcagtca	ggaagctggg	gggtgctggg	atctgccagc	agctctgtgt	gctccccagg	540
tgggctgccc	ggggccctgg	ctctctggct	tctacaccag	tggccctgcg	aagcctcagc	600
gcccagggtc	tttgcaagcc	ctgttctctg	atctctggag	agggtgctc	cagggtgtggg	660
cctggccagg	gcttcagac	cagtccaggc	actgcactg	cacattgacc	cccttctcca	720
ctctctctca	tccttcagta	gccacaaagc	tggcagcagg	gtactcgcag	atthgtgagg	780
agagccgagc	ccttgctgac	tgtcctccc	tgcacggcag	agtgaggcgg	ggctcctggc	840
ctccttaggg	gctgcccgt	ctgggccaca	cagccagaga	aacactgtcc	atctggctgg	900
cttggcctgt	gggtgcaaa	agaggtgttt	tcactttgct	ttggaaacat	ggaaattacc	960
aagtgactta	accataacac	caaatgatga	tttttaaat	tatgaaaatt	atggaaccag	1020
atgggacaca	gggaaacgag	aacgatactg	aaaaggctcg	agtcctgtgg	gggggtccagg	1080
tcccgaaga	cagctaata	ggtgcccctg	tctgttatc	gggtaaatgt	agcagctggg	1140
tccaggctcg	ggcagagcag	cagggtgtgc	cgccctctcc	cgggggtccc		1200
acgggggtcc	caggggcctc	ttactctgcc	acgatgtggg	agttccacca	ccacagggac	1260
ttgagcggca	gctccggctc	ttacgtagaa	acgcgcaact	ccagtcccta	ggttctgtcc	1320
gaggttgcta	tgggtccatc	ccatcttgcc	gctcactctg	cgactgtgcg	gagaaacgca	1380
agtgcccccg	aagggtgggc	gtggcctctg	atgaatgcac	acgttgggtg	gaggtggatt	1440
ccgtttgtac	gaagcgcctc	ttcacgcgag	cgttcacctc	ggtctcccct	ttgcttggtc	1500

```

cagttccaga aacgccgctg gactgcgagg tctccctgtg gtcgtcctgg ggactgtgcg 1560
gaggccactg tgggaggctc gggaccaaga gcaggactcg ctacgtccgg gtccagcccg 1620
ccaacaacgg gagccctgc cccgagctcg aagaagaggc tgagtgcgtc cctgataact 1680
gcgtctaaga ccagagcccc gcagccctg gggcccccgg gagccatggg gtgtcggggg 1740
ctcctgtgca ggctcatgct gcaggcggcc gagggcacag ggggtttcgc gctgctcctg 1800
accgcggtga ggccgcgcgg accatctctg cactgaaggg ccctctgggt gccggcacgg 1860
gcattgggaa acagcctcct cctttcccaa ccttgcttct taggggcccc cgtgtcccg 1920
ctgctctcag cctcctcctc ctgcaggata aagtcattcc caaggctcca gctactctaa 1980
attatgtctc cttataagtt attgctgctc caggagattg tccttcctcg tccagggggc 2040
tggtccccc gtggttgca atacctcaga cctggtgctc taggctgtgc tgagccact 2100
ctcccaggg cgcattccaag cgggggccac ttgagaagtg aataaatggg gcggtttcgg 2160
aagcgtcagt gtttccatgt tatggatctc tctgcgtttg aataaagact atctctgtt 2220
ctcaaaaaaa aaaaaaaaaa aaaaaaaaaa aaaaaaaaaa a 2261

```

```

<210> 55
<211> 813
<212> DNA
<213> Homo sapiens

```

```

<220>
<221> SITE
<222> (30)
<223> n equals a,t,g, or c

```

```

<220>
<221> SITE
<222> (54)
<223> n equals a,t,g, or c

```

```

<220>
<221> SITE
<222> (81)
<223> n equals a,t,g, or c

```

```

<220>
<221> SITE
<222> (97)
<223> n equals a,t,g, or c

```

```

<220>
<221> SITE
<222> (807)
<223> n equals a,t,g, or c

```

```

<400> 55
ttttaaccc ctttagggc ttccggggcn cggtatggtt ggtgtgggaa tttntggggc 60
gggttaacca atttaccacc nggraaccag ctatggncca tggatttasg ccagctcgaa 120
attamcccty cactaarggg amcaaaagct ggagctccac cgcggtggcg gccgtctag 180
aactagtgga tccccgggc tgcaggaatt cggcacgagg tttaggagga atacaattca 240
agaacctcac tttgaagcta accatagcaa acctagttta cctctagaat gcaggtgaca 300
gtgtgcaacc actcctgtca tctgtctttt gtatgctctt cataagcaat cttcctcttc 360
cttcccagtt catttattta gectctgatt cattctctc ttctccacc ctttctctt 420
ctaccagcca gctaccaat acttattctc tctgaaagga tcttatcctt taccagaagt 480
tcagttatta cctctaagaa ccagtccctt cccttttaaa ttttcttttc cctgaaagag 540

```

tcagttttat	gtattagaaa	aatgaaggta	tatgttttcc	tttaaaggga	ttataatcat	600
tacagaaaaat	tttaaagttt	tcaaaaaatt	aaaaaaaaaa	cccagaattt	cccaaacct	660
gactatccaa	agagaagcac	ttaatgttat	gtgtgtgac	ttgagtga	atttaaaatt	720
tatattagaa	cattttataa	attatccaaa	ctttttgtac	gaaaaaaaca	gaaaatgtaa	780
aaaaaaaaaa	aaaaactcga	ggggggnccc	gca			813

<210> 56

<211> 1392

<212> DNA

<213> Homo sapiens

<220>

<221> SITE

<222> (11)

<223> n equals a,t,g, or c

<220>

<221> SITE

<222> (17)

<223> n equals a,t,g, or c

<220>

<221> SITE

<222> (26)

<223> n equals a,t,g, or c

<220>

<221> SITE

<222> (71)

<223> n equals a,t,g, or c

<220>

<221> SITE

<222> (86)

<223> n equals a,t,g, or c

<400> 56

cttaagaacc	ncceccnatt	ctttgnccca	tccagtccttt	tctaacttgg	tctggtcttt	60
tgtcccatcc	nttttgaaga	ttttgnccga	wttttgamca	aggccaccaa	cctgcagcaa	120
tgamcaaagt	tccaatgcca	aataagtaaa	gaggggagcc	accacgaacc	cgggtgagggc	180
atcgagattt	tggtttccaa	catagcacaa	gccagtcagt	tcattctgcat	ccaccagtct	240
cataatcaag	atgacaatgg	ttttcactgc	ggggatggcc	caggctgcaa	tgtggaaata	300
agagctgtgc	atttcaatgg	cttcatgacc	ccattttgagt	cctgctgcca	aaaaccaagt	360
gagtgtcaga	ataaccacc	aaatggagct	ggccattcca	aaaaagtaca	tcagcaagaa	420
aattattgca	catcctgtgt	tcttaagtcc	ttcttggtatg	agaacagggt	ctgctgcctc	480
ttcaaaatca	caggatatcc	tttcccggcc	tacagtcagc	ctgacaataa	gcaatgctat	540
aaatattata	gcacatactg	agaaatatga	tggggcgctc	agggtaggaa	aacctagaag	600
aatcgatcag	gaaggtcagt	actgtgaagg	cagtggagat	gaaacacagg	ctggcccaca	660
cagccatcca	gatatcagtg	aactccttgg	ctgagcggct	gtataagcca	gcacatagc	720
cacacttgag	cacacagttc	aggctccttt	tcaccagat	gtactgatca	gaattgggtc	780
ccacagagtg	acactcttcc	ccaggctgga	tgggggtttt	gtgaggttaag	ggcacctctt	840
catcacctgg	cccttccatg	cacatgtggt	tgtggtcggt	ctgtggtggg	aatttgctgc	900
agttcagact	ctctggccag	gcaaatccaa	attccttcag	gacgggttca	cagcgtctct	960
tgactgaaag	acacatgccg	ccgcatgggc	caatggggat	gttgatcttc	tctgtgcaca	1020

ttggcacata	aacagaacaa	aggaagaact	ggaaaagtaa	caaaatgaac	acacacaaaa	1080
aaaacaatga	cttggaagtt	tgaccaaagt	ctcccacaaa	gctgagttga	atgcttccag	1140
gcaatctagg	tatctacttt	taaaccaacc	tatcgggagt	ctgtctgttt	actctgacct	1200
caaacaagg	cgctgataa	tccatcactt	acatactttt	cttagtttgg	ctttagtaat	1260
cctttccaga	agaatgtaca	taaataaata	aataaaataa	atagggtctg	aagaattgca	1320
taagccctct	ttgcacaaca	cttgaaataa	atcttaaaaa	actgaaaaaa	aaaaaaaaaa	1380
aaaaactcgt	ag					1392

<210> 57

<211> 1807

<212> DNA

<213> Homo sapiens

<220>

<221> SITE

<222> (1807)

<223> n equals a,t,g, or c

<400> 57

ccgggatcg	acccacgcgt	ccgctttaca	tatcatactt	tgggggttaa	ggagattcct	60
cagactcatc	cagcccttgg	gtgctgacca	gcagagtcac	tagtggatgc	tgaagttaca	120
tgagctacat	gttaaataat	taaagtctcc	aaaataaaac	accccaacgt	tgaccttacc	180
cggctgatgg	ttagccccct	gctgcctgct	ccatgtgtct	tatgagagcc	cgtagttaca	240
gtgtcctcta	atltgaaatc	cataagttaa	caagtctata	tcagggtgcag	ctggctttga	300
ttaaaggcca	tttttaaaac	ttaaaaactc	aacacctcac	agattataat	agaaaaagaa	360
atggcctcag	tttgatctcg	ttcagaatga	cccagattgt	ttctgctttg	ggtgcagctg	420
tttagttcag	agttatatta	cagagaatta	ttttctgaga	taactctaaa	ctagaatggt	480
caaaactaat	tgataattga	agtatcaaga	tacgtagaac	acctcagaga	ttttcttcca	540
ggaacttcca	caaactttga	atccttgat	ctttatattg	tattcatact	actagtagca	600
aaatacaggt	tttttgtttt	gttttgtttt	ggcttcatag	agtatctcaa	attgaaactt	660
ttctgcacaa	agaataaaat	taaggatttt	ataaactcaa	attggcacct	actgaattaa	720
aatacataaa	atcattttaa	tataattcag	catatgggaa	gtaacattgc	actaatatgg	780
aatcactgc	cagagacagt	ctattttctt	ttaatattgt	actacttagt	cacaaacccc	840
acattattcc	agtttggaat	tacttattaa	ggagaattgg	aaatacatat	gcccattgct	900
aaattttata	gctttaattt	gtgttatttc	tttattgacg	ggaagaggta	catctttttt	960
tccttactga	aaacaaatat	ggattaattg	cctcaaattt	gtataagtga	ttggctagtg	1020
attcttggtt	tcagaaggga	gagtggata	gatagaaaat	gacaaagatg	gcaatataca	1080
cttaattgtt	ttattgtatg	ttgttactga	agtacttaga	tttttaaaat	ttcaaactct	1140
aatcacttc	ttgtaggagg	gttttcatta	actgcagtat	atacagttca	ctacatatgg	1200
gttggttgag	ttttttgtgt	gctgtatttc	tttctgtttt	ttaatacctg	gttttgtaga	1260
tatctaactc	tggtctcttt	tggttggttc	gaaactggat	tttttttttc	ttaagcagtg	1320
cttaatttgt	gttttttaat	tttgattcag	aagtagtccc	agctcatagg	tgttcatact	1380
gttacatcca	gaacatttgt	caggctctct	gtcagcttcc	atgtacatat	ggtatagaaa	1440
ccatggagtt	aggcacttcc	tggatttttt	ttttatgaga	aaaatactgt	atttaaaatg	1500
taaaataaac	tttttaaaaag	caggcactaa	tatatatttc	ttccagcctt	tgattacaaa	1560
tttgctcttg	cacatgttaa	gatgaattat	ctcctaaaaa	tatcattgtt	cttgggagca	1620
gtgtatgtta	ctttacatag	cagcggttcc	tgtcatgtgt	tcatgtcaga	atattttttg	1680
ttttaaactt	tcttattgcc	tttggtgtgt	gattagtaca	gtacaagtgc	gattttcaaaa	1740
agatcttgaa	agtaatatat	ttaatcaatt	aaaatgttta	tctgtaaaaa	aaaaaaaaaa	1800
aaaaaan						1807

<210> 58

<211> 44

<212> PRT
 <213> Homo sapiens

<220>
 <221> SITE
 <222> (44)
 <223> Xaa equals stop translation

<400> 58

Met Gly Ile Phe Asp Tyr Lys Leu Phe Ser His Tyr Phe Lys Ala Cys
 1 5 10 15

Phe Ile Phe Phe Leu Ile Leu Leu Thr His Leu Cys Leu Ser Leu Phe
 20 25 30

Tyr Tyr Lys Leu Phe Ile Val Gln Ser Leu Pro Xaa
 35 40

<210> 59
 <211> 140
 <212> PRT
 <213> Homo sapiens

<220>
 <221> SITE
 <222> (140)
 <223> Xaa equals stop translation

<400> 59

Met Lys Lys Ile Val Asp Gln Asn Thr Lys Leu Ala Pro Glu Thr Lys
 1 5 10 15

Ala Val Ile His Trp Ile Met Asp Ile Pro Phe Val Leu Ser Ala Asn
 20 25 30

Leu His Gly Gly Asp Leu Val Ala Asn Tyr Pro Tyr Asp Glu Thr Arg
 35 40 45

Ser Gly Ser Ala His Glu Tyr Ser Ser Ser Pro Asp Asp Ala Ile Phe
 50 55 60

Gln Ser Leu Ala Arg Ala Tyr Ser Ser Phe Asn Pro Ala Met Ser Asp
 65 70 75 80

Pro Asn Arg Pro Pro Cys Arg Lys Asn Asp Asp Asp Ser Ser Phe Val
 85 90 95

Asp Gly Thr Thr Asn Gly Val Leu Gly Thr Ala Tyr Leu Glu Gly Cys
 100 105 110

Lys Thr Ser Ile Thr Leu Ala Ala Thr Val Leu Arg Ser Pro Trp Ser
 115 120 125

Leu Ala Val Arg Ser Ser His Leu Lys Arg Leu Xaa

130

135

140

<210> 60

<211> 2

<212> PRT

<213> Homo sapiens

<220>

<221> SITE

<222> (2)

<223> Xaa equals stop translation

<400> 60

Phe Xaa

1

<210> 61

<211> 255

<212> PRT

<213> Homo sapiens

<220>

<221> SITE

<222> (222)

<223> Xaa equals any of the naturally occurring L-amino acids

<220>

<221> SITE

<222> (255)

<223> Xaa equals stop translation

<400> 61

Met	Gln	Gly	Val	Ile	Leu	Leu	Leu	Phe	Ala	Lys	Tyr	Tyr	His	Leu	Pro
1				5					10					15	

Phe	Leu	Arg	Asp	Val	Gln	Thr	Asp	Cys	Thr	Arg	Thr	Gly	Leu	Gly	Gly
		20						25					30		

Tyr	Trp	Gly	Asn	Lys	Gly	Gly	Val	Ser	Val	Arg	Leu	Ala	Ala	Phe	Gly
		35					40						45		

His	Met	Leu	Cys	Phe	Leu	Asn	Cys	His	Leu	Pro	Ala	His	Met	Asp	Lys
	50					55						60			

Ala	Glu	Gln	Arg	Lys	Asp	Asn	Phe	Gln	Thr	Ile	Leu	Ser	Leu	Gln	Gln
65					70					75				80	

Phe	Gln	Gly	Pro	Gly	Ala	Gln	Gly	Ile	Leu	Asp	His	Asp	Leu	Val	Phe
			85					90						95	

Trp	Phe	Gly	Asp	Leu	Asn	Phe	Arg	Ile	Glu	Ser	Tyr	Asp	Leu	His	Phe
			100					105						110	

Val Lys Phe Ala Ile Asp Ser Asp Gln Leu His Gln Leu Trp Glu Lys
 115 120 125
 Asp Gln Leu Asn Met Ala Lys Asn Thr Trp Pro Ile Leu Lys Gly Phe
 130 135 140
 Gln Glu Gly Pro Leu Asn Phe Ala Pro Thr Phe Lys Phe Asp Val Gly
 145 150 155 160
 Thr Asn Lys Tyr Asp Thr Ser Ala Lys Lys Arg Lys Pro Ala Trp Thr
 165 170 175
 Asp Arg Ile Leu Trp Lys Val Lys Ala Pro Gly Gly Gly Pro Ser Pro
 180 185 190
 Ser Gly Arg Lys Ser His Arg Leu Gln Val Thr Gln His Ser Tyr Arg
 195 200 205
 Ser His Met Glu Tyr Thr Val Ser Asp His Lys Pro Val Xaa Ala Gln
 210 215 220
 Phe Leu Leu Gln Phe Ala Phe Gln Gly Arg His Ala Thr Gly Ala Ala
 225 230 235 240
 Gly Gly Gly Gln Met Ser Gly Cys Gly Pro Ser Arg Arg Trp Xaa
 245 250 255

<210> 62
 <211> 7
 <212> PRT
 <213> Homo sapiens

 <220>
 <221> SITE
 <222> (7)
 <223> Xaa equals stop translation

<400> 62
 Gly Val Ser Leu His Gln Xaa
 1 5

<210> 63
 <211> 49
 <212> PRT
 <213> Homo sapiens

<220>
 <221> SITE
 <222> (49)
 <223> Xaa equals stop translation

<400> 63
 Glu Leu Ser Cys Trp Gln Asp Leu Leu Glu Leu Ala Arg Gln Leu Trp

1 5 10 15
 Leu Trp Leu Leu Leu Arg Ser Trp Val Val Arg Ser Pro Ser Ala Gln
 20 25 30
 Trp Trp Gly Val Lys Phe Thr Gln Leu Arg Ser Arg Arg Gln Arg Cys
 35 40 45

Xaa

<210> 64
 <211> 30
 <212> PRT
 <213> Homo sapiens

<220>
 <221> SITE
 <222> (30)
 <223> Xaa equals stop translation

<400> 64
 Met Glu Leu Leu Phe Phe Leu Leu Leu Pro Tyr Phe Leu Leu Phe Ile
 1 5 10 15
 Cys Leu Ile Ser Cys Ile Ser Gln Ile Tyr Ile Tyr Leu Xaa
 20 25 30

<210> 65
 <211> 56
 <212> PRT
 <213> Homo sapiens

<220>
 <221> SITE
 <222> (56)
 <223> Xaa equals stop translation

<400> 65
 Met Thr Asn Tyr Phe Trp Ala Ser Ala Ser Ser Cys Thr Val Phe Pro
 1 5 10 15
 Leu Ala Phe Leu Cys Ser Ser Cys Val Gly Pro Pro Ser Phe Ser Cys
 20 25 30
 Ala His His Ile Leu His Gln Glu Ser Trp His Leu Phe Phe Ser Ser
 35 40 45
 Ala Trp Asn Ala Phe Leu Cys Xaa
 50 55

<210> 66

<211> 36
 <212> PRT
 <213> Homo sapiens

<220>
 <221> SITE
 <222> (36)
 <223> Xaa equals stop translation

<400> 66
 Met Thr Met Lys Leu Ser Val Phe Leu Ser Val Gln Asn Leu Gly Leu
 1 5 10 15
 Leu Ser Val Val Leu Lys Arg Ala Leu Ala Leu Ser Thr Pro Ser Leu
 20 25 30
 Glu Ile Cys Xaa
 35

<210> 67
 <211> 14
 <212> PRT
 <213> Homo sapiens

<220>
 <221> SITE
 <222> (14)
 <223> Xaa equals stop translation

<400> 67
 Gly Asn Ser Leu Thr Leu Ala Ile Leu Leu Leu Ser Phe Xaa
 1 5 10

<210> 68
 <211> 45
 <212> PRT
 <213> Homo sapiens

<220>
 <221> SITE
 <222> (45)
 <223> Xaa equals stop translation

<400> 68
 Asp Leu Gln Ile Gln Trp Pro Ile Leu Leu Ser Leu Ser Cys Glu Gly
 1 5 10 15
 Val Phe Gln Val Leu Lys Gln Ser Lys Asn His Leu Gly Pro Ser Leu
 20 25 30
 Arg Lys His Phe Ser Gly Gln Val Gly Phe Arg Leu Xaa
 35 40 45

<210> 69
<211> 15
<212> PRT
<213> Homo sapiens

<220>
<221> SITE
<222> (15)
<223> Xaa equals stop translation

<400> 69
Gly Phe Tyr Thr Phe Leu Pro Ser Leu Pro Gly Ala Leu Tyr Xaa
1 5 10 15

<210> 70
<211> 3
<212> PRT
<213> Homo sapiens

<220>
<221> SITE
<222> (3)
<223> Xaa equals stop translation

<400> 70
Met Tyr Xaa
1

<210> 71
<211> 13
<212> PRT
<213> Homo sapiens

<220>
<221> SITE
<222> (13)
<223> Xaa equals stop translation

<400> 71
Met Glu Arg Val Arg Thr Ser Met Glu Cys Phe Cys Xaa
1 5 10

<210> 72
<211> 48
<212> PRT
<213> Homo sapiens

<400> 72
Met Leu Val Ile Leu Leu Asp Met Phe Phe Val Val Val Val Thr Trp
1 5 10 15

Asn Phe Cys Ile Leu Asn Lys Phe Gly Asp Gln Ile Gln Lys Lys Lys
20 25 30

Lys Lys Lys Lys Lys Lys Lys Lys Lys Lys Lys Lys Lys Lys Lys
35 40 45

<210> 73
<211> 49
<212> PRT
<213> Homo sapiens

<220>
<221> SITE
<222> (49)
<223> Xaa equals stop translation

<400> 73
Gly Thr Pro Ser Leu Leu Pro Tyr Ser Cys Ser Arg Phe Ser Gly Leu
1 5 10 15

Asn Gln Val Asn Arg Gly Pro Glu Pro Pro Leu Cys Leu Leu Glu Ile
20 25 30

Pro Lys Thr Glu Met Cys Ser Arg Ser Arg Ser Lys Thr Leu Ser Leu
35 40 45

Xaa

<210> 74
<211> 60
<212> PRT
<213> Homo sapiens

<220>
<221> SITE
<222> (60)
<223> Xaa equals stop translation

<400> 74
Met Gly Gly His Gly Ser Ala Gly Leu Cys Leu Lys His Thr Leu Gln
1 5 10 15

Leu Gln Leu Phe Ser Leu Val Phe Cys Leu Lys Leu Ile Leu Thr Glu
20 25 30

Ser Asp Phe Val Phe Ile Ser Phe Gln Gly Leu Gly Cys Leu Trp Ala
35 40 45

Ser Pro Gly Gly Leu Glu Val Gly Lys Gly Lys Xaa

50

55

60

<210> 75
<211> 45
<212> PRT
<213> Homo sapiens

<220>
<221> SITE
<222> (45)
<223> Xaa equals stop translation

<400> 75
Met Ser Gly Leu Ser Gly Ala Pro Trp Gly Thr Trp Ala Leu Pro Leu
1 5 10 15

Ala His Leu Ser Leu Leu Ser Arg Thr Cys Leu Ser Ser Ser Pro Leu
20 25 30

Gly Phe Arg Leu Phe Ser Arg His Leu Arg His Leu Xaa
35 40 45

<210> 76
<211> 39
<212> PRT
<213> Homo sapiens

<220>
<221> SITE
<222> (39)
<223> Xaa equals stop translation

<400> 76
Met Leu Ser Leu Pro Ala Pro Leu Arg Val Asn Arg Gly Leu Trp Gln
1 5 10 15

Leu Cys Thr Gly Ala Gly Leu Trp Leu Leu Gln Gly Ala Leu Pro Val
20 25 30

Thr Arg Ser Trp Ala Val Xaa
35

<210> 77
<211> 134
<212> PRT
<213> Homo sapiens

<220>
<221> SITE
<222> (134)
<223> Xaa equals stop translation

<400> 77

Met Phe Thr Asp Leu Ser Ser Ser Trp Cys Met Thr Met Thr Pro Val
 1 5 10 15

Gly Ser Met Thr Ser Ser Ala Ser Ser Pro Ala Leu Ser Arg Arg Cys
 20 25 30

Arg Lys Gly Arg Gln Thr Leu Gly Arg Arg Cys Ser Gly Thr Val Ser
 35 40 45

Thr Gln Val Ser Gly Gln Glu Glu Glu Leu Gln Glu Leu Arg Asp Gly
 50 55 60

Ser Ala Gly Pro Val His Gly Gly Glu Gly Ala His Leu Pro Gly Leu
 65 70 75 80

His His Gly Trp Leu Pro Asp Gln Leu His Gly Gly His Arg Leu His
 85 90 95

Arg Leu Gln Trp Gly Pro Glu Glu Gln Pro Val Pro Ala Leu Pro Gln
 100 105 110

Ser Pro Thr Ala Gln Pro Leu Pro Ala Gly Pro Ala Cys Ser Gly Arg
 115 120 125

His Leu Pro Gly Leu Xaa
 130

<210> 78

<211> 9

<212> PRT

<213> Homo sapiens

<220>

<221> SITE

<222> (9)

<223> Xaa equals stop translation

<400> 78

Trp Ala Phe Ser Ala Trp Leu Leu Xaa
 1 5

<210> 79

<211> 10

<212> PRT

<213> Homo sapiens

<220>

<221> SITE

<222> (10)

<223> Xaa equals stop translation

<400> 79

Ile Phe Leu Ile Leu Leu Ile Cys Val Xaa
1 5 10

<210> 80

<211> 91

<212> PRT

<213> Homo sapiens

<220>

<221> SITE

<222> (91)

<223> Xaa equals stop translation

<400> 80

Met Ala Asp Arg Ala Pro Arg Gln Pro Thr Ser Leu Gly Ser Ser Pro
1 5 10 15

Gln Arg Cys Gly Ser Pro Pro Ser Ala Pro Pro Pro Pro Pro Asp Arg
20 25 30

Gly Glu Arg Ile Glu Asp Cys Leu Ala Pro Leu Cys Pro Pro Val Val
35 40 45

Gly Gly Arg Gly Gly Cys Ala Leu Ser Arg Phe Leu Pro Val Asp Leu
50 55 60

Ser Leu Leu Arg Ser Ser Arg Thr Ala Ala Val Glu Leu Val Ala Gly
65 70 75 80

Gly Val Cys Arg Ser Pro Ser Gln Glu Asn Xaa
85 90

<210> 81

<211> 43

<212> PRT

<213> Homo sapiens

<220>

<221> SITE

<222> (43)

<223> Xaa equals stop translation

<400> 81

Ile Ile Val Arg Val Gly Phe Ala Leu Ser Cys Ser Val Ser Cys Pro
1 5 10 15

Phe Phe Ala Leu Leu Ser Cys Asp Ala Phe His His Ile Met Thr Gln
20 25 30

Gln Glu Gly Pro His Lys Val Pro Ala Pro Xaa
35 40

<210> 82
<211> 43
<212> PRT
<213> Homo sapiens

<220>
<221> SITE
<222> (43)
<223> Xaa equals stop translation

<400> 82
Met Lys Ile Phe Met Tyr Val Pro Ile Ile Leu Phe Arg Ser Gln Cys
1 5 10 15
Leu Cys Leu Ser Leu Thr Glu Ile Ile Glu Leu Leu Ser Ser Ser Lys
20 25 30
Leu Asn Lys Thr Leu Glu Leu Lys Ala Ser Xaa
35 40

<210> 83
<211> 88
<212> PRT
<213> Homo sapiens

<220>
<221> SITE
<222> (88)
<223> Xaa equals stop translation

<400> 83
Met Glu Gly Pro Leu Gln Ser Phe Lys Arg Arg Leu Trp Gly Gly Val
1 5 10 15
Glu Cys Ala Leu Gly Leu Asp Thr Cys Arg Gly Ala Pro Gly Trp Pro
20 25 30
Arg Arg Leu Ala Leu Ala Arg Gly Pro Val Leu Arg Gly Pro Gln Arg
35 40 45
Leu Thr Leu Gly Gln Ala Pro Ala Arg Arg Gly Arg Ser Pro Gly Leu
50 55 60
Pro Gly Arg His Val His Ile Ala Ile Ala Thr Arg Gln Leu Arg Leu
65 70 75 80
Gly Phe Thr Lys Ser Leu Pro Xaa
85

<210> 84
<211> 52
<212> PRT
<213> Homo sapiens

<220>
<221> SITE
<222> (52)
<223> Xaa equals stop translation

<400> 84
Met Thr Thr Ser Gln Ala Ile Ser Cys Leu Ala His Trp Ala Cys Ala
1 5 10 15
Trp Glu Thr Ala Cys Cys Trp Met Ala Arg Arg Arg Ala His Cys Gly
20 25 30
Ser Val Gly Pro Arg Ser Leu Pro Ala Ala Val Gly Gly Arg Gly Ala
35 40 45
Gly Arg Thr Xaa
50

<210> 85
<211> 77
<212> PRT
<213> Homo sapiens

<220>
<221> SITE
<222> (77)
<223> Xaa equals stop translation

<400> 85
Val Gly His His Ser Arg Pro Trp Thr Ser Pro Asn Leu Gly Thr Ala
1 5 10 15
Leu Phe Pro Asn Val Ser Ile Leu Leu Thr Gly Ala Ser Ser Ala Gln
20 25 30
Pro Trp Glu Pro Ile Gln Gly Leu Val Arg Thr Gly Leu Pro Gly Val
35 40 45
Gly Val Trp Gly Val Pro Ser Gly Val Ile Asp Ser Pro Thr Ala Pro
50 55 60
Ala Leu Thr Gly Pro Gln Val Ala Ala Met Val Asn Xaa
65 70 75

<210> 86
<211> 33
<212> PRT
<213> Homo sapiens

<220>
<221> SITE
<222> (33)

<223> Xaa equals stop translation

<400> 86

Lys Ala Ile Trp Phe Leu Ile Leu Cys Thr Thr His Ser Ile Leu Ile
1 5 10 15

Ile Thr Phe Ile Tyr Lys Lys Asn Lys Glu Asn Asn Ser Lys Leu Cys
20 25 30

Xaa

<210> 87

<211> 40

<212> PRT

<213> Homo sapiens

<220>

<221> SITE

<222> (40)

<223> Xaa equals stop translation

<400> 87

Met Ala Gly Lys Ala Ser Leu Thr Ser Ser Ala Ser Val Ile Ala Trp
1 5 10 15

Pro Ser Leu Leu Cys Leu Leu Leu Leu Trp Met Pro Val Ile Gly Phe
20 25 30

Cys Ala Tyr His Thr Pro Met Xaa
35 40

<210> 88

<211> 67

<212> PRT

<213> Homo sapiens

<220>

<221> SITE

<222> (67)

<223> Xaa equals stop translation

<400> 88

Met Cys Cys Pro Ser Leu Leu Lys Phe Tyr Phe Arg Phe Ser Ile Gly
1 5 10 15

Tyr Leu Phe Cys Phe Leu Tyr Phe Phe Ser Leu Ser Leu Pro Pro Ser
20 25 30

Arg Pro Pro Arg Pro Ile Pro Phe Leu Pro Leu Asp Phe His Pro Leu
35 40 45

Gly Cys Leu Ala His Leu Tyr Ala Pro Ala Leu Gly Thr Gly Pro Asn

50

55

60

Thr Trp Xaa
65

<210> 89
<211> 56
<212> PRT
<213> Homo sapiens

<220>
<221> SITE
<222> (56)
<223> Xaa equals stop translation

<400> 89
Met Ser Val Gly Leu Ser Ser Ser Phe Leu Ile Leu Phe Met Pro Cys
1 5 10 15

Gln Val Tyr Cys Leu Tyr Pro Leu Val Asn Phe Phe Ile Ser Val Ile
20 25 30

Ile Arg Leu Lys Lys Lys Lys Lys Lys Lys Glu Ser Ser Leu Leu
35 40 45

Gly Asn Ala Trp Thr Leu Cys Xaa
50 55

<210> 90
<211> 47
<212> PRT
<213> Homo sapiens

<220>
<221> SITE
<222> (13)
<223> Xaa equals any of the naturally occurring L-amino acids

<220>
<221> SITE
<222> (47)
<223> Xaa equals stop translation

<400> 90
Met His Leu Ile Ile Phe Phe Leu Ser Leu Leu Cys Xaa Phe Leu Lys
1 5 10 15

Leu Pro Pro Lys Tyr Leu Ser Thr Pro Leu Asn Val Trp Ser Gln Asp
20 25 30

Lys Phe Leu Ile Leu Ala Leu Gln Phe Lys Met Tyr Lys Lys Xaa
35 40 45

<210> 91
 <211> 2
 <212> PRT
 <213> Homo sapiens

<220>
 <221> SITE
 <222> (2)
 <223> Xaa equals stop translation

<400> 91
 Leu Xaa
 1

<210> 92
 <211> 35
 <212> PRT
 <213> Homo sapiens

<220>
 <221> SITE
 <222> (35)
 <223> Xaa equals stop translation

<400> 92
 Leu Glu Leu Thr Val Leu Ser Ser Leu Arg Thr Phe Glu Tyr Thr Leu
 1 5 10 15
 Pro Ile Ser Leu Pro Tyr Phe Leu Phe Ala Ala Phe Ala Leu Glu Leu
 20 25 30
 Cys Phe Xaa
 35

<210> 93
 <211> 56
 <212> PRT
 <213> Homo sapiens

<220>
 <221> SITE
 <222> (56)
 <223> Xaa equals stop translation

<400> 93
 Met Trp Ala Leu Lys Ser Leu Phe Leu Leu Thr Pro Ser Pro Val Ile
 1 5 10 15
 Arg Phe Tyr Phe Ala Ala Leu Trp Ile Arg Ala Ala Gly Arg Leu Leu
 20 25 30
 Gly Gly Gly Gly Ser Pro Thr Pro Pro Thr Ser Leu Ala Pro Gly Phe

35

40

45

Ser Glu Ala Gly Gly Leu Cys Xaa
50 55

<210> 94

<211> 2

<212> PRT

<213> Homo sapiens

<220>

<221> SITE

<222> (2)

<223> Xaa equals stop translation

<400> 94

Met Xaa

1

<210> 95

<211> 24

<212> PRT

<213> Homo sapiens

<220>

<221> SITE

<222> (24)

<223> Xaa equals stop translation

<400> 95

Val Gly His Phe Trp Val Val Ile Trp Leu Val Arg Ser Met Ser Asp
1 5 10 15

Arg Met Asn Lys Asn Ala Leu Xaa
20

<210> 96

<211> 23

<212> PRT

<213> Homo sapiens

<220>

<221> SITE

<222> (23)

<223> Xaa equals stop translation

<400> 96

Met Leu Met Pro Pro Pro Pro Pro Ala Arg Leu Phe Phe Leu Tyr Phe
1 5 10 15

Val Leu Phe Leu Leu Gly Xaa
20

<210> 97
<211> 5
<212> PRT
<213> Homo sapiens

<220>
<221> SITE
<222> (5)
<223> Xaa equals stop translation

<400> 97
Phe Gly Phe Val Xaa
1 5

<210> 98
<211> 2
<212> PRT
<213> Homo sapiens

<220>
<221> SITE
<222> (2)
<223> Xaa equals stop translation

<400> 98
Met Xaa
1

<210> 99
<211> 18
<212> PRT
<213> Homo sapiens

<220>
<221> SITE
<222> (18)
<223> Xaa equals stop translation

<400> 99
Met Val Gly Leu Trp Phe Val Phe Ser Leu Tyr Ser Leu Val Thr Phe
1 5 10 15

Leu Xaa

<210> 100
<211> 39
<212> PRT
<213> Homo sapiens

<220>

<221> SITE

<222> (39)

<223> Xaa equals stop translation

<400> 100

Met Leu Gly Cys Leu Leu Val Val Phe Leu Cys Cys Leu Ile Leu Gly
1 5 10 15

Lys Leu Leu Cys Lys Ala Lys Asn Pro Val Asn Thr Ile Leu Ser Phe
20 25 30

Thr Gly Arg His Tyr Trp Xaa
35

<210> 101

<211> 60

<212> PRT

<213> Homo sapiens

<220>

<221> SITE

<222> (60)

<223> Xaa equals stop translation

<400> 101

Met Trp Arg Leu Ala Pro Arg Arg Leu Arg Gln Val His Ala Lys Pro
1 5 10 15

Ala Trp Leu Ser Ser Gly Phe Leu Leu Thr Arg Trp Met Pro Val Pro
20 25 30

Arg Pro Pro Asp Arg Ala Leu Gln His Trp Arg Gly Leu Trp Trp Gly
35 40 45

Pro Arg Cys Arg Thr Gly Thr Ala Ser Ala His Xaa
50 55 60

<210> 102

<211> 41

<212> PRT

<213> Homo sapiens

<220>

<221> SITE

<222> (41)

<223> Xaa equals stop translation

<400> 102

Met Leu Phe Ile Ser Asn Leu Pro Leu Pro Ser Gln Phe Ile Tyr Leu
1 5 10 15

Ala Ser Asp Ser Phe Phe Ser Ser Pro Thr Pro Phe Ser Ser Thr Ser

20 25 30
 Gln Pro Thr Asn Thr Tyr Ser Leu Xaa
 35 40

 <210> 103
 <211> 36
 <212> PRT
 <213> Homo sapiens

 <220>
 <221> SITE
 <222> (36)
 <223> Xaa equals stop translation

 <400> 103
 Ala His Val Val Val Val Val Leu Trp Trp Glu Phe Ala Ala Val Gln
 1 5 10 15
 Thr Leu Trp Pro Gly Lys Ser Lys Phe Leu Gln Asp Gly Phe Thr Ala
 20 25 30
 Ser Leu Asp Xaa
 35

<210> 104
 <211> 33
 <212> PRT
 <213> Homo sapiens

 <220>
 <221> SITE
 <222> (33)
 <223> Xaa equals stop translation

 <400> 104
 Tyr Leu Leu Lys Ile Ser Leu Phe Leu Gly Ala Val Tyr Val Thr Leu
 1 5 10 15
 His Ser Ser Gly Ser Cys His Val Phe Met Ser Glu Tyr Phe Trp Phe
 20 25 30
 Xaa

<210> 105
 <211> 109
 <212> PRT
 <213> Homo sapiens

 <400> 105
 Asn Asn His Leu Leu Lys Asn Leu Lys Lys Ile Val Asp Gln Asn Ser

1 5 10 15
 Lys Leu Ala Pro Glu Thr Lys Ala Val Ile His Trp Ile Met Asp Ile
 20 25 30
 Pro Phe Val Leu Ser Ala Asn Leu His Gly Gly Asp Leu Val Ala Asn
 35 40 45
 Tyr Pro Tyr Asp Glu Thr Arg Ser Gly Thr Ala His Glu Tyr Ser Ser
 50 55 60
 Cys Pro Asp Asp Ala Ile Phe Gln Ser Leu Ala Arg Ala Tyr Ser Ser
 65 70 75 80
 Phe Asn Pro Val Met Ser Asp Pro Asn Arg Pro Pro Cys Cys Lys Asn
 85 90 95
 Asp Asp Asp Ser Ser Phe Asp Asp Gly Thr Thr Asn Gly
 100 105

<210> 106
 <211> 109
 <212> PRT
 <213> Homo sapiens

<400> 106
 Asn Asn His Leu Leu Lys Asn Met Lys Lys Ile Val Asp Gln Asn Thr
 1 5 10 15
 Lys Leu Ala Pro Glu Thr Lys Ala Val Ile His Trp Ile Met Asp Ile
 20 25 30
 Pro Phe Val Leu Ser Ala Asn Leu His Gly Gly Asp Leu Val Ala Asn
 35 40 45
 Tyr Pro Tyr Asp Glu Thr Arg Ser Gly Ser Ala His Glu Tyr Ser Ser
 50 55 60
 Ser Pro Asp Asp Ala Ile Phe Gln Ser Leu Ala Arg Ala Tyr Ser Ser
 65 70 75 80
 Phe Asn Pro Ala Met Ser Asp Pro Asn Arg Pro Pro Cys Arg Lys Asn
 85 90 95
 Asp Asp Asp Ser Ser Phe Val Asp Gly Thr Thr Asn Gly
 100 105

<210> 107
 <211> 98
 <212> PRT
 <213> Homo sapiens

<400> 107

Ala Trp Tyr Ser Val Pro Gly Gly Met Gln Asp Phe Asn Tyr Leu Ser
 1 5 10 15
 Ser Ser Asn Cys Phe Glu Ile Thr Val Glu Leu Thr Cys Glu Lys Phe
 20 25 30
 Pro Pro Glu Glu Thr Leu Lys Ser Tyr Trp Glu Asp Asn Lys Asn Ser
 35 40 45
 Leu Ile Asn Tyr Leu Glu Gln Ile His Arg Gly Val Lys Gly Phe Val
 50 55 60
 Arg Asp Leu Gln Gly Asn Pro Asn Ala Asn Ala Thr Ile Ser Val Asp
 65 70 75 80
 Gly Ile Asp His Asp Val Thr Ser Ala Lys Asp Gly Asp Tyr Trp Arg
 85 90 95
 Leu Leu

<210> 108
 <211> 94
 <212> PRT
 <213> Homo sapiens

<400> 108
 Ala Trp Tyr Ser Val Pro Gly Gly Met Gln Asp Phe Asn Tyr Leu Ser
 1 5 10 15
 Ser Asn Cys Phe Glu Ile Thr Val Glu Leu Ser Cys Glu Lys Phe Pro
 20 25 30
 Pro Glu Glu Thr Leu Lys Thr Tyr Trp Glu Asp Asn Lys Asn Ser Leu
 35 40 45
 Ile Ser Tyr Leu Glu Gln Ile His Arg Gly Val Lys Gly Phe Val Arg
 50 55 60
 Asp Leu Gln Gly Asn Pro Ile Ala Met Ala Thr Asn Leu Arg Gly Arg
 65 70 75 80
 Asn Arg Pro Arg Cys Tyr Ile Arg Lys Gly Trp Arg Leu Leu
 85 90

<210> 109
 <211> 381
 <212> PRT
 <213> Homo sapiens

<400> 109
 Leu His Phe Val Lys Phe Ala Ile Asp Ser Asp Gln Leu His Gln Leu
 1 5 10 15

Trp Glu Lys Asp Gln Leu Asn Met Ala Lys Asn Thr Trp Pro Ile Leu
 20 25 30
 Lys Gly Phe Gln Glu Gly Pro Leu Asn Phe Ala Pro Thr Phe Lys Phe
 35 40 45
 Asp Val Gly Thr Asn Lys Tyr Asp Thr Ser Ala Lys Lys Arg Lys Pro
 50 55 60
 Ala Trp Thr Asp Arg Ile Leu Trp Lys Val Lys Ala Pro Gly Gly Gly
 65 70 75 80
 Pro Ser Pro Ser Gly Arg Lys Ser His Arg Leu Gln Val Thr Gln His
 85 90 95
 Ser Tyr Arg Ser His Met Glu Tyr Thr Val Ser Asp His Lys Pro Val
 100 105 110
 Ala Ala Gln Phe Leu Leu Gln Phe Ala Phe Arg Asp Asp Met Pro Leu
 115 120 125
 Val Arg Leu Glu Val Ala Asp Glu Trp Val Arg Pro Glu Gln Ala Val
 130 135 140
 Val Arg Tyr Arg Met Glu Thr Val Phe Ala Arg Ser Ser Trp Asp Trp
 145 150 155 160
 Ile Gly Leu Tyr Arg Val Gly Phe Arg His Cys Lys Asp Tyr Val Ala
 165 170 175
 Tyr Val Trp Ala Lys His Glu Asp Val Asp Gly Asn Thr Tyr Gln Val
 180 185 190
 Thr Phe Ser Glu Glu Ser Leu Pro Lys Gly His Gly Asp Phe Ile Leu
 195 200 205
 Gly Tyr Tyr Ser His Asn His Ser Ile Leu Ile Gly Ile Thr Glu Pro
 210 215 220
 Phe Gln Ile Ser Leu Pro Ser Ser Glu Leu Ala Ser Ser Ser Thr Asp
 225 230 235 240
 Ser Ser Gly Thr Ser Ser Glu Gly Glu Asp Asp Ser Thr Leu Glu Leu
 245 250 255
 Leu Ala Pro Lys Ser Arg Ser Pro Ser Pro Gly Lys Ser Lys Arg His
 260 265 270
 Arg Ser Arg Ser Pro Gly Leu Ala Arg Phe Pro Gly Leu Ala Leu Arg
 275 280 285
 Pro Ser Ser Arg Glu Arg Arg Gly Ala Ser Arg Ser Pro Ser Pro Gln
 290 295 300

Ser Arg Arg Leu Ser Arg Val Ala Pro Asp Arg Ser Ser Asn Gly Ser
305 310 315 320

Ser Arg Gly Ser Ser Glu Glu Gly Pro Ser Gly Leu Pro Gly Pro Trp
325 330 335

Ala Phe Pro Pro Ala Val Pro Arg Ser Leu Gly Leu Leu Pro Ala Leu
340 345 350

Arg Leu Glu Thr Val Asp Pro Gly Gly Gly Gly Ser Trp Gly Pro Asp
355 360 365

Arg Glu Ala Leu Ala Pro Asn Ser Leu Ser Pro Ser Pro
370 375 380

<210> 110

<211> 382

<212> PRT

<213> Homo sapiens

<220>

<221> SITE

<222> (77)

<223> Xaa equals any of the naturally occurring L-amino acids

<220>

<221> SITE

<222> (78)

<223> Xaa equals any of the naturally occurring L-amino acids

<220>

<221> SITE

<222> (79)

<223> Xaa equals any of the naturally occurring L-amino acids

<220>

<221> SITE

<222> (80)

<223> Xaa equals any of the naturally occurring L-amino acids

<220>

<221> SITE

<222> (81)

<223> Xaa equals any of the naturally occurring L-amino acids

<220>

<221> SITE

<222> (82)

<223> Xaa equals any of the naturally occurring L-amino acids

<220>

<221> SITE

<222> (83)

<223> Xaa equals any of the naturally occurring L-amino acids

<220>
<221> SITE
<222> (84)
<223> Xaa equals any of the naturally occurring L-amino acids

<220>
<221> SITE
<222> (85)
<223> Xaa equals any of the naturally occurring L-amino acids

<220>
<221> SITE
<222> (113)
<223> Xaa equals any of the naturally occurring L-amino acids

<220>
<221> SITE
<222> (127)
<223> Xaa equals any of the naturally occurring L-amino acids

<220>
<221> SITE
<222> (128)
<223> Xaa equals any of the naturally occurring L-amino acids

<220>
<221> SITE
<222> (129)
<223> Xaa equals any of the naturally occurring L-amino acids

<220>
<221> SITE
<222> (130)
<223> Xaa equals any of the naturally occurring L-amino acids

<220>
<221> SITE
<222> (131)
<223> Xaa equals any of the naturally occurring L-amino acids

<220>
<221> SITE
<222> (132)
<223> Xaa equals any of the naturally occurring L-amino acids

<220>
<221> SITE
<222> (133)
<223> Xaa equals any of the naturally occurring L-amino acids

<220>
<221> SITE
<222> (134)
<223> Xaa equals any of the naturally occurring L-amino acids

<220>
<221> SITE
<222> (135)
<223> Xaa equals any of the naturally occurring L-amino acids

<220>
<221> SITE
<222> (136)
<223> Xaa equals any of the naturally occurring L-amino acids

<220>
<221> SITE
<222> (137)
<223> Xaa equals any of the naturally occurring L-amino acids

<220>
<221> SITE
<222> (138)
<223> Xaa equals any of the naturally occurring L-amino acids

<220>
<221> SITE
<222> (139)
<223> Xaa equals any of the naturally occurring L-amino acids

<220>
<221> SITE
<222> (140)
<223> Xaa equals any of the naturally occurring L-amino acids

<220>
<221> SITE
<222> (146)
<223> Xaa equals stop translation

<220>
<221> SITE
<222> (150)
<223> Xaa equals any of the naturally occurring L-amino acids

<220>
<221> SITE
<222> (153)
<223> Xaa equals any of the naturally occurring L-amino acids

<220>
<221> SITE
<222> (155)
<223> Xaa equals any of the naturally occurring L-amino acids

<220>
<221> SITE
<222> (177)
<223> Xaa equals any of the naturally occurring L-amino acids

<220>
<221> SITE
<222> (212)
<223> Xaa equals any of the naturally occurring L-amino acids

<220>
<221> SITE
<222> (229)
<223> Xaa equals any of the naturally occurring L-amino acids

<220>
<221> SITE
<222> (230)
<223> Xaa equals any of the naturally occurring L-amino acids

<220>
<221> SITE
<222> (231)
<223> Xaa equals any of the naturally occurring L-amino acids

<220>
<221> SITE
<222> (232)
<223> Xaa equals any of the naturally occurring L-amino acids

<220>
<221> SITE
<222> (233)
<223> Xaa equals any of the naturally occurring L-amino acids

<220>
<221> SITE
<222> (234)
<223> Xaa equals any of the naturally occurring L-amino acids

<220>
<221> SITE
<222> (235)
<223> Xaa equals any of the naturally occurring L-amino acids

<220>
<221> SITE
<222> (236)
<223> Xaa equals any of the naturally occurring L-amino acids

<220>
<221> SITE
<222> (237)
<223> Xaa equals any of the naturally occurring L-amino acids

<220>
<221> SITE
<222> (238)
<223> Xaa equals any of the naturally occurring L-amino acids

<220>
<221> SITE
<222> (239)
<223> Xaa equals any of the naturally occurring L-amino acids

<220>
<221> SITE
<222> (240)
<223> Xaa equals any of the naturally occurring L-amino acids

<220>
<221> SITE
<222> (241)
<223> Xaa equals any of the naturally occurring L-amino acids

<220>
<221> SITE
<222> (242)
<223> Xaa equals any of the naturally occurring L-amino acids

<220>
<221> SITE
<222> (243)
<223> Xaa equals any of the naturally occurring L-amino acids

<220>
<221> SITE
<222> (244)
<223> Xaa equals any of the naturally occurring L-amino acids

<220>
<221> SITE
<222> (245)
<223> Xaa equals any of the naturally occurring L-amino acids

<220>
<221> SITE
<222> (246)
<223> Xaa equals any of the naturally occurring L-amino acids

<220>
<221> SITE
<222> (247)
<223> Xaa equals any of the naturally occurring L-amino acids

<220>
<221> SITE
<222> (248)
<223> Xaa equals any of the naturally occurring L-amino acids

<220>
<221> SITE
<222> (249)
<223> Xaa equals any of the naturally occurring L-amino acids

<220>
<221> SITE
<222> (250)
<223> Xaa equals any of the naturally occurring L-amino acids

<220>
<221> SITE
<222> (251)
<223> Xaa equals any of the naturally occurring L-amino acids

<220>
<221> SITE
<222> (252)
<223> Xaa equals any of the naturally occurring L-amino acids

<220>
<221> SITE
<222> (253)
<223> Xaa equals any of the naturally occurring L-amino acids

<220>
<221> SITE
<222> (254)
<223> Xaa equals any of the naturally occurring L-amino acids

<220>
<221> SITE
<222> (260)
<223> Xaa equals any of the naturally occurring L-amino acids

<220>
<221> SITE
<222> (261)
<223> Xaa equals any of the naturally occurring L-amino acids

<220>
<221> SITE
<222> (262)
<223> Xaa equals any of the naturally occurring L-amino acids

<220>
<221> SITE
<222> (263)
<223> Xaa equals any of the naturally occurring L-amino acids

<220>
<221> SITE
<222> (264)
<223> Xaa equals any of the naturally occurring L-amino acids

<220>
<221> SITE
<222> (265)
<223> Xaa equals any of the naturally occurring L-amino acids

<220>
<221> SITE
<222> (266)
<223> Xaa equals any of the naturally occurring L-amino acids

<220>
<221> SITE
<222> (267)
<223> Xaa equals any of the naturally occurring L-amino acids

<220>
<221> SITE
<222> (268)
<223> Xaa equals any of the naturally occurring L-amino acids

<220>
<221> SITE
<222> (269)
<223> Xaa equals any of the naturally occurring L-amino acids

<220>
<221> SITE
<222> (270)
<223> Xaa equals any of the naturally occurring L-amino acids

<220>
<221> SITE
<222> (271)
<223> Xaa equals any of the naturally occurring L-amino acids

<220>
<221> SITE
<222> (272)
<223> Xaa equals any of the naturally occurring L-amino acids

<220>
<221> SITE
<222> (273)
<223> Xaa equals any of the naturally occurring L-amino acids

<220>
<221> SITE
<222> (274)
<223> Xaa equals any of the naturally occurring L-amino acids

<220>
<221> SITE
<222> (275)
<223> Xaa equals any of the naturally occurring L-amino acids

<220>
<221> SITE
<222> (276)
<223> Xaa equals any of the naturally occurring L-amino acids

<220>
<221> SITE
<222> (277)
<223> Xaa equals any of the naturally occurring L-amino acids

<220>
<221> SITE
<222> (278)
<223> Xaa equals any of the naturally occurring L-amino acids

<220>
<221> SITE
<222> (279)
<223> Xaa equals any of the naturally occurring L-amino acids

<220>
<221> SITE
<222> (295)
<223> Xaa equals any of the naturally occurring L-amino acids

<220>
<221> SITE
<222> (296)
<223> Xaa equals any of the naturally occurring L-amino acids

<220>
<221> SITE
<222> (297)
<223> Xaa equals any of the naturally occurring L-amino acids

<220>
<221> SITE
<222> (298)
<223> Xaa equals any of the naturally occurring L-amino acids

<220>
<221> SITE
<222> (299)
<223> Xaa equals any of the naturally occurring L-amino acids

<220>
<221> SITE
<222> (300)
<223> Xaa equals any of the naturally occurring L-amino acids

<220>
<221> SITE
<222> (301)
<223> Xaa equals any of the naturally occurring L-amino acids

<220>
<221> SITE
<222> (302)
<223> Xaa equals any of the naturally occurring L-amino acids

<220>
<221> SITE
<222> (303)
<223> Xaa equals any of the naturally occurring L-amino acids

<220>
<221> SITE
<222> (304)
<223> Xaa equals any of the naturally occurring L-amino acids

<220>
<221> SITE
<222> (305)
<223> Xaa equals any of the naturally occurring L-amino acids

<220>
<221> SITE
<222> (306)
<223> Xaa equals any of the naturally occurring L-amino acids

<220>
<221> SITE
<222> (307)
<223> Xaa equals any of the naturally occurring L-amino acids

<220>
<221> SITE
<222> (308)
<223> Xaa equals any of the naturally occurring L-amino acids

<220>
<221> SITE
<222> (309)
<223> Xaa equals any of the naturally occurring L-amino acids

<220>
<221> SITE
<222> (310)
<223> Xaa equals any of the naturally occurring L-amino acids

<220>
<221> SITE
<222> (311)
<223> Xaa equals any of the naturally occurring L-amino acids

<220>
<221> SITE
<222> (316)
<223> Xaa equals any of the naturally occurring L-amino acids

<220>
<221> SITE
<222> (317)
<223> Xaa equals any of the naturally occurring L-amino acids

<220>
<221> SITE
<222> (318)
<223> Xaa equals any of the naturally occurring L-amino acids

<220>
<221> SITE
<222> (319)
<223> Xaa equals any of the naturally occurring L-amino acids

<220>
<221> SITE
<222> (320)
<223> Xaa equals any of the naturally occurring L-amino acids

<220>
<221> SITE
<222> (321)
<223> Xaa equals any of the naturally occurring L-amino acids

<220>
<221> SITE
<222> (322)
<223> Xaa equals any of the naturally occurring L-amino acids

<220>
<221> SITE
<222> (323)
<223> Xaa equals any of the naturally occurring L-amino acids

<220>
<221> SITE
<222> (324)
<223> Xaa equals any of the naturally occurring L-amino acids

<220>
<221> SITE
<222> (325)
<223> Xaa equals any of the naturally occurring L-amino acids

<220>
<221> SITE
<222> (326)
<223> Xaa equals any of the naturally occurring L-amino acids

<220>
<221> SITE
<222> (327)
<223> Xaa equals any of the naturally occurring L-amino acids

<220>
<221> SITE
<222> (328)
<223> Xaa equals any of the naturally occurring L-amino acids

<220>
 <221> SITE
 <222> (329)
 <223> Xaa equals any of the naturally occurring L-amino acids

<220>
 <221> SITE
 <222> (330)
 <223> Xaa equals any of the naturally occurring L-amino acids

<220>
 <221> SITE
 <222> (331)
 <223> Xaa equals any of the naturally occurring L-amino acids

<220>
 <221> SITE
 <222> (332)
 <223> Xaa equals any of the naturally occurring L-amino acids

<220>
 <221> SITE
 <222> (333)
 <223> Xaa equals any of the naturally occurring L-amino acids

<220>
 <221> SITE
 <222> (334)
 <223> Xaa equals any of the naturally occurring L-amino acids

<220>
 <221> SITE
 <222> (335)
 <223> Xaa equals any of the naturally occurring L-amino acids

<400> 110
 Leu His Phe Val Lys Phe Ala Ile Asp Ser Asp Gln Leu His Gln Leu
 1 5 10 15
 Trp Glu Lys Asp Gln Leu Asn Met Ala Lys Asn Thr Trp Pro Ile Leu
 20 25 30
 Lys Gly Phe Gln Glu Gly Pro Leu Asn Phe Ala Pro Thr Phe Lys Phe
 35 40 45
 Asp Val Gly Thr Asn Lys Tyr Asp Thr Ser Ala Lys Lys Arg Lys Pro
 50 55 60
 Ala Trp Thr Asp Arg Ile Leu Trp Lys Val Lys Ala Xaa Xaa Xaa Xaa
 65 70 75 80
 Xaa Xaa Xaa Xaa Xaa Arg Lys Ser His Arg Leu Gln Val Thr Gln His
 85 90 95

Ser Tyr Arg Ser His Met Glu Tyr Thr Val Ser Asp His Lys Pro Val
 100 105 110
 Xaa Ala Gln Phe Leu Leu Gln Phe Ala Phe Gln Gly Arg His Xaa Xaa
 115 120 125
 Xaa Xaa Xaa Xaa Xaa Xaa Xaa Xaa Xaa Xaa Xaa Xaa Pro Ser Arg Arg
 130 135 140
 Trp Xaa Gly Tyr Arg Xaa Glu Thr Xaa Phe Xaa Arg Ser Ser Trp Asp
 145 150 155 160
 Trp Ile Gly Leu Tyr Arg Val Gly Phe Arg His Cys Lys Asp Tyr Val
 165 170 175
 Xaa Tyr Val Trp Ala Lys His Glu Asp Val Asp Gly Asn Thr Tyr Gln
 180 185 190
 Val Thr Phe Ser Glu Glu Ser Leu Pro Lys Gly His Gly Asp Phe Ile
 195 200 205
 Leu Gly Tyr Xaa Ser His Asn His Ser Ile Leu Ile Gly Ile Thr Glu
 210 215 220
 Pro Phe Gln Ile Xaa Xaa Xaa Xaa Xaa Xaa Xaa Xaa Xaa Xaa Xaa Xaa
 225 230 235 240
 Xaa Xaa Xaa Xaa Xaa Xaa Xaa Xaa Xaa Xaa Xaa Xaa Xaa Xaa Leu Glu
 245 250 255
 Leu Leu Ala Xaa Xaa Xaa Xaa Xaa Xaa Xaa Xaa Xaa Xaa Xaa Xaa Xaa Xaa
 260 265 270
 Xaa Xaa Xaa Xaa Xaa Xaa Xaa Leu Ala Arg Phe Pro Gly Leu Ala Leu
 275 280 285
 Arg Pro Ser Ser Arg Glu Xaa Xaa Xaa Xaa Xaa Xaa Xaa Xaa Xaa Xaa
 290 295 300
 Xaa Xaa Xaa Xaa Xaa Xaa Xaa Val Ala Pro Asp Xaa Xaa Xaa Xaa Xaa
 305 310 315 320
 Xaa Xaa Xaa Xaa Xaa Xaa Xaa Xaa Xaa Xaa Xaa Xaa Xaa Xaa Xaa Pro
 325 330 335
 Trp Ala Phe Pro Pro Ala Val Pro Arg Ser Leu Gly Leu Leu Pro Ala
 340 345 350
 Leu Arg Leu Glu Thr Val Asp Pro Gly Gly Gly Gly Ser Trp Gly Pro
 355 360 365
 Asp Arg Glu Ala Leu Ala Pro Asn Ser Leu Ser Pro Ser Pro
 370 375 380

<210> 111
<211> 119
<212> PRT
<213> Homo sapiens

<400> 111
Glu Val Thr Glu Leu Thr Pro Cys Glu Thr Glu Asn Pro Met Gly Gly
1 5 10 15
Tyr Gly Lys Thr Ile Ser His Val Ile Ile Gly Leu Lys Thr Ala Lys
20 25 30
Gly Thr Lys Gln Leu Lys Leu Asp Pro Ser Ile Phe Glu Ser Leu Gln
35 40 45
Lys Glu Arg Val Glu Ala Gly Asp Val Ile Tyr Ile Glu Ala Asn Ser
50 55 60
Gly Ala Val Lys Arg Gln Gly Arg Cys Asp Thr Tyr Ala Thr Glu Phe
65 70 75 80
Asp Leu Glu Ala Glu Glu Tyr Val Pro Leu Pro Lys Gly Asp Val His
85 90 95
Lys Lys Lys Glu Ile Ile Gln Asp Val Thr Leu His Asp Leu Asp Val
100 105 110
Ala Asn Ala Arg Pro Gln Gly
115

<210> 112
<211> 119
<212> PRT
<213> Homo sapiens

<400> 112
Glu Val Thr Glu Leu Thr Pro Cys Glu Thr Glu Asn Pro Met Gly Gly
1 5 10 15
Tyr Gly Lys Thr Ile Ser His Val Ile Ile Gly Leu Lys Thr Ala Lys
20 25 30
Gly Thr Lys Gln Leu Lys Leu Asp Pro Ser Ile Phe Glu Ser Leu Gln
35 40 45
Lys Glu Arg Val Glu Ala Gly Asp Val Ile Tyr Ile Glu Ala Asn Ser
50 55 60
Gly Ala Val Lys Arg Gln Gly Arg Cys Asp Thr Tyr Ala Thr Glu Phe
65 70 75 80
Asp Leu Glu Ala Glu Glu Tyr Val Pro Leu Pro Lys Gly Asp Val His
85 90 95

Lys Lys Lys Glu Ile Ile Gln Asp Val Thr Leu His Asp Leu Asp Val
 100 105 110

Ala Asn Ala Arg Pro Arg Gly
 115

<210> 113
 <211> 121
 <212> PRT
 <213> Homo sapiens

<400> 113
 Met Lys Ile Glu Glu Val Lys Ser Thr Thr Lys Thr Gln Arg Ile Ala
 1 5 10 15

Ser His Ser His Val Lys Gly Leu Gly Leu Asp Glu Ser Gly Leu Ala
 20 25 30

Lys Gln Ala Ala Ser Gly Leu Val Gly Gln Glu Asn Ala Arg Glu Ala
 35 40 45

Cys Gly Val Ile Val Glu Leu Ile Lys Ser Lys Lys Met Ala Gly Arg
 50 55 60

Ala Val Leu Leu Ala Gly Pro Pro Gly Thr Gly Lys Thr Ala Leu Ala
 65 70 75 80

Leu Ala Ile Ala Gln Glu Leu Gly Ser Lys Val Pro Phe Cys Pro Met
 85 90 95

Val Gly Ser Glu Val Tyr Ser Thr Glu Ile Lys Lys Thr Glu Val Leu
 100 105 110

Met Glu Asn Phe Arg Arg Ala Ile Gly
 115 120

<210> 114
 <211> 121
 <212> PRT
 <213> Homo sapiens

<400> 114
 Met Lys Ile Glu Glu Val Lys Ser Thr Thr Lys Thr Gln Arg Ile Ala
 1 5 10 15

Ser His Ser His Val Lys Gly Leu Gly Leu Asp Glu Ser Gly Leu Ala
 20 25 30

Lys Gln Ala Ala Ser Gly Leu Val Gly Gln Glu Asn Ala Arg Glu Ala
 35 40 45

Cys Gly Val Ile Val Glu Leu Ile Lys Ser Lys Lys Met Ala Gly Arg
 50 55 60

Ala Val Leu Leu Ala Gly Pro Pro Gly Thr Gly Lys Thr Ala Leu Ala
 65 70 75 80

Leu Ala Ile Ala Gln Glu Leu Gly Ser Lys Val Pro Phe Cys Pro Met
 85 90 95

Val Gly Ser Glu Val Tyr Ser Thr Glu Ile Lys Lys Thr Glu Val Leu
 100 105 110

Met Glu Asn Phe Arg Arg Pro Leu Gly
 115 120

<210> 115

<211> 128

<212> PRT

<213> Homo sapiens

<400> 115

Ile Leu Ser Met Met Gly Gln Leu Met Lys Pro Lys Lys Thr Glu Ile
 1 5 10 15

Thr Asp Lys Leu Arg Gly Glu Ile Asn Lys Val Val Asn Lys Tyr Ile
 20 25 30

Asp Gln Gly Ile Ala Glu Leu Val Pro Gly Val Leu Phe Val Asp Glu
 35 40 45

Val His Met Leu Asp Ile Glu Cys Phe Thr Tyr Leu His Arg Ala Leu
 50 55 60

Glu Ser Ser Ile Ala Pro Ile Val Ile Phe Ala Ser Asn Arg Gly Asn
 65 70 75 80

Cys Val Ile Arg Gly Thr Glu Asp Ile Thr Ser Pro His Gly Ile Pro
 85 90 95

Leu Asp Leu Leu Asp Arg Val Met Ile Ile Arg Thr Met Leu Tyr Thr
 100 105 110

Pro Gln Glu Met Lys Gln Ile Ile Lys Ile Arg Ala Gln Thr Glu Gly
 115 120 125

<210> 116

<211> 127

<212> PRT

<213> Homo sapiens

<220>

<221> SITE

<222> (97)

<223> Xaa equals any of the naturally occurring L-amino acids

<400> 116

Ile Leu Ser Met Met Gly Gln Leu Met Lys Pro Arg Arg Gln Lys Ser
 1 5 10 15

Arg Gln Thr Ser Gly Glu Ile Asn Lys Val Val Asn Lys Tyr Ile Asp
 20 25 30

Gln Gly Ile Ala Glu Leu Val Pro Gly Val Leu Phe Val Asp Glu Val
 35 40 45

His Met Leu Asp Ile Glu Cys Phe Thr Tyr Leu His Arg Ala Leu Glu
 50 55 60

Ser Ser Ile Ala Pro Ile Val Ile Phe Ala Ser Asn Arg Gly Asn Cys
 65 70 75 80

Val Ile Arg Gly Thr Glu Asp Ile Thr Ser Pro His Gly Ile Pro Leu
 85 90 95

Xaa Leu Leu Asp Arg Val Met Ile Ile Arg Thr Met Leu Tyr Thr Pro
 100 105 110

Gln Glu Met Lys Gln Ile Ile Lys Ile Arg Ala Gln Thr Glu Gly
 115 120 125

<210> 117

<211> 81

<212> PRT

<213> Homo sapiens

<400> 117

Ala Gly Met Gln Trp Cys Asp Leu Ser Ser Leu Gln Pro Pro Pro Pro
 1 5 10 15

Gly Phe Lys Arg Phe Ser His Leu Ser Leu Pro Asn Ser Trp Asn Tyr
 20 25 30

Arg His Leu Pro Ser Cys Pro Thr Asn Phe Cys Ile Phe Val Glu Thr
 35 40 45

Gly Phe His His Val Gly Gln Ala Cys Leu Glu Leu Leu Thr Ser Gly
 50 55 60

Gly Leu Leu Ala Ser Ala Ser Gln Ser Ala Gly Ile Thr Gly Val Ser
 65 70 75 80

His

<210> 118

<211> 81
 <212> PRT
 <213> Homo sapiens

<220>
 <221> SITE
 <222> (20)
 <223> Xaa equals stop translation

<400> 118
 Ala Gly Val Gln Trp Leu Asn Leu Ser Ser Leu Gln Pro Pro His Pro
 1 5 10 15
 Gly Phe Lys Xaa Phe Ser Cys Leu Ser Leu Pro Cys Ser Trp Asp Tyr
 20 25 30
 Arg Ser Ala Pro Pro Gly Pro Gly Asn Phe Cys Ile Phe Arg Arg Asp
 35 40 45
 Gly Val Arg His Val Gly Gln Asn Ser Leu Glu Leu Leu Asn Ser Gly
 50 55 60
 Asp Pro Ala Ala Leu Ala Pro Gln Ser Gly Gly Thr Thr Gly Val Ser
 65 70 75 80
 His

<210> 119
 <211> 43
 <212> PRT
 <213> Homo sapiens

<400> 119
 Phe Cys Ile Phe Val Glu Thr Gly Phe His His Val Gly Gln Ala Cys
 1 5 10 15
 Leu Glu Leu Leu Thr Ser Gly Gly Leu Leu Ala Ser Ala Ser Gln Ser
 20 25 30
 Ala Gly Ile Thr Gly Val Ser His His Ala Arg
 35 40

<210> 120
 <211> 43
 <212> PRT
 <213> Homo sapiens

<400> 120
 Phe Val Phe Val Val Gln Met Gly Leu His His Val Gly Gln Ala Gly
 1 5 10 15
 Leu Lys Leu Pro Thr Ser Gly Asp Pro Pro Ala Ser Ala Ser Gln Ser

20 25 30

Ala Gly Ile Thr Gly Val Ser His His Ala Gln
35 40

<210> 121
<211> 39
<212> PRT
<213> Homo sapiens

<400> 121
Leu Gln Pro Pro Pro Gly Phe Lys Arg Phe Ser His Leu Ser Leu
1 5 10 15
Pro Asn Ser Trp Asn Tyr Arg His Leu Pro Ser Cys Pro Thr Asn Phe
20 25 30
Cys Ile Phe Val Glu Thr Gly
35

<210> 122
<211> 39
<212> PRT
<213> Homo sapiens

<400> 122
Met Gln Pro Pro Pro Gly Leu Lys Gln Phe Ser Arg Leu Ser Pro
1 5 10 15
Pro Ser Ser Trp Asp Tyr Arg Cys Val Pro Pro Arg Leu Ala Asn Phe
20 25 30
Cys Ile Cys Ser Thr Asn Gly
35

<210> 123
<211> 17
<212> PRT
<213> Homo sapiens

<400> 123
Ala Ser Ala Ser Gln Ser Ala Gly Ile Thr Gly Val Ser His His Ala
1 5 10 15
Arg

<210> 124
<211> 17
<212> PRT
<213> Homo sapiens

<400> 124

Ala Ser Ala Ser His Val Ala Gly Thr Thr Gly Val His His Gln Ala
 1 5 10 15

Arg

<210> 125

<211> 255

<212> PRT

<213> Homo sapiens

<400> 125

Asp Tyr Ser Phe Arg Ile Val Asp Thr Asp Ser Gly Arg Val Arg Gln
 1 5 10 15

Ala Val Tyr Gly His Gly Asp Val Val Thr Cys Ile Ala Arg Ser Glu
 20 25 30

Thr Ser Leu Phe Ser Asp Cys Tyr Val Val Thr Gly Ser Met Asp Cys
 35 40 45

Thr Val Val Leu Trp His Trp Asn Gly Thr Thr Gly Phe Ile Ala Gly
 50 55 60

Glu Tyr Asn Gln Pro Gly Glu Val Pro Ser Pro Arg Ser Ile Leu Thr
 65 70 75 80

Gly His Glu Ala Ser Ile Ser Ala Leu Cys Val Ser Ala Glu His Gly
 85 90 95

Leu Val Val Ser Gly Cys Glu Asp Gly Val Ile Leu Ile His Thr Thr
 100 105 110

Ser Ser Asp Leu Leu Arg Arg Ile Arg Gly His Gly Ile Val Thr Gln
 115 120 125

Leu Ser Met Ser Arg Glu Cys Ile Leu Leu Ser Leu Phe Asp Ser Lys
 130 135 140

Arg Met Val Thr Tyr Ser Ala Thr Ala Lys Lys Leu Asp Glu Val Leu
 145 150 155 160

Val Asp Asp Lys Ile Glu Cys Val Thr Val Thr Arg Asp Gly Glu Phe
 165 170 175

Ala Val Thr Gly Ala Val Asn Gly Arg Ile Asn Ile Trp Arg Met Phe
 180 185 190

Pro Leu Thr Lys Leu Tyr Thr Tyr Gln Pro Leu Asn Ser Ala Val Arg
 195 200 205

Ser Val Ala Val Val Ala Ser His Arg Phe Ile Leu Gly Gly Leu Asp

210 215 220
 Ser Gly Ala Ile Val Val Phe Asn Ala Asp Phe Asn Arg Trp His Tyr
 225 230 235 240
 Glu Tyr Lys His Arg Tyr Ile Gln Asn Thr Ser Ala Ala Lys Pro
 245 250 255

<210> 126
 <211> 257
 <212> PRT
 <213> Homo sapiens

<220>
 <221> SITE
 <222> (249)
 <223> Xaa equals stop translation

<220>
 <221> SITE
 <222> (251)
 <223> Xaa equals stop translation

<400> 126
 Asp Lys Ser Phe Arg Val Tyr Ser Thr Glu Thr Gly Lys Leu Thr Gln
 1 5 10 15
 Ile Val Phe Gly His Trp Asp Val Val Thr Cys Leu Ala Arg Ser Glu
 20 25 30
 Ser Tyr Ile Gly Gly Asp Cys Tyr Ile Val Ser Gly Ser Arg Asp Ala
 35 40 45
 Thr Leu Leu Leu Trp Tyr Trp Ser Gly Arg His His Ile Ile Gly Asp
 50 55 60
 Asn Pro Asn Ser Ser Asp Tyr Pro Ala Pro Arg Ala Val Leu Thr Gly
 65 70 75 80
 His Asp His Glu Val Val Cys Val Ser Val Cys Ala Glu Leu Gly Leu
 85 90 95
 Val Ile Ser Gly Ala Lys Glu Gly Pro Cys Leu Val His Thr Ile Thr
 100 105 110
 Gly Asp Leu Leu Arg Ala Leu Glu Gly Pro Glu Asn Cys Leu Phe Pro
 115 120 125
 Arg Leu Ile Ser Val Ser Ser Glu Gly His Cys Ile Ile Tyr Tyr Glu
 130 135 140
 Arg Gly Arg Phe Ser Asn Phe Ser Ile Asn Gly Lys Leu Leu Ala Gln
 145 150 155 160

Met Glu Ile Asn Asp Ser Thr Arg Ala Ile Leu Leu Ser Ser Asp Gly
 165 170 175

Gln Asn Leu Val Thr Gly Gly Asp Asn Gly Val Val Glu Val Trp Gln
 180 185 190

Ala Cys Asp Phe Lys Gln Leu Tyr Ile Tyr Pro Gly Cys Asp Ala Gly
 195 200 205

Ile Arg Ala Met Asp Leu Ser His Asp Gln Arg Thr Leu Ile Thr Gly
 210 215 220

Met Ala Ser Gly Ser Ile Val Ala Phe Asn Ile Asp Phe Asn Arg Trp
 225 230 235 240

His Tyr Glu His Gln Asn Arg Tyr Xaa Arg Xaa Arg Lys Asn Gln Lys
 245 250 255

Pro

<210> 127
 <211> 126
 <212> PRT
 <213> Homo sapiens

<400> 127
 Pro Glu Lys Gln Gln Lys Lys Lys Ser Tyr Lys Asn Ser Gly Thr Ile
 1 5 10 15

Arg Val Lys Ile Cys Arg Val Glu Thr Glu Tyr Ser Phe Leu Asp Tyr
 20 25 30

Val Met Gly Gly Cys Gln Ile Asn Phe Thr Val Gly Val Asp Phe Thr
 35 40 45

Gly Ser Asn Gly Asp Pro Ser Ser Pro Asp Ser Leu His Tyr Leu Ser
 50 55 60

Pro Thr Gly Val Asn Glu Tyr Leu Met Ala Leu Trp Ser Val Gly Ser
 65 70 75 80

Val Val Gln Asp Tyr Asp Ser Asp Lys Leu Phe Pro Ala Phe Gly Phe
 85 90 95

Gly Ala Gln Val Pro Pro Asp Trp Gln Val Ser His Glu Phe Ala Leu
 100 105 110

Asn Phe Asn Pro Ser Asn Pro Tyr Cys Ala Gly Ile Gln Gly
 115 120 125

<210> 128
 <211> 126

<212> PRT

<213> Homo sapiens

<400> 128

```

Pro Lys Tyr Arg Asp Lys Lys Lys Asn Tyr Lys Ser Ser Gly Thr Val
 1             5             10             15
Val Leu Ala Gln Cys Thr Val Glu Lys Val His Thr Phe Leu Asp Tyr
             20             25             30
Ile Met Gly Gly Cys Gln Ile Ser Phe Thr Val Ala Ile Asp Phe Thr
             35             40             45
Ala Ser Asn Gly Asp Pro Arg Ser Ser Gln Ser Leu His Cys Leu Ser
             50             55             60
Pro Arg Gln Pro Asn His Tyr Leu Gln Ala Leu Arg Ala Val Gly Gly
             65             70             75             80
Ile Cys Gln Asp Tyr Asp Ser Asp Lys Arg Phe Pro Ala Phe Gly Phe
             85             90             95
Gly Ala Arg Ile Pro Pro Asn Phe Glu Val Ser His Asp Phe Ala Ile
             100            105            110
Asn Phe Asp Pro Glu Asn Pro Glu Cys Glu Glu Ser Gln Gly
             115            120            125

```

<210> 129

<211> 140

<212> PRT

<213> Homo sapiens

<400> 129

```

Arg Leu Tyr Gly Pro Thr Asn Phe Ala Pro Ile Ile Asn His Val Ala
 1             5             10             15
Arg Phe Ala Ala Gln Ala Ala His Gln Gly Thr Ala Ser Gln Tyr Phe
             20             25             30
Met Leu Leu Leu Leu Thr Asp Gly Ala Val Thr Asp Val Glu Ala Thr
             35             40             45
Arg Glu Ala Val Val Arg Ala Ser Asn Leu Pro Met Ser Val Ile Ile
             50             55             60
Val Gly Val Gly Gly Ala Asp Phe Glu Ala Met Glu Gln Leu Asp Ala
             65             70             75             80
Asp Gly Gly Pro Leu His Thr Arg Ser Gly Gln Ala Ala Ala Arg Asp
             85             90             95
Ile Val Gln Phe Val Pro Tyr Arg Arg Phe Gln Asn Ala Pro Arg Glu
             100            105            110

```

Ala Leu Ala Gln Thr Val Leu Ala Glu Val Pro Thr Gln Leu Val Ser
 115 120 125

Tyr Phe Arg Ala Gln Gly Trp Ala Pro Leu Lys Pro
 130 135 140

<210> 130

<211> 140

<212> PRT

<213> Homo sapiens

<400> 130

Gln Leu Tyr Gly Pro Thr Asn Val Ala Pro Ile Ile Asn Arg Val Ala
 1 5 10 15

Glu Pro Ala Gln Arg Glu Gln Ser Thr Gly Gln Ala Thr Lys Tyr Ser
 20 25 30

Val Leu Leu Val Leu Thr Asp Gly Val Val Ser Asp Met Ala Glu Thr
 35 40 45

Arg Thr Ala Ile Val Arg Ala Ser Arg Leu Pro Met Ser Ile Ile Ile
 50 55 60

Val Gly Val Gly Asn Ala Asp Phe Ser Asp Met Arg Leu Leu Asp Gly
 65 70 75 80

Asp Asp Gly Pro Leu Arg Cys Pro Arg Gly Val Pro Ala Ala Arg Asp
 85 90 95

Ile Val Gln Phe Val Pro Phe Arg Asp Phe Lys Asp Ala Ala Pro Ser
 100 105 110

Ala Leu Ala Lys Cys Val Leu Ala Glu Val Pro Arg Gln Val Val Glu
 115 120 125

Tyr Tyr Ala Ser Gln Gly Ile Ser Pro Gly Ala Pro
 130 135 140

<210> 131

<211> 176

<212> PRT

<213> Homo sapiens

<400> 131

Lys Thr Pro Glu Leu Arg Asp Asp Asp Phe Leu Gly Gly Ala Glu Cys
 1 5 10 15

Ser Leu Gly Gln Ile Val Ser Ser Gln Val Leu Thr Leu Pro Leu Met
 20 25 30

Leu Lys Pro Gly Lys Pro Ala Gly Arg Gly Thr Ile Thr Val Ser Ala

35 40 45
 Gln Glu Leu Lys Asp Asn Arg Val Val Thr Met Glu Val Glu Ala Arg
 50 55 60
 Asn Leu Asp Lys Lys Asp Phe Leu Gly Lys Ser Asp Pro Phe Leu Glu
 65 70 75 80
 Phe Phe Arg Gln Gly Asp Gly Lys Trp His Leu Val Tyr Arg Ser Glu
 85 90 95
 Val Ile Lys Asn Asn Leu Asn Pro Thr Trp Lys Arg Phe Ser Val Pro
 100 105 110
 Val Gln His Phe Cys Gly Gly Asn Pro Ser Thr Pro Ile Gln Val Gln
 115 120 125
 Cys Ser Asp Tyr Asp Ser Asp Gly Ser His Asp Leu Ile Gly Thr Phe
 130 135 140
 His Thr Ser Leu Ala Gln Leu Gln Ala Val Pro Ala Glu Phe Glu Cys
 145 150 155 160
 Ile His Pro Glu Lys Gln Gln Lys Lys Lys Ser Tyr Lys Asn Ser Gly
 165 170 175

<210> 132

<211> 184

<212> PRT

<213> Homo sapiens

<400> 132

Arg Thr Thr Ser Pro Arg Asn Asp Thr Phe Leu Gly Ser Thr Glu Cys
 1 5 10 15
 Thr Leu Gly Gln Ile Val Ser Gln Thr Lys Val Thr Lys Pro Leu Leu
 20 25 30
 Leu Lys Asn Gly Lys Thr Ala Gly Lys Ser Thr Ile Thr Ile Val Ala
 35 40 45
 Glu Glu Val Ser Gly Thr Asn Asp Tyr Val Gln Leu Thr Phe Arg Ala
 50 55 60
 Tyr Lys Leu Asp Asn Lys Asp Leu Phe Ser Lys Ser Asp Pro Phe Met
 65 70 75 80
 Glu Ile Tyr Lys Thr Asn Glu Asp Gln Ser Asp Gln Leu Val Trp Arg
 85 90 95
 Thr Glu Val Val Lys Asn Asn Leu Asn Pro Ser Trp Glu Pro Phe Arg

[illegible]

```
<210> 133
<211> 118
<212> PRT
<213> Homo sapiens
```

```

<400> 133
Gly Ser Val Lys Arg Gly Glu Lys Asp Leu Arg Leu Gly Asp Arg Val
  1             5             10             15
Leu Val Gly Gly Thr Lys Thr Gly Val Val Arg Tyr Val Gly Glu Thr
      20             25             30
Asp Phe Ala Lys Gly Glu Trp Cys Gly Val Glu Leu Asp Glu Pro Leu
      35             40             45
Gly Lys Asn Asp Gly Ala Val Ala Gly Thr Arg Tyr Phe Gln Cys Pro
      50             55             60
Pro Lys Phe Gly Leu Phe Ala Pro Ile His Lys Val Ile Arg Ile Gly
  65             70             75             80
Phe Pro Ser Thr Ser Pro Ala Lys Ala Lys Lys Thr Lys Arg Met Ala
      85             90             95
Met Gly Val Ser Ala Leu Thr His Ser Pro Ser Ser Ser Ser Ile Ser
      100             105             110
Ser Val Ser Ser Val Ala
      115

```

```
<210> 134
<211> 124
<212> PRT
<213> Homo sapiens
```

<400> 134

Gly Ser Leu Gln Gln Arg Asp Gly Ala Lys Ala Glu Val Gly Asp Gln
 1 5 10 15
 Val Leu Val Ala Gly Gln Lys Gln Gly Ile Val Arg Phe Tyr Gly Lys
 20 25 30
 Thr Asp Phe Ala Pro Gly Tyr Trp Tyr Gly Ile Glu Leu Asp Gln Pro
 35 40 45
 Thr Gly Lys His Asp Gly Ser Val Phe Gly Val Arg Tyr Phe Thr Cys
 50 55 60
 Pro Pro Arg His Gly Val Phe Ala Pro Ala Ser Arg Ile Gln Arg Ile
 65 70 75 80
 Gly Gly Ser Thr Asp Ser Pro Gly Asp Ser Val Gly Ala Lys Lys Val
 85 90 95
 His Gln Val Thr Met Thr Gln Pro Lys Arg Thr Phe Thr Thr Val Arg
 100 105 110
 Thr Pro Lys Asp Ile Ala Ser Glu Asn Ser Ile Ser
 115 120

<210> 135
 <211> 41
 <212> PRT
 <213> Homo sapiens

<400> 135
 Gln Arg Leu Ile Tyr Leu Pro Pro Pro Trp Gly Thr Cys Lys Ala Val
 1 5 10 15
 Thr Met Asp Ser Asp Leu Asp Phe Phe Asp Ser Tyr Ser Ile Thr Ala
 20 25 30
 Cys Arg Ile Asp Cys Glu Thr Arg Tyr
 35 40

<210> 136
 <211> 41
 <212> PRT
 <213> Homo sapiens

<220>
 <221> SITE
 <222> (32)
 <223> Xaa equals any of the naturally occurring L-amino acids

<400> 136
 Gln Gln Leu Ile Tyr Leu Pro Pro Pro Trp Gly Thr Cys Lys Ala Val
 1 5 10 15

Thr Met Asp Ser Asp Leu Asp Phe Phe Asp Ser Tyr Ser Ile Thr Xaa
20 25 30

Cys Arg Ile Asp Cys Glu Thr Arg Tyr
35 40

<210> 137

<211> 19

<212> PRT

<213> Homo sapiens

<400> 137

Gly Asp Ala Pro Tyr Cys Thr Pro Glu Gln Tyr Lys Glu Cys Ala Asp
1 5 10 15

Pro Ala Leu

<210> 138

<211> 19

<212> PRT

<213> Homo sapiens

<400> 138

Gly Asp Ala Pro Tyr Cys Thr Pro Glu Gln Tyr Lys Glu Cys Ala Asp
1 5 10 15

Pro Ala Leu

<210> 139

<211> 16

<212> PRT

<213> Homo sapiens

<400> 139

Leu Val Glu Asn Cys Asn Cys Arg Met Val His Met Pro Gly Asp Ala
1 5 10 15

<210> 140

<211> 16

<212> PRT

<213> Homo sapiens

<400> 140

Leu Val Glu Asn Cys Asn Cys Arg Met Val His Met Pro Gly Gln Ala
1 5 10 15

<210> 141
 <211> 63
 <212> PRT
 <213> Homo sapiens

<400> 141
 Gly Pro Leu Ile Asn Thr Pro Leu Asp Cys Glu Val Ser Val Trp Ser
 1 5 10 15
 Pro Trp Gly Leu Cys Lys Gly Gln Cys Gly Glu Lys Gly Val Lys His
 20 25 30
 Arg Thr Arg Tyr Ile His Met His Pro Ala Asn Asn Gly Ala Pro Cys
 35 40 45
 Pro Ser Leu Glu Glu Lys Arg Leu Cys Ile Pro Asp Asn Cys Val
 50 55 60

<210> 142
 <211> 63
 <212> PRT
 <213> Homo sapiens

<400> 142
 Gly Pro Val Pro Glu Thr Pro Leu Asp Cys Glu Val Ser Leu Trp Ser
 1 5 10 15
 Ser Trp Gly Leu Cys Gly Gly His Cys Gly Arg Leu Gly Thr Lys Ser
 20 25 30
 Arg Thr Arg Tyr Val Arg Val Gln Pro Ala Asn Asn Gly Ser Pro Cys
 35 40 45
 Pro Glu Leu Glu Glu Glu Ala Glu Cys Val Pro Asp Asn Cys Val
 50 55 60

<210> 143
 <211> 171
 <212> PRT
 <213> Homo sapiens

<400> 143
 Cys Ser Val Tyr Val Pro Met Cys Thr Glu Lys Ile Asn Ile Pro Ile
 1 5 10 15
 Gly Pro Cys Gly Gly Met Cys Leu Ser Val Lys Arg Arg Cys Glu Pro
 20 25 30
 Val Leu Arg Glu Phe Gly Phe Ala Trp Pro Asp Thr Leu Asn Cys Ser

```

          35              40              45
Lys Phe Pro Pro Gln Asn Asp His Asn His Met Cys Met Glu Gly Pro
   50              55              60
Gly Asp Glu Glu Val Pro Leu Pro His Lys Thr Pro Ile Gln Pro Gly
   65              70              75              80
Glu Glu Cys His Ser Val Gly Ser Asn Ser Asp Gln Tyr Ile Trp Val
          85              90              95
Lys Arg Ser Leu Asn Cys Val Leu Lys Cys Gly Tyr Asp Ala Gly Leu
          100              105              110
Tyr Ser Arg Ser Ala Lys Glu Phe Thr Asp Ile Trp Met Ala Val Trp
          115              120              125
Ala Ser Leu Cys Phe Ile Ser Thr Thr Phe Thr Val Leu Thr Phe Leu
          130              135              140
Ile Asp Ser Ser Arg Phe Ser Tyr Pro Glu Arg Pro Ile Ile Phe Leu
          145              150              155              160
Ser Met Cys Tyr Asn Ile Tyr Ser Ile Ala Tyr
          165              170

```

<210> 144
 <211> 171
 <212> PRT
 <213> Homo sapiens

```

<400> 144
Cys Ser Val Tyr Val Pro Met Cys Thr Glu Lys Ile Asn Ile Pro Ile
   1              5              10              15
Gly Pro Cys Gly Gly Met Cys Leu Ser Val Lys Arg Arg Cys Glu Pro
          20              25              30
Val Leu Lys Glu Phe Gly Phe Ala Trp Pro Glu Ser Leu Asn Cys Ser
          35              40              45
Lys Phe Pro Pro Gln Asn Asp His Asn His Met Cys Met Glu Gly Pro
   50              55              60
Gly Asp Glu Glu Val Pro Leu Pro His Lys Thr Pro Ile Gln Pro Gly
   65              70              75              80
Glu Glu Cys His Ser Val Gly Thr Asn Ser Asp Gln Tyr Ile Trp Val
          85              90              95
Lys Arg Ser Leu Asn Cys Val Leu Lys Cys Gly Tyr Asp Ala Gly Leu
          100              105              110
Tyr Ser Arg Ser Ala Lys Glu Phe Thr Asp Ile Trp Met Ala Val Trp

```

115 120 125
 Ala Ser Leu Cys Phe Ile Ser Thr Ala Phe Thr Val Leu Thr Phe Leu
 130 135 140
 Ile Asp Ser Ser Arg Phe Ser Tyr Pro Glu Arg Pro Ile Ile Phe Leu
 145 150 155 160
 Ser Met Cys Tyr Asn Ile Tyr Ser Ile Ala Tyr
 165 170

<210> 145
 <211> 176
 <212> PRT
 <213> Homo sapiens

<400> 145
 Phe Leu Ser Met Cys Tyr Asn Ile Tyr Ser Ile Ala Tyr Ile Val Arg
 1 5 10 15
 Leu Thr Val Gly Arg Glu Arg Ile Ser Cys Asp Phe Glu Glu Ala Ala
 20 25 30
 Glu Pro Val Leu Ile Gln Glu Gly Leu Lys Asn Thr Gly Cys Ala Ile
 35 40 45
 Ile Phe Leu Leu Met Tyr Phe Phe Gly Met Ala Ser Ser Ile Trp Trp
 50 55 60
 Val Ile Leu Thr Leu Thr Trp Phe Leu Ala Ala Gly Leu Lys Trp Gly
 65 70 75 80
 His Glu Ala Ile Glu Met His Ser Ser Tyr Phe His Ile Ala Ala Trp
 85 90 95
 Ala Ile Pro Ala Val Lys Thr Ile Val Ile Leu Ile Met Arg Leu Val
 100 105 110
 Asp Ala Asp Glu Leu Thr Gly Leu Cys Tyr Val Gly Asn Gln Asn Leu
 115 120 125
 Asp Ala Leu Thr Gly Phe Val Val Ala Pro Leu Phe Thr Tyr Leu Val
 130 135 140
 Ile Gly Thr Leu Phe Ile Ala Ala Gly Leu Val Ala Leu Phe Lys Ile
 145 150 155 160
 Arg Ser Asn Leu Gln Lys Asp Gly Thr Lys Thr Asp Lys Leu Glu Arg
 165 170 175

<210> 146
<211> 177
<212> PRT
<213> Homo sapiens

<220>
<221> SITE
<222> (147)
<223> Xaa equals any of the naturally occurring L-amino acids

<220>
<221> SITE
<222> (148)
<223> Xaa equals any of the naturally occurring L-amino acids

<220>
<221> SITE
<222> (149)
<223> Xaa equals any of the naturally occurring L-amino acids

<220>
<221> SITE
<222> (150)
<223> Xaa equals any of the naturally occurring L-amino acids

<220>
<221> SITE
<222> (151)
<223> Xaa equals any of the naturally occurring L-amino acids

<220>
<221> SITE
<222> (152)
<223> Xaa equals any of the naturally occurring L-amino acids

<220>
<221> SITE
<222> (153)
<223> Xaa equals any of the naturally occurring L-amino acids

<220>
<221> SITE
<222> (154)
<223> Xaa equals any of the naturally occurring L-amino acids

<220>
<221> SITE
<222> (155)
<223> Xaa equals any of the naturally occurring L-amino acids

<220>
<221> SITE
<222> (156)
<223> Xaa equals any of the naturally occurring L-amino acids

<220>
 <221> SITE
 <222> (157)
 <223> Xaa equals any of the naturally occurring L-amino acids

<220>
 <221> SITE
 <222> (158)
 <223> Xaa equals any of the naturally occurring L-amino acids

<220>
 <221> SITE
 <222> (159)
 <223> Xaa equals any of the naturally occurring L-amino acids

<220>
 <221> SITE
 <222> (160)
 <223> Xaa equals any of the naturally occurring L-amino acids

<220>
 <221> SITE
 <222> (161)
 <223> Xaa equals any of the naturally occurring L-amino acids

<220>
 <221> SITE
 <222> (166)
 <223> Xaa equals any of the naturally occurring L-amino acids

<400> 146
 Tyr Phe Ser Val Cys Ala Ile Ile Phe Ile Ala Leu Leu Ile Val Arg
 1 5 10 15
 Leu Thr Val Gly Arg Glu Arg Ile Ser Cys Asp Phe Glu Glu Ala Ala
 20 25 30
 Glu Pro Val Leu Ile Gln Glu Gly Leu Lys Asn Thr Gly Cys Ala Ile
 35 40 45
 Ile Phe Leu Leu Met Tyr Phe Phe Gly Met Ala Ser Ser Ile Trp Trp
 50 55 60
 Val Ile Leu Thr Leu Thr Trp Phe Leu Ala Ala Gly Leu Lys Trp Gly
 65 70 75 80
 His Glu Ala Ile Glu Met His Ser Ser Tyr Phe His Ile Ala Ala Trp
 85 90 95
 Ala Ile Pro Ala Val Lys Thr Ile Val Ile Leu Ile Met Arg Leu Val
 100 105 110
 Asp Ala Asp Glu Leu Thr Gly Leu Cys Tyr Val Gly Asn Gln Asn Leu
 115 120 125

Asp Ala Leu Thr Gly Phe Val Val Ala Pro Leu Phe Thr Tyr Leu Ala
 130 135 140

Leu Glu Xaa Xaa Xaa Xaa Xaa Xaa Xaa Xaa Xaa Xaa Xaa Xaa Xaa
 145 150 155 160

Xaa Lys Ile Phe Lys Xaa Asp Gly Thr Lys Asp Gln Thr Lys Leu Glu
 165 170 175

Lys

<210> 147

<211> 255

<212> PRT

<213> Homo sapiens

<400> 147

Leu Leu Ser Glu Pro Val Val Thr Val Ser Ser Asn Leu Lys Ser Ile
 1 5 10 15

Ser Tyr Gln Val Tyr Ile Arg Tyr Ser Trp Leu Ser Leu Lys Ala Ile
 20 25 30

Phe Lys Thr Ser Lys Thr Gln His Leu Thr Asp Tyr Asn Arg Lys Arg
 35 40 45

Asn Gly Leu Ser Leu Ile Ser Phe Arg Met Thr Gln Ile Val Phe Cys
 50 55 60

Phe Gly Cys Ser Cys Leu Val Gln Ser Tyr Ile Thr Glu Asn Tyr Phe
 65 70 75 80

Leu Arg Asn Leu Lys Thr Arg Met Phe Lys Thr Asn Ser Ile Ile Glu
 85 90 95

Val Ser Arg Tyr Val Glu His Leu Arg Asp Phe Ser Ser Gly Thr Ser
 100 105 110

Thr Asn Phe Arg Ile Leu Val Ser Leu Phe Gly Ile His Thr Thr Ser
 115 120 125

Arg Lys Ile Gln Val Phe Cys Phe Val Leu Phe Cys Phe Gly Phe Ile
 130 135 140

Glu Tyr Leu Lys Leu Lys Leu Phe Cys Pro Lys Asn Lys Ile Lys Asp
 145 150 155 160

Phe Ile Lys Leu Lys Leu Ala Pro Thr Glu Leu Lys Tyr Ile Lys Cys
 165 170 175

Ile Lys Tyr Asn Ser Ala Tyr Gly Ser Asn Ile Ala Leu Ile Trp Lys
 180 185 190

Ser Leu Pro Glu Thr Val Tyr Phe Leu Leu Ile Cys Tyr Tyr Leu Val
195 200 205

Thr Thr Pro His Tyr Ser Ser Leu Glu Leu Leu Ile Lys Glu Asn Trp
210 215 220

Lys Tyr Ile Cys Pro Cys Leu Asn Phe Ile Ala Leu Ile Cys Val Ile
225 230 235 240

Ser Leu Leu Thr Gly Arg Gly Thr Ser Phe Phe Pro Tyr Ser Lys
245 250 255

<210> 148

<211> 252

<212> PRT

<213> Homo sapiens

<220>

<221> SITE

<222> (3)

<223> Xaa equals stop translation

<220>

<221> SITE

<222> (18)

<223> Xaa equals stop translation

<220>

<221> SITE

<222> (28)

<223> Xaa equals stop translation

<220>

<221> SITE

<222> (36)

<223> Xaa equals stop translation

<220>

<221> SITE

<222> (53)

<223> Xaa equals stop translation

<220>

<221> SITE

<222> (58)

<223> Xaa equals stop translation

<220>

<221> SITE

<222> (81)

<223> Xaa equals any of the naturally occurring L-amino acids

<220>

<221> SITE

<222> (82)
<223> Xaa equals any of the naturally occurring L-amino acids

<220>
<221> SITE
<222> (83)
<223> Xaa equals any of the naturally occurring L-amino acids

<220>
<221> SITE
<222> (84)
<223> Xaa equals any of the naturally occurring L-amino acids

<220>
<221> SITE
<222> (85)
<223> Xaa equals any of the naturally occurring L-amino acids

<220>
<221> SITE
<222> (86)
<223> Xaa equals any of the naturally occurring L-amino acids

<220>
<221> SITE
<222> (87)
<223> Xaa equals any of the naturally occurring L-amino acids

<220>
<221> SITE
<222> (88)
<223> Xaa equals any of the naturally occurring L-amino acids

<220>
<221> SITE
<222> (89)
<223> Xaa equals any of the naturally occurring L-amino acids

<220>
<221> SITE
<222> (90)
<223> Xaa equals any of the naturally occurring L-amino acids

<220>
<221> SITE
<222> (91)
<223> Xaa equals any of the naturally occurring L-amino acids

<220>
<221> SITE
<222> (92)
<223> Xaa equals any of the naturally occurring L-amino acids

<220>
<221> SITE

<222> (93)
<223> Xaa equals any of the naturally occurring L-amino acids

<220>
<221> SITE
<222> (94)
<223> Xaa equals any of the naturally occurring L-amino acids

<220>
<221> SITE
<222> (95)
<223> Xaa equals any of the naturally occurring L-amino acids

<220>
<221> SITE
<222> (96)
<223> Xaa equals any of the naturally occurring L-amino acids

<220>
<221> SITE
<222> (160)
<223> Xaa equals any of the naturally occurring L-amino acids

<220>
<221> SITE
<222> (161)
<223> Xaa equals any of the naturally occurring L-amino acids

<220>
<221> SITE
<222> (162)
<223> Xaa equals any of the naturally occurring L-amino acids

<220>
<221> SITE
<222> (163)
<223> Xaa equals any of the naturally occurring L-amino acids

<220>
<221> SITE
<222> (164)
<223> Xaa equals any of the naturally occurring L-amino acids

<220>
<221> SITE
<222> (165)
<223> Xaa equals any of the naturally occurring L-amino acids

<220>
<221> SITE
<222> (166)
<223> Xaa equals any of the naturally occurring L-amino acids

<220>
<221> SITE

<222> (167)
<223> Xaa equals any of the naturally occurring L-amino acids

<220>
<221> SITE
<222> (168)
<223> Xaa equals any of the naturally occurring L-amino acids

<220>
<221> SITE
<222> (169)
<223> Xaa equals any of the naturally occurring L-amino acids

<220>
<221> SITE
<222> (170)
<223> Xaa equals any of the naturally occurring L-amino acids

<220>
<221> SITE
<222> (171)
<223> Xaa equals any of the naturally occurring L-amino acids

<220>
<221> SITE
<222> (172)
<223> Xaa equals any of the naturally occurring L-amino acids

<220>
<221> SITE
<222> (173)
<223> Xaa equals any of the naturally occurring L-amino acids

<220>
<221> SITE
<222> (174)
<223> Xaa equals any of the naturally occurring L-amino acids

<220>
<221> SITE
<222> (175)
<223> Xaa equals any of the naturally occurring L-amino acids

<220>
<221> SITE
<222> (218)
<223> Xaa equals stop translation

<400> 148
Val Leu Xaa Glu Pro Val Val Thr Val Ser Ser Asn Leu Lys Ser Ile
1 5 10 15
Ser Xaa Gln Val Tyr Ile Arg Cys Ser Trp Leu Xaa Leu Lys Ala Ile
20 25 30

Phe Lys Thr Xaa Lys Leu Asn Thr Ser Gln Ile Ile Ile Glu Lys Glu
 35 40 45
 Met Ala Ser Val Xaa Ser Arg Ser Glu Xaa Pro Arg Leu Phe Leu Leu
 50 55 60
 Trp Val Gln Leu Phe Ser Ser Glu Leu Tyr Tyr Arg Glu Leu Phe Ser
 65 70 75 80
 Xaa Xaa Xaa Xaa Xaa Xaa Xaa Xaa Xaa Xaa Xaa Xaa Xaa Xaa Xaa Xaa
 85 90 95
 Ser Arg Tyr Val Glu His Leu Arg Asp Phe Ser Ser Gly Thr Ser Thr
 100 105 110
 Asn Phe Glu Ser Leu Tyr Leu Tyr Leu Val Phe Ile Leu Leu Val Ala
 115 120 125
 Lys Tyr Arg Phe Phe Val Leu Phe Cys Phe Gly Phe Ile Glu Tyr Leu
 130 135 140
 Lys Leu Lys Leu Phe Cys Thr Lys Asn Lys Ile Lys Asp Phe Ile Xaa
 145 150 155 160
 Xaa Xaa Xaa Xaa Xaa Xaa Xaa Xaa Xaa Xaa Xaa Xaa Xaa Xaa Xaa Ile
 165 170 175
 Ile Gln His Met Gly Ser Asn Ile Ala Leu Ile Trp Lys Ser Leu Pro
 180 185 190
 Glu Thr Val Tyr Phe Leu Leu Ile Cys Tyr Tyr Leu Val Thr Asn Pro
 195 200 205
 Thr Leu Phe Gln Phe Gly Ile Thr Tyr Xaa Gly Glu Leu Glu Ile His
 210 215 220
 Met Pro Met Leu Lys Phe Tyr Ser Phe Asn Leu Cys Tyr Phe Phe Ile
 225 230 235 240
 Asp Gly Lys Arg Tyr Ile Phe Phe Ser Leu Leu Lys
 245 250

INTERNATIONAL SEARCH REPORT

International application No.

PCT/US00/06014

A. CLASSIFICATION OF SUBJECT MATTER

IPC(7) : C07K 14/47; C12N 5/10, 5/16, 15/12, 15/63, 15/64

US CL : Please See Extra Sheet.

According to International Patent Classification (IPC) or to both national classification and IPC

B. FIELDS SEARCHED

Minimum documentation searched (classification system followed by classification symbols)

U.S. : 530/350; 536/23.1, 23.5, 24.3, 24.31; 435/69.1, 71.1, 71.2, 471, 325, 252.3, 254.11, 320.1

Documentation searched other than minimum documentation to the extent that such documents are included in the fields searched
NONEElectronic data base consulted during the international search (name of data base and, where practicable, search terms used)
WEST, CAS ONLINE, MEDLINE, CAPLUS

C. DOCUMENTS CONSIDERED TO BE RELEVANT

Category*	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
A	WO 96/09386 A2 (THE GOVERNMENT OF THE UNITED STATES OF AMERICA) 28 March 1996 (28/03/1996), see entire document.	1-12, 14-16, 21

☐ Further documents are listed in the continuation of Box C.
 ☐ See patent family annex.

* Special categories of cited documents:	
A document defining the general state of the art which is not considered to be of particular relevance	*T* later document published after the international filing date or priority date and not in conflict with the application but cited to understand the principle or theory underlying the invention
E earlier document published on or after the international filing date	*X* document of particular relevance; the claimed invention cannot be considered novel or cannot be considered to involve an inventive step when the document is taken alone
L document which may throw doubts on priority claim(s) or which is cited to establish the publication date of another citation or other special reason (as specified)	*Y* document of particular relevance; the claimed invention cannot be considered to involve an inventive step when the document is combined with one or more other such documents, such combination being obvious to a person skilled in the art
O document referring to an oral disclosure, use, exhibition or other means	*A* document member of the same patent family
P document published prior to the international filing date but later than the priority date claimed	

Date of the actual completion of the international search

23 JUNE 2000

Date of mailing of the international search report

02 AUG 2000

 Name and mailing address of the ISA/US
 Commissioner of Patents and Trademarks
 Box PCT
 Washington, D.C. 20231

Facsimile No. (703) 305-3230

Authorized officer

PREMA MERTZ

Telephone No. (703) 308-0196

INTERNATIONAL SEARCH REPORT

International application No.
PCT/US00/06014

Box I Observations where certain claims were found unsearchable (Continuation of item 1 of first sheet)

This international report has not been established in respect of certain claims under Article 17(2)(a) for the following reasons:

1. ☐ Claims Nos.:
because they relate to subject matter not required to be searched by this Authority, namely:
2. ☐ Claims Nos.:
because they relate to parts of the international application that do not comply with the prescribed requirements to such an extent that no meaningful international search can be carried out, specifically:
3. ☐ Claims Nos.:
because they are dependent claims and are not drafted in accordance with the second and third sentences of Rule 6.4(a).

Box II Observations where unity of invention is lacking (Continuation of item 2 of first sheet)

This International Searching Authority found multiple inventions in this international application, as follows:

Please See Extra Sheet.

1. ☐ As all required additional search fees were timely paid by the applicant, this international search report covers all searchable claims.
2. ☐ As all searchable claims could be searched without effort justifying an additional fee, this Authority did not invite payment of any additional fee.
3. ☐ As only some of the required additional search fees were timely paid by the applicant, this international search report covers only those claims for which fees were paid, specifically claims Nos.:
4. ☒ No required additional search fees were timely paid by the applicant. Consequently, this international search report is restricted to the invention first mentioned in the claims; it is covered by claims Nos.:
1-12, 14-16, 21 (SEQ ID NO:11, 58)

Remark on Protest

- ☐ The additional search fees were accompanied by the applicant's protest.
☐ No protest accompanied the payment of additional search fees.

INTERNATIONAL SEARCH REPORT

International application No.
PCT/US00/06014

A. CLASSIFICATION OF SUBJECT MATTER:

US CL :

530/350; 536/23.1, 23.5, 24.3, 24.31; 435/69.1, 71.1, 71.2, 471, 325, 252.3, 254.11, 320.1

BOX II. OBSERVATIONS WHERE UNITY OF INVENTION WAS LACKING

This ISA found multiple inventions as follows:

This application contains the following inventions or groups of inventions which are not so linked as to form a single inventive concept under PCT Rule 13.1. In order for all inventions to be searched, the appropriate additional search fees must be paid.

Group I, claims 1-12, 14-16, 21, drawn to a nucleic acid of SEQ ID NO:11 encoding a protein of SEQ ID NO:58, a vector, a host cell, a method of making the protein and the protein of SEQ ID NO:58.

Group II, claim 13, drawn to an antibody that binds the protein of SEQ ID NO:58.

Group III, claim 17, drawn to a method of treating a condition comprising administering the protein of SEQ ID NO:58.

Group IV, claim 18, drawn to a method of diagnosing a pathological condition using the polynucleotide encoding a protein of SEQ ID NO:58.

Group V, claim 19, drawn to a method of diagnosing a pathological condition by determining the amount of protein of SEQ ID NO:58.

Group VI, claim 20, drawn to a method of identifying a binding partner of the protein of SEQ ID NO:58.

Group VII, claims 22-23, drawn to a method of identifying an activity in a biological assay.

Group VIII, claim 17, drawn to a method for treating a condition comprising administering the nucleic acid of SEQ ID NO:11.

The inventions listed as Groups I-VIII do not relate to a single inventive concept under PCT Rule 13.1 because, under PCT Rule 13.2, they lack the same or corresponding special technical features for the following reasons:

Pursuant to 37 C.F.R. § 1.475 (d), the ISA/US considers that where multiple products and processes are claimed, the main invention shall consist of the first invention of the category first mentioned in the claims and the first recited invention of each of the other categories related thereto. Accordingly, the main invention (Group I) comprises the first-recited product, a nucleic acid encoding a protein of SEQ ID NO:58, a vector, a host cell, a method of making the protein of SEQ ID NO:58, and the protein of SEQ ID NO:58. Further pursuant to 37 C.F.R. § 1.475 (d), the ISA/US considers that any feature which the subsequently recited products and methods share with the main invention does not constitute a special technical feature within the meaning of PCT Rule 13.2 and that each of such products and methods accordingly defines a separate invention.

This application contains claims directed to more than one species of the generic invention. These species are deemed to lack Unity of Invention because they are not so linked as to form a single inventive concept under PCT Rule 13.1. In order for more than one species to be searched, the appropriate additional search fees must be paid. The species are as follows:

The polynucleotides set forth in SEQ ID NO:11-57 encoding the polypeptides set forth in SEQ ID NO:58-104.